



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:50 AM GMT

PDB ID : 2EJF
Title : Crystal Structure Of The Biotin Protein Ligase (Mutations R48A and K111A) and Biotin Carboxyl Carrier Protein Complex From Pyrococcus Horikoshii OT3
Authors : Bagautdinov, B.; Matsuura, Y.; Bagautdinova, S.; Kunishima, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2007-03-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	<div> <div>2%</div> <div>81% 18%</div> </div>
1	B	235	<div> <div>6%</div> <div>73% 22%</div> </div>
2	C	74	<div> <div>15%</div> <div>69% 23% 7%</div> </div>
2	D	74	<div> <div>0%</div> <div>78% 15% 7%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5131 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 235aa long hypothetical biotin--[acetyl-CoA-carboxylase] ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	235	Total	C	N	O	S	0	0	0
			1827	1179	307	336	5			
1	B	227	Total	C	N	O	S	0	0	0
			1766	1141	296	324	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	ALA	ARG	ENGINEERED	UNP O57883
A	111	ALA	LYS	ENGINEERED	UNP O57883
B	48	ALA	ARG	ENGINEERED	UNP O57883
B	111	ALA	LYS	ENGINEERED	UNP O57883


- Molecule 2 is a protein called 149aa long hypothetical methylmalonyl-CoA decarboxylase gamma chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	69	Total	C	N	O	S	0	0	0
			519	326	95	95	3			
2	D	69	Total	C	N	O	S	0	0	0
			519	326	95	95	3			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	76	MET	-	INITIATING METHIONINE	UNP O59021
D	76	MET	-	INITIATING METHIONINE	UNP O59021

- Molecule 3 is BIOTIN (three-letter code: BTN) (formula: C₁₀H₁₆N₂O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	S	0	0
			15	10	2	2	1		
3	A	1	Total	C	N	O	S	0	0
			16	10	2	3	1		
3	B	1	Total	C	N	O	S	0	0
			16	10	2	3	1		

- Molecule 4 is ADENOSINE (three-letter code: ADN) (formula: C₁₀H₁₃N₅O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			19	10	5	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			19	10	5	4		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	180	Total	O	0	0
			180	180		
6	B	153	Total	O	0	0
			153	153		
6	C	42	Total	O	0	0
			42	42		
6	D	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 235aa long hypothetical biotin--[acetyl-CoA-carboxylase] ligase




- Molecule 1: 235aa long hypothetical biotin--[acetyl-CoA-carboxylase] ligase



- Molecule 2: 149aa long hypothetical methylmalonyl-CoA decarboxylase gamma chain



- Molecule 2: 149aa long hypothetical methylmalonyl-CoA decarboxylase gamma chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.85Å 63.12Å 75.64Å 90.00° 95.86° 90.00°	Depositor
Resolution (Å)	37.62 – 2.00 40.97 – 2.01	Depositor EDS
% Data completeness (in resolution range)	95.6 (37.62-2.00) 95.6 (40.97-2.01)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.01Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.209 , 0.245 0.219 , 0.255	Depositor DCC
R_{free} test set	2090 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	22.9	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 61.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 42137 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5131	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADN, GOL, BTN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1855	0.64	0/2504
1	B	0.36	0/1792	0.61	0/2420
2	C	0.40	0/523	0.71	0/704
2	D	0.28	0/523	0.62	0/704
All	All	0.37	0/4693	0.63	0/6332

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1827	0	1918	37	0
1	B	1766	0	1858	53	0
2	C	519	0	559	13	0
2	D	519	0	561	9	0
3	A	16	0	15	0	0
3	B	16	0	15	0	0
3	C	15	0	15	0	0
4	A	19	0	13	0	0
4	B	19	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	6	0	8	0	0
6	A	180	0	0	10	0
6	B	153	0	0	9	0
6	C	42	0	0	3	0
6	D	34	0	0	0	0
All	All	5131	0	4975	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:69:LYS:HE2	6:B:3079:HOH:O	1.77	0.83
1:A:24:GLU:HG2	1:A:46:HIS:HE1	1.47	0.79
1:A:11:ARG:HD3	6:A:2120:HOH:O	1.84	0.79
1:B:208:ASP:OD2	1:B:224:LYS:HE3	1.83	0.78
1:A:29:SER:O	1:A:123:LYS:NZ	2.20	0.75
1:B:68:PRO:HD2	1:B:124:ILE:HD13	1.69	0.75
1:A:108:ASN:HB3	6:A:2079:HOH:O	1.88	0.74
1:B:226:ILE:HG23	1:B:227:TYR:HD1	1.54	0.73
2:C:104:VAL:HG23	2:C:125:ASP:OD1	1.90	0.71
1:B:51:ARG:HD2	6:B:3150:HOH:O	1.92	0.70
1:A:51:ARG:NH1	6:A:2128:HOH:O	2.23	0.70
1:B:135:LYS:HE3	6:B:3124:HOH:O	1.91	0.70
1:A:186:MET:HG3	1:A:188:LEU:HG	1.75	0.69
1:B:154:LEU:HD12	1:B:158:ARG:CZ	2.22	0.69
1:B:226:ILE:HG23	1:B:227:TYR:CD1	2.27	0.68
1:A:32:GLU:HG3	1:A:35:THR:OG1	1.94	0.68
1:A:24:GLU:HG2	1:A:46:HIS:CE1	2.30	0.67
2:C:108:LEU:HD21	2:C:122:SER:HB2	1.78	0.66
1:A:193:LYS:HB2	1:A:235:LEU:HD11	1.77	0.65
1:B:154:LEU:HD12	1:B:158:ARG:NH1	2.12	0.64
1:A:205:GLU:O	1:A:206:ASP:HB2	1.97	0.63
1:B:178:ILE:O	1:B:182:VAL:HG23	1.98	0.63
1:B:18:GLU:OE2	1:B:41:LYS:HE3	1.99	0.62
1:B:176:MET:CE	1:B:209:ASP:HB2	2.29	0.61
1:A:193:LYS:HB2	1:A:235:LEU:CD1	2.31	0.61
1:A:188:LEU:HD12	6:A:2094:HOH:O	1.98	0.61
1:B:205:GLU:HG3	1:B:216:ARG:HG2	1.83	0.60
1:A:52:LYS:HG3	6:A:2053:HOH:O	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:GLU:HG3	6:A:2088:HOH:O	2.00	0.60
2:D:90:LYS:HE3	2:D:136:GLY:HA2	1.82	0.59
1:B:190:VAL:HG22	1:B:191:ARG:N	2.17	0.59
1:A:232:LEU:HD23	1:A:232:LEU:C	2.24	0.58
1:B:81:LEU:HD22	1:B:167:LEU:HB3	1.86	0.58
2:C:101:ARG:HG3	2:C:127:VAL:HG22	1.85	0.57
1:A:81:LEU:HD23	1:A:178:ILE:HD13	1.87	0.57
1:B:216:ARG:NH1	1:B:222:VAL:HG22	2.21	0.56
1:B:194:ILE:CD1	1:B:230:VAL:HG22	2.34	0.56
1:B:140:ALA:HB2	6:B:3099:HOH:O	2.04	0.56
1:A:42:GLN:HB2	1:A:55:SER:HB2	1.87	0.56
1:B:204:ALA:HA	1:B:215:ILE:HD13	1.86	0.55
2:C:98:VAL:HG23	6:C:1429:HOH:O	2.07	0.55
1:B:216:ARG:HG2	1:B:216:ARG:HH11	1.72	0.54
1:B:176:MET:HE3	1:B:209:ASP:HB2	1.89	0.54
1:B:124:ILE:N	1:B:124:ILE:HD12	2.24	0.53
1:A:232:LEU:HD23	1:A:233:ARG:N	2.23	0.53
2:D:92:LEU:O	2:D:92:LEU:HD23	2.08	0.53
1:A:56:PRO:HG2	1:A:134:ASN:HB2	1.91	0.52
1:A:49:LEU:HD12	1:A:51:ARG:HH22	1.74	0.52
1:B:176:MET:HE2	1:B:209:ASP:HB2	1.92	0.52
1:B:8:ILE:HG12	6:B:3042:HOH:O	2.10	0.51
2:D:131:ILE:HA	2:D:146:ILE:HD13	1.91	0.51
1:A:75:LEU:N	1:A:76:PRO:CD	2.74	0.51
1:B:81:LEU:CD2	1:B:167:LEU:HB3	2.41	0.50
1:A:32:GLU:HG3	1:A:35:THR:HG1	1.76	0.50
1:B:193:LYS:HA	1:B:201:GLU:HG2	1.94	0.50
2:D:92:LEU:HD22	2:D:93:ARG:NH1	2.28	0.49
1:B:206:ASP:CG	1:B:207:ILE:H	2.16	0.49
1:B:84:VAL:O	1:B:88:GLU:HG3	2.13	0.49
1:A:186:MET:SD	1:A:188:LEU:HD21	2.53	0.48
1:B:202:GLY:HA3	1:B:216:ARG:O	2.14	0.48
1:B:191:ARG:HG3	1:B:191:ARG:HH11	1.78	0.48
1:B:103:ASN:OD1	2:D:115:LYS:HD2	2.13	0.47
1:B:41:LYS:NZ	6:B:3066:HOH:O	2.46	0.47
1:B:195:LEU:O	1:B:195:LEU:HD13	2.14	0.47
1:B:205:GLU:HG3	6:B:3132:HOH:O	2.12	0.47
1:B:46:HIS:NE2	6:B:3148:HOH:O	2.31	0.47
1:A:158:ARG:HB2	1:A:158:ARG:NH1	2.29	0.47
2:D:92:LEU:HD22	2:D:93:ARG:HG2	1.96	0.47
2:C:91:VAL:HA	2:C:111:LEU:HD23	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:TRP:HA	1:A:102:PRO:HA	1.66	0.46
1:B:81:LEU:HD23	1:B:178:ILE:HD13	1.98	0.45
1:B:205:GLU:HG3	1:B:216:ARG:CG	2.46	0.45
1:A:42:GLN:HB2	1:A:55:SER:CB	2.46	0.45
1:B:195:LEU:C	1:B:195:LEU:HD13	2.36	0.45
2:C:81:ASN:N	2:C:124:ARG:HH21	2.15	0.45
1:A:178:ILE:O	1:A:182:VAL:HG23	2.17	0.45
1:A:221:GLU:OE2	6:A:2145:HOH:O	2.21	0.45
1:B:56:PRO:HG2	1:B:134:ASN:HB2	1.97	0.45
1:A:17:GLN:O	1:A:40:ASP:HB2	2.16	0.45
1:B:175:PRO:O	1:B:178:ILE:HG22	2.17	0.45
1:B:48:ALA:O	1:B:51:ARG:HG3	2.17	0.44
2:C:148:LEU:N	6:C:1414:HOH:O	2.27	0.44
2:C:109:LEU:N	2:C:109:LEU:HD23	2.33	0.44
1:B:205:GLU:CG	6:B:3132:HOH:O	2.65	0.44
2:D:92:LEU:CD2	2:D:93:ARG:NH1	2.80	0.43
1:A:41:LYS:HB2	1:A:41:LYS:HE3	1.85	0.43
1:B:190:VAL:CG2	1:B:191:ARG:N	2.81	0.43
1:A:194:ILE:HD12	1:A:194:ILE:N	2.34	0.43
1:A:193:LYS:HG3	1:A:201:GLU:HG3	2.00	0.43
1:B:190:VAL:HG22	1:B:191:ARG:H	1.82	0.43
1:B:144:LYS:HD3	1:B:150:GLU:HG2	2.01	0.43
1:B:208:ASP:OD2	1:B:212:ARG:HB2	2.19	0.42
2:C:81:ASN:CG	2:C:82:VAL:H	2.22	0.42
1:B:194:ILE:HD12	1:B:230:VAL:HG22	2.01	0.41
2:C:103:ARG:HA	2:C:125:ASP:OD2	2.20	0.41
2:D:92:LEU:HD23	2:D:92:LEU:C	2.40	0.41
1:A:51:ARG:NH2	6:A:2113:HOH:O	2.53	0.41
1:A:77:LYS:HE3	6:A:2071:HOH:O	2.19	0.41
1:B:216:ARG:NH1	1:B:216:ARG:HG2	2.35	0.41
1:A:223:LYS:HE2	6:A:2169:HOH:O	2.19	0.41
2:C:92:LEU:HD23	2:C:92:LEU:HA	1.94	0.41
2:C:113:ALA:HB3	2:C:118:ASN:ND2	2.36	0.41
1:A:1:MET:SD	1:B:157:PHE:CG	3.14	0.41
1:B:80:PHE:HB3	1:B:182:VAL:CG2	2.49	0.41
1:B:227:TYR:CD2	2:D:90:LYS:HD2	2.55	0.41
1:A:19:ILE:O	1:A:42:GLN:HA	2.20	0.41
1:B:101:TRP:HA	1:B:102:PRO:HA	1.70	0.40
2:C:148:LEU:CB	6:C:1414:HOH:O	2.68	0.40
1:B:109:TYR:CD1	1:B:109:TYR:N	2.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/235 (99%)	222 (95%)	11 (5%)	0	100	100
1	B	223/235 (95%)	214 (96%)	9 (4%)	0	100	100
2	C	67/74 (90%)	66 (98%)	1 (2%)	0	100	100
2	D	67/74 (90%)	62 (92%)	5 (8%)	0	100	100
All	All	590/618 (96%)	564 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/203 (100%)	198 (98%)	5 (2%)	55	55
1	B	197/203 (97%)	189 (96%)	8 (4%)	37	32
2	C	58/63 (92%)	57 (98%)	1 (2%)	68	71
2	D	58/63 (92%)	54 (93%)	4 (7%)	19	13
All	All	516/532 (97%)	498 (96%)	18 (4%)	43	40

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	GLU
1	A	102	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	122	ASP
1	A	154	LEU
1	A	197	ASP
1	B	23	ASN
1	B	50	ASN
1	B	73	LYS
1	B	102	PRO
1	B	122	ASP
1	B	154	LEU
1	B	176	MET
1	B	191	ARG
2	C	125	ASP
2	D	112	GLU
2	D	125	ASP
2	D	132	LEU
2	D	134	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BTN	A	1401	-	12,17,17	1.34	3 (25%)	13,23,23	1.85	3 (23%)
4	ADN	A	2001	-	16,21,21	2.21	3 (18%)	16,31,31	3.11	3 (18%)
3	BTN	B	1402	-	12,17,17	1.37	3 (25%)	13,23,23	1.92	3 (23%)
4	ADN	B	2002	-	16,21,21	2.17	3 (18%)	16,31,31	3.09	3 (18%)
5	GOL	B	3001	-	5,5,5	1.07	0	5,5,5	0.86	0
3	BTN	C	1400	2	14,16,17	1.62	2 (14%)	13,21,23	1.68	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BTN	A	1401	-	-	0/5/28/28	0/2/2/2
4	ADN	A	2001	-	-	0/2/22/22	0/3/3/3
3	BTN	B	1402	-	-	0/5/28/28	0/2/2/2
4	ADN	B	2002	-	-	0/2/22/22	0/3/3/3
5	GOL	B	3001	-	-	0/4/4/4	0/0/0/0
3	BTN	C	1400	2	-	0/5/27/28	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1400	BTN	C2-S1	-2.96	1.77	1.82
3	B	1402	BTN	C2-S1	-2.58	1.78	1.82
3	A	1401	BTN	C2-S1	-2.41	1.78	1.82
3	B	1402	BTN	C5-N1	2.00	1.49	1.45
3	A	1401	BTN	O3-C3	2.06	1.27	1.23
4	B	2002	ADN	C5-C4	2.23	1.45	1.40
3	B	1402	BTN	C3-N2	2.35	1.38	1.35
4	A	2001	ADN	C5-C4	2.37	1.45	1.40
3	A	1401	BTN	C3-N2	2.46	1.38	1.35
4	B	2002	ADN	O4'-C1'	3.30	1.45	1.41
3	C	1400	BTN	C3-N2	3.72	1.40	1.35
4	A	2001	ADN	O4'-C1'	3.89	1.46	1.41
4	A	2001	ADN	C8-N7	6.85	1.47	1.34
4	B	2002	ADN	C8-N7	6.96	1.48	1.34

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2001	ADN	N3-C2-N1	-10.84	120.59	128.89
4	B	2002	ADN	N3-C2-N1	-10.71	120.69	128.89
3	B	1402	BTN	C5-N1-C3	-4.71	108.96	112.49
3	A	1401	BTN	C5-N1-C3	-4.17	109.37	112.49
4	B	2002	ADN	C2'-C1'-N9	-3.68	108.66	114.29
4	A	2001	ADN	C2'-C1'-N9	-3.68	108.67	114.29
3	C	1400	BTN	C5-N1-C3	-3.26	110.05	112.49
4	B	2002	ADN	C4'-O4'-C1'	-2.90	106.53	109.72
4	A	2001	ADN	C2-N1-C6	2.02	122.37	118.77
3	B	1402	BTN	C4-C5-N1	2.55	105.25	102.37
3	A	1401	BTN	C4-C5-N1	2.83	105.56	102.37
3	A	1401	BTN	N2-C3-N1	3.14	111.06	108.88
3	B	1402	BTN	N2-C3-N1	3.22	111.11	108.88
3	C	1400	BTN	C4-C5-N1	3.82	106.67	102.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	235/235 (100%)	0.04	4 (1%) 73 73	9, 21, 42, 58	0
1	B	227/235 (96%)	0.23	15 (6%) 22 22	12, 22, 62, 71	0
2	C	69/74 (93%)	0.71	11 (15%) 3 3	19, 34, 56, 63	0
2	D	69/74 (93%)	0.32	1 (1%) 78 78	20, 32, 44, 51	0
All	All	600/618 (97%)	0.22	31 (5%) 31 33	9, 24, 54, 71	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	LEU	6.3
1	B	200	PHE	6.0
1	B	220	GLY	5.8
2	C	149	GLY	5.1
1	A	196	GLY	5.0
2	C	126	GLY	4.5
2	C	81	ASN	3.9
1	A	197	ASP	3.8
1	B	219	SER	3.8
1	B	227	TYR	3.8
2	D	104	VAL	3.7
1	B	195	LEU	3.4
2	C	148	LEU	3.4
1	B	229	ASP	3.0
2	C	103	ARG	2.9
2	C	82	VAL	2.9
1	B	221	GLU	2.7
2	C	125	ASP	2.6
1	A	30	TYR	2.4
2	C	101	ARG	2.3
1	B	194	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	222	VAL	2.3
1	B	164	LEU	2.3
2	C	127	VAL	2.3
2	C	83	VAL	2.2
1	B	218	ASP	2.1
1	B	226	ILE	2.1
1	A	219	SER	2.1
1	B	230	VAL	2.1
1	B	109	TYR	2.0
2	C	129	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	BTN	C	1400	15/16	0.87	0.15	1.58	25,49,53,54	0
4	ADN	A	2001	19/19	0.90	0.15	0.75	17,28,42,45	0
5	GOL	B	3001	6/6	0.87	0.14	0.61	24,26,31,33	0
4	ADN	B	2002	19/19	0.92	0.14	0.11	15,23,41,44	0
3	BTN	B	1402	16/16	0.96	0.10	0.10	15,17,30,30	0
3	BTN	A	1401	16/16	0.96	0.10	-0.18	5,14,21,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.