



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:20 PM GMT

PDB ID : 4EJX  
Title : Structure of ceruloplasmin-myeloperoxidase complex  
Authors : Samygina, V.R.; Sokolov, A.V.; Bourenkov, G.; Vasilyev, V.B.; Bartunik, H.  
Deposited on : 2012-04-07  
Resolution : 4.69 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 4.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1095 (5.80-3.60)
Clashscore	102246	1010 (5.76-3.64)
Ramachandran outliers	100387	1137 (5.80-3.60)
Sidechain outliers	100360	1118 (5.80-3.60)
RSRZ outliers	91569	1098 (5.80-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1065	<div> <div></div> <div>66% 27% . .</div> </div>
2	B	114	<div> <div></div> <div>75% 15% . 9%</div> </div>
3	D	467	<div> <div></div> <div>84% 15% .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CU	A	1101	-	-	X	-
5	NAG	D	606	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 13093 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ceruloplasmin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1034	Total	C	N	O	S	0	0	0
			8380	5339	1400	1603	38			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	236	GLN	GLU	CONFLICT	UNP P00450
A	252	SER	PRO	CONFLICT	UNP P00450

- Molecule 2 is a protein called Myeloperoxidase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	104	Total	C	N	O	S	0	0	0
			837	529	148	155	5			


- Molecule 3 is a protein called Myeloperoxidase heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	466	Total	C	N	O	S	0	0	0
			3731	2351	687	666	27			

- Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	Cu	0	0
			7	7		

- Molecule 5 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



- 
- Chemical structure of HEM (heme) showing a central iron atom coordinated by four nitrogen atoms in a porphyrin-like ring, with various side chains and a central heme group.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 7 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	3	Total	C	N	O	0	0
			39	22	2	15		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Ceruloplasmin





#### • Molecule 2: Myeloperoxidase light chain





● Molecule 3: Myeloperoxidase heavy chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.25Å 106.25Å 834.62Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 4.69 15.00 – 4.69	Depositor EDS
% Data completeness (in resolution range)	98.9 (15.00-4.69) 98.9 (15.00-4.69)	Depositor EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 4.64Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.366 , 0.401 0.360 , 0.405	Depositor DCC
$R_{free}$ test set	752 reflections (5.22%)	DCC
Wilson B-factor (Å <sup>2</sup> )	135.2	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , -10.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.32$ , $\langle L^2 \rangle = 0.15$	Xtriage
Outliers	0 of 15168 reflections	Xtriage
$F_o, F_c$ correlation	0.73	EDS
Total number of atoms	13093	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, BMA, NAG, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.42	6/8615 (0.1%)	0.55	7/11689 (0.1%)
2	B	0.47	0/862	0.63	0/1174
3	D	0.47	1/3817 (0.0%)	0.59	2/5180 (0.0%)
All	All	0.44	7/13294 (0.1%)	0.57	9/18043 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
3	D	0	1
All	All	0	3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	GLU	C-N	10.20	1.57	1.34
1	A	940	HIS	C-N	6.80	1.49	1.34
1	A	945	ARG	C-N	5.84	1.47	1.34
3	D	210	LEU	C-N	5.58	1.46	1.34
1	A	252	SER	CB-OG	5.33	1.49	1.42

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	GLU	O-C-N	10.35	139.26	122.70
1	A	65	ILE	C-N-CA	-8.46	100.55	121.70
1	A	66	GLU	CA-C-N	-8.22	99.11	117.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	ILE	O-C-N	7.79	135.16	122.70
3	D	210	LEU	O-C-N	-7.49	110.72	122.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	270	GLY	Mainchain
1	A	947	PHE	Mainchain
3	D	210	LEU	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8380	0	7954	325	8
2	B	837	0	798	15	6
3	D	3731	0	3729	113	7
4	A	7	0	0	2	0
5	A	28	0	26	0	0
5	D	28	0	26	0	0
6	D	43	0	30	4	0
7	D	39	0	34	0	0
All	All	13093	0	12597	401	13

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 401 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:ASP:HA	1:A:346:ASN:CB	1.36	1.45
1:A:345:ASP:CA	1:A:346:ASN:HB2	1.55	1.36
1:A:743:ASN:CA	1:A:744:VAL:HB	1.62	1.30
1:A:340:LYS:CB	1:A:341:SER:HA	1.41	1.27
1:A:989:ARG:HG3	3:D:202:ARG:NE	1.49	1.27

The worst 5 of 13 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:925:LYS:CB	3:D:462:ARG:NH2[1_455]	1.50	0.70
1:A:706:SER:CB	3:D:157:ASN:ND2[10_555]	1.71	0.49
2:B:27:ARG:NH2	2:B:41:PHE:CE2[10_555]	1.83	0.37
1:A:925:LYS:CA	3:D:462:ARG:NH2[1_455]	1.91	0.29
2:B:27:ARG:CZ	2:B:41:PHE:CZ[10_555]	1.98	0.22

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1030/1065 (97%)	897 (87%)	110 (11%)	23 (2%)	8	50
2	B	102/114 (90%)	98 (96%)	4 (4%)	0	100	100
3	D	464/467 (99%)	446 (96%)	15 (3%)	3 (1%)	30	74
All	All	1596/1646 (97%)	1441 (90%)	129 (8%)	26 (2%)	12	57

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	119	ASN
1	A	166	ALA
1	A	350	LYS
1	A	744	VAL
1	A	890	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	909/937 (97%)	853 (94%)	56 (6%)	23	61
2	B	90/97 (93%)	88 (98%)	2 (2%)	60	84
3	D	411/412 (100%)	402 (98%)	9 (2%)	60	84
All	All	1410/1446 (98%)	1343 (95%)	67 (5%)	31	69

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	635	ASP
1	A	739	LEU
3	D	216	LEU
1	A	655	THR
1	A	702	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	677	ASN
1	A	740	GLN
2	B	54	ASN
1	A	657	ASN
3	D	421	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	NAG	D	602	3,7	14,14,15	0.50	0	15,19,21	2.26	3 (20%)
7	NAG	D	603	7	14,14,15	0.51	0	15,19,21	1.16	1 (6%)
7	BMA	D	604	7	11,11,12	0.75	0	14,15,17	1.99	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	D	602	3,7	-	0/6/23/26	0/1/1/1
7	NAG	D	603	7	-	0/6/23/26	0/1/1/1
7	BMA	D	604	7	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	604	BMA	O3-C3-C2	-4.03	102.72	110.00
7	D	602	NAG	C3-C4-C5	-2.65	105.58	110.20
7	D	604	BMA	C2-C3-C4	2.09	114.59	111.04
7	D	602	NAG	O3-C3-C2	2.28	113.62	109.11
7	D	603	NAG	C1-O5-C5	3.50	116.68	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	NAG	A	1108	1	14,14,15	0.62	0	15,19,21	1.57	2 (13%)
5	NAG	A	1109	1	14,14,15	0.54	0	15,19,21	0.92	1 (6%)
6	HEM	D	601	3	30,50,50	2.50	15 (50%)	24,82,82	2.83	10 (41%)
5	NAG	D	605	3	14,14,15	0.51	0	15,19,21	1.18	1 (6%)
5	NAG	D	606	3	14,14,15	0.52	0	15,19,21	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1108	1	-	0/6/23/26	0/1/1/1
5	NAG	A	1109	1	-	0/6/23/26	0/1/1/1
6	HEM	D	601	3	-	0/10/54/54	0/0/8/8
5	NAG	D	605	3	-	0/6/23/26	0/1/1/1
5	NAG	D	606	3	-	0/6/23/26	0/1/1/1

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	601	HEM	C2D-C3D	-5.50	1.38	1.54
6	D	601	HEM	C3B-C4B	-4.26	1.48	1.51
6	D	601	HEM	C3D-C4D	-2.84	1.47	1.51
6	D	601	HEM	C4C-NC	-2.27	1.33	1.36
6	D	601	HEM	C1C-NC	-2.07	1.33	1.36

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	601	HEM	CBD-CAD-C3D	-4.87	99.37	113.55
6	D	601	HEM	CBA-CAA-C2A	-3.08	107.00	112.53
6	D	601	HEM	CMA-C3A-C4A	-2.05	124.98	128.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
6	D	601	HEM	CMA-C3A-C2A	2.06	129.54	125.24
5	A	1109	NAG	C4-C3-C2	2.30	114.80	111.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	601	HEM	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	1034/1065 (97%)	-0.20	11 (1%) 82 76	20, 20, 20, 20	7 (0%)
2	B	104/114 (91%)	0.02	1 (0%) 84 78	20, 20, 20, 20	0
3	D	466/467 (99%)	-0.12	3 (0%) 90 86	20, 20, 20, 20	0
All	All	1604/1646 (97%)	-0.16	15 (0%) 85 80	20, 20, 20, 20	7 (0%)

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	341	SER	4.0
1	A	706	SER	3.1
1	A	257	CYS	2.9
1	A	1039	GLN	2.8
3	D	487	ARG	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
7	NAG	D	603	14/15	0.92	0.30	0.38	20,20,20,20	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	BMA	D	604	11/12	0.82	0.28	-0.65	20,20,20,20	0
7	NAG	D	602	14/15	0.77	0.34	-	20,20,20,20	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	D	606	14/15	0.79	0.40	1.00	20,20,20,20	0
5	NAG	D	605	14/15	0.72	0.36	0.94	20,20,20,20	0
4	CU	A	1107	1/1	0.86	0.40	0.72	20,20,20,20	1
4	CU	A	1104	1/1	0.84	0.26	0.27	20,20,20,20	0
6	HEM	D	601	43/43	0.89	0.30	-0.04	20,20,20,20	0
4	CU	A	1103	1/1	0.84	0.28	-0.06	20,20,20,20	0
4	CU	A	1106	1/1	0.91	0.17	-1.61	20,20,20,20	0
4	CU	A	1102	1/1	0.89	0.18	-2.42	20,20,20,20	0
4	CU	A	1105	1/1	0.93	0.07	-2.59	20,20,20,20	0
4	CU	A	1101	1/1	0.96	0.07	-4.33	20,20,20,20	0
5	NAG	A	1109	14/15	0.74	0.42	-	20,20,20,20	0
5	NAG	A	1108	14/15	0.81	0.34	-	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.