



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:12 PM GMT

PDB ID : 1EK2  
Title : CRYSTAL STRUCTURE OF MURINE SOLUBLE EPOXIDE HYDROLASE  
COMPLEXED WITH CDU INHIBITOR  
Authors : Argiriadi, M.A.; Morisseau, C.; Goodrow, M.H.; Dowdy, D.L.; Hammock,  
B.D.; Christianson, D.W.  
Deposited on : 2000-03-06  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	554	
1	B	554	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EPOXIDE HYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	487	Total	C	N	O	S	61	0	0
			3879	2501	648	701	29			
1	B	541	Total	C	N	O	S	71	0	0
			4299	2766	719	783	31			

- Molecule 2 is N-CYCLOHEXYL-N'-DECYLUREA (three-letter code: CDU) (formula:  $C_{17}H_{34}N_2O$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			20	17	2	1		
2	B	1	Total	C	N	O	0	0
			20	17	2	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total 10	O 10	0	0
3	B	9	Total 9	O 9	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: EPOXIDE HYDROLASE



#### • Molecule 1: EPOXIDE HYDROLASE



B519	K455	G389	G323	L259	P192	Q130	L66
D520	T486	V390	I324	C260	G197	D131	A67
H523	G458	A391	P325	L261	M198	P325	D68
W524	F458	E392	Q326	C262	T199	R133	E69
T525	R459	L395	A327	H263	G199	D134	S70
Q526	G460	E396	V328	G264	T200	S135	Y71
I527	P461	E396	F328	F265	L202	L136	R72
E528	L462	S400	I330	P266	L202	A137	K73
B529	M463	R401	G331	E267	V203	Q138	S74
F530	M464	T402	H332	S268	H204	M139	S75
Y531	Y465	F403	Q333	W269	N205		K76
R466	R466	F403	H334	F270	T206	L143	A77
M467	M467	K404	A335	S271	A207	S144	C78
		S405	G336	W272	S208	Q145	
R470		F406	V337	R273	A209	H146	L82
		F407	H338	Y274	L210	F147	P83
W474	W474	R408	V339	Q275	R211	D148	E84
S475	S475	A409		I276	E212	F149	N85
C476	C476	S410	H342	P277	L213	L150	F86
K477	K477	D411		A278	L213	I151	S87
		E412	Y346		K215	E152	I88
G478	G478	T413	P347	Q281	Y216	S153	S89
L479	L479	G414	E348		T217	C154	Q90
G480	G480	F415	H349		T219	Q155	
R481	R481	I416	V350	R285	G218	Y156	I91
K482	K482	A417		W286	T219		
L483	L483	V418	V353	L287	Q220	G157	Q94
Y485	Y485	H419	A354	A288	F221	M158	A95
		K420	S355	I289	P222	I159	Y96
L488	L488	A421	L356	D290		K160	
M489	M489	T422	K357	M291	P225	P161	R99
V490	V490	E423	T358	K292	L226	E162	S100
T491	T491	I424		Q293	P227	P163	I101
A492	A492	G425	H361	I294	P229	Q164	I102
E493	E493	G426	P362	G296	P229	I165	R103
K494	K494	I427	P363	S297	C230	Y166	P104
D495	D495	L428	D364	S297	N231	N167	M105
L496	L496	V429		S298	P232	F168	L106
V497	V497	M430	V367	S299	N233	L169	Q107
L498	L498	T431	S368	P300	D234	L170	A108
R499	R499	P432	P369	P301	Y235	D171	A109
P500	P500	E433	H370	E302	S236	T172	I110
B501	B501	L437	K371	I303	H237	L173	A111
M502	M502		V372	E304	G238	K174	L112
S503	S503	S438	I373	E305	Y239	A175	K113
K504	K504	R439	H374	Y306	Y240	K176	K114
M505	M505	I440	S375	A307	T241	P177	K115
M506	M506	T441	I376	M308	Y242	N178	G116
B507	B507	T442	I377	E309	K243	E179	F117
V508	V508	E443	V378	L310	P244	V180	T118
M509	M509	E444	F379	L311	G245	V181	T119
I510	I510	E445	H380		I246	F182	C120
P511	P511	I446	Y381	E314	L246	I183	I121
F512	F512	E447	Q382	M315	L248	D184	V122
L513	L513		V383	V316	H249	D185	T123
V514	V514	I450	I383	T317		F186	N124
B515	B515	Q451	V384	F318		G187	N125
G516	G516	Q452	F385	L319		S188	W126
H517	H517	F453	Q386	D320		M189	L127
I518	I518	R454	P388	K321		L190	D128
				L322		D129	

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	151.90 Å   143.00 Å   60.00 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	20.00 – 3.00	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-3.00)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.211 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8237	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CDU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/3981	0.64	0/5397
1	B	0.42	0/4413	0.61	0/5984
All	All	0.42	0/8394	0.62	0/11381

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3879	0	3863	431	0
1	B	4299	0	4270	485	0
2	A	20	0	34	5	0
2	B	20	0	34	1	0
3	A	10	0	0	0	0
3	B	9	0	0	1	0
All	All	8237	0	8201	893	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 56.

The worst 5 of 893 close contacts within the same asymmetric unit are listed below, sorted by



their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:LEU:HG	1:B:17:PRO:HA	1.29	1.10
1:A:484:LEU:HD13	1:B:61:SER:HB2	1.37	1.06
1:A:348:GLU:HA	1:B:133:ARG:HG3	1.37	1.01
1:A:122:VAL:HG22	1:A:151:ILE:HG13	1.44	1.00
1:B:122:VAL:HG22	1:B:151:ILE:HG13	1.46	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/554 (87%)	350 (73%)	98 (20%)	33 (7%)	1	7
1	B	539/554 (97%)	393 (73%)	110 (20%)	36 (7%)	1	8
All	All	1020/1108 (92%)	743 (73%)	208 (20%)	69 (7%)	1	7

5 of 69 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	18	SER
1	A	61	SER
1	A	207	ALA
1	A	231	ASN
1	A	232	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	424/480 (88%)	293 (69%)	131 (31%)	0	2
1	B	468/480 (98%)	316 (68%)	152 (32%)	0	1
All	All	892/960 (93%)	609 (68%)	283 (32%)	0	1

5 of 283 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	541	LEU
1	B	96	MET
1	B	481	ARG
1	B	4	ARG
1	B	53	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	517	HIS
1	B	107	GLN
1	B	419	HIS
1	B	85	ASN
1	B	146	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CDU	A	1100	-	20,20,20	1.77	6 (30%)	22,22,22	1.67	4 (18%)
2	CDU	B	1200	-	20,20,20	1.77	6 (30%)	22,22,22	1.67	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CDU	A	1100	-	-	0/15/23/23	0/1/1/1
2	CDU	B	1200	-	-	0/15/23/23	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1100	CDU	C2-C1	2.02	1.59	1.51
2	B	1200	CDU	C2-C1	2.03	1.59	1.51
2	B	1200	CDU	C1-C6	2.45	1.59	1.53
2	A	1100	CDU	C1-C6	2.45	1.59	1.53
2	A	1100	CDU	C3-C4	2.59	1.60	1.53

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1200	CDU	O-C8-N2	-4.26	114.84	122.75
2	A	1100	CDU	O-C8-N2	-4.26	114.84	122.75
2	B	1200	CDU	O-C8-N1	-2.81	115.77	122.76
2	A	1100	CDU	O-C8-N1	-2.78	115.84	122.76
2	A	1100	CDU	C5-N1-C8	3.33	131.85	123.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1100	CDU	5	0
2	B	1200	CDU	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.