



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:25 AM GMT

PDB ID : 3EMZ
Title : Crystal structure of xylanase XynB from *Paenibacillus barcinonensis* complexed with a conduramine derivative
Authors : Sanz-Aparicio, J.; Isorna, P.
Deposited on : 2008-09-25
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4546 (2.10-2.06)
Clashscore	102246	5101 (2.10-2.06)
Ramachandran outliers	100387	5048 (2.10-2.06)
Sidechain outliers	100360	5049 (2.10-2.06)
RSRZ outliers	91569	4556 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-1,4-beta-xylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	331	2715	1717	471	517	10	0	0	0

- Molecule 2 is (1S,2S,3R,6R)-6-[(4-PHENOXYBENZYL)AMINO]CYCLOHEX-4-ENE-1,2,3-TRIOL (three-letter code: HXH) (formula: C₁₉H₂₁NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	24	19	1	4	7	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	268	Total	O	0	0
			268	268		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endo-1,4-beta-xylanase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.02Å 79.72Å 91.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.85 – 2.08 18.85 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.6 (18.85-2.08) 91.2 (18.85-1.98)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.97Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.216 , 0.264 0.328 , 0.340	Depositor DCC
R_{free} test set	1277 reflections (7.22%)	DCC
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.469	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 45.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 21039 reflections	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	3007	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HXH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2785	0.62	1/3778 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	239	VAL	N-CA-C	-6.20	94.26	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	176	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2715	0	2574	78	0
2	A	24	0	21	1	0
3	A	268	0	0	9	0
All	All	3007	0	2595	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:TYR:HB3	1:A:126:ILE:HD13	1.60	0.81
1:A:5:ILE:HG21	1:A:321:PRO:HG3	1.62	0.80
1:A:271:TYR:HB3	1:A:328:ILE:HD11	1.63	0.79
1:A:3:THR:HG23	1:A:4:GLU:H	1.45	0.79
1:A:81:HIS:O	3:A:1001:HOH:O	2.04	0.75
1:A:314:VAL:HG11	1:A:328:ILE:HD12	1.69	0.73
1:A:18:ILE:H	1:A:39:ASN:HD21	1.39	0.69
1:A:331:GLN:O	1:A:332:ASP:HB2	1.93	0.68
1:A:325:PHE:CZ	1:A:329:ILE:HD11	2.29	0.68
1:A:28:GLN:HB2	3:A:1256:HOH:O	1.95	0.67
1:A:82:THR:H	1:A:130:ASP:HB2	1.61	0.64
1:A:270:ARG:O	1:A:274:ILE:HG12	1.99	0.63
1:A:271:TYR:CB	1:A:328:ILE:HD11	2.30	0.61
1:A:23:HIS:O	1:A:26:MET:HG3	2.00	0.61
1:A:83:LEU:HD13	1:A:115:ILE:HD12	1.83	0.61
1:A:111:LEU:O	1:A:115:ILE:HG12	2.01	0.61
1:A:2:SER:N	1:A:36:LYS:HZ2	2.01	0.58
1:A:122:TYR:CB	1:A:126:ILE:HD13	2.32	0.57
1:A:48:LYS:HG2	1:A:82:THR:CG2	2.34	0.57
1:A:81:HIS:HB3	3:A:1001:HOH:O	2.04	0.57
1:A:5:ILE:HD11	3:A:1140:HOH:O	2.05	0.56
1:A:17:LYS:HE2	3:A:1168:HOH:O	2.05	0.56
1:A:271:TYR:CG	1:A:328:ILE:HD11	2.42	0.55
1:A:271:TYR:CD1	1:A:328:ILE:HD11	2.42	0.55
1:A:218:PRO:O	1:A:270:ARG:NH2	2.30	0.55
1:A:137:GLU:HG2	1:A:145:ARG:HA	1.88	0.54
1:A:55:ARG:HH21	1:A:58:GLU:CD	2.12	0.53
1:A:29:THR:HG23	1:A:30:GLU:HG3	1.90	0.52
1:A:17:LYS:C	1:A:18:ILE:HD12	2.30	0.51
1:A:32:GLU:HG3	3:A:1219:HOH:O	2.10	0.51
1:A:50:GLU:HG2	1:A:88:GLN:O	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:GLY:HA3	1:A:313:PHE:CZ	2.46	0.50
1:A:299:TRP:HH2	2:A:1000:HXX:H16	1.76	0.50
1:A:241:GLU:HG2	1:A:291:TRP:CE3	2.47	0.50
1:A:217:GLY:C	1:A:270:ARG:NH2	2.65	0.50
1:A:320:GLN:HG3	1:A:321:PRO:HD2	1.94	0.49
1:A:18:ILE:H	1:A:39:ASN:ND2	2.07	0.49
1:A:36:LYS:NZ	3:A:1205:HOH:O	2.47	0.48
1:A:211:GLY:HA2	1:A:213:TRP:CE2	2.48	0.48
1:A:136:ILE:CD1	1:A:190:ILE:HA	2.43	0.48
1:A:136:ILE:HD13	1:A:190:ILE:HA	1.96	0.48
1:A:64:ALA:O	1:A:68:VAL:HG23	2.14	0.48
1:A:137:GLU:HG3	1:A:142:LEU:HD22	1.96	0.47
1:A:49:PHE:CD1	1:A:90:PRO:HD2	2.50	0.47
1:A:238:HIS:CD2	1:A:287:SER:HB3	2.50	0.47
1:A:314:VAL:CG1	1:A:328:ILE:HD12	2.43	0.46
1:A:166:HIS:HD2	3:A:1048:HOH:O	1.97	0.46
1:A:105:GLU:CD	1:A:105:GLU:H	2.18	0.46
1:A:266:LEU:O	1:A:270:ARG:HB2	2.14	0.46
1:A:53:HIS:HD2	1:A:114:HIS:ND1	2.13	0.46
1:A:5:ILE:HG22	1:A:326:TRP:CH2	2.52	0.45
1:A:217:GLY:C	1:A:270:ARG:HH21	2.20	0.45
1:A:204:VAL:HG13	1:A:204:VAL:O	2.17	0.45
1:A:45:ASN:O	1:A:51:GLU:HG3	2.17	0.45
1:A:139:LYS:NZ	1:A:141:ASP:OD1	2.50	0.44
1:A:82:THR:OG1	1:A:85:TRP:HB2	2.17	0.44
1:A:53:HIS:HE1	1:A:56:GLU:O	2.00	0.44
1:A:17:LYS:O	1:A:18:ILE:HD12	2.18	0.44
1:A:21:ALA:HB2	1:A:291:TRP:CE3	2.52	0.43
1:A:3:THR:HG23	1:A:4:GLU:N	2.23	0.43
1:A:55:ARG:HB2	1:A:58:GLU:HB3	2.01	0.43
1:A:5:ILE:HA	1:A:6:PRO:HD3	1.85	0.43
1:A:259:PRO:HD3	1:A:311:TRP:CD2	2.54	0.43
1:A:5:ILE:HD11	1:A:37:HIS:HE2	1.84	0.43
1:A:249:HIS:O	1:A:250:GLU:HG2	2.18	0.43
1:A:12:TYR:HB2	1:A:18:ILE:CD1	2.49	0.42
1:A:123:LYS:O	1:A:124:ASP:HB2	2.20	0.42
1:A:174:LEU:HD12	1:A:203:PRO:O	2.20	0.42
1:A:11:SER:O	1:A:332:ASP:N	2.49	0.42
1:A:212:HIS:HA	1:A:243:ASP:OD1	2.17	0.42
1:A:238:HIS:CG	1:A:287:SER:HB3	2.55	0.41
1:A:148:LYS:HG3	3:A:1101:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:ILE:O	1:A:227:ILE:HG12	2.21	0.41
1:A:5:ILE:HG13	1:A:5:ILE:O	2.20	0.41
1:A:39:ASN:HD22	1:A:39:ASN:C	2.25	0.40
1:A:12:TYR:HB2	1:A:18:ILE:HD13	2.02	0.40
1:A:39:ASN:ND2	1:A:39:ASN:H	2.19	0.40
1:A:180:ASN:HA	1:A:180:ASN:HD22	1.66	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	329/331 (99%)	312 (95%)	15 (5%)	2 (1%)	30 24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	THR
1	A	3	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/288 (100%)	278 (96%)	10 (4%)	43 43

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	26	MET
1	A	28	GLN
1	A	39	ASN
1	A	53	HIS
1	A	58	GLU
1	A	142	LEU
1	A	268	GLN
1	A	312	PRO
1	A	331	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	53	HIS
1	A	87	ASN
1	A	160	GLN
1	A	166	HIS
1	A	180	ASN
1	A	192	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HXH	A	1000	-	26,26,26	2.35	7 (26%)	29,35,35	0.98	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HXH	A	1000	-	-	0/9/26/26	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	HXH	C4-C5	-8.07	1.34	1.50
2	A	1000	HXH	C16-C15	2.26	1.42	1.38
2	A	1000	HXH	C13-C12	2.83	1.43	1.38
2	A	1000	HXH	C25-C26	3.18	1.45	1.38
2	A	1000	HXH	C10-N1	3.34	1.57	1.46
2	A	1000	HXH	O20-C21	3.45	1.47	1.39
2	A	1000	HXH	O20-C14	3.76	1.47	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	HXH	C21-O20-C14	3.65	127.91	118.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	HXH	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/331 (100%)	2.09	144 (43%) 0 0	6, 13, 24, 40	0

All (144) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	252	GLN	10.5
1	A	2	SER	10.4
1	A	250	GLU	9.2
1	A	304	PRO	7.6
1	A	28	GLN	7.4
1	A	251	ASP	7.3
1	A	297	TYR	6.3
1	A	3	THR	6.2
1	A	249	HIS	6.0
1	A	303	PHE	5.9
1	A	199	ASP	5.8
1	A	82	THR	5.7
1	A	305	VAL	4.6
1	A	254	THR	4.6
1	A	257	THR	4.5
1	A	57	HIS	4.4
1	A	195	ARG	4.3
1	A	97	ALA	4.3
1	A	299	TRP	4.3
1	A	98	SER	4.2
1	A	319	LEU	4.0
1	A	155	GLU	4.0
1	A	18	ILE	4.0
1	A	328	ILE	4.0
1	A	308	ARG	4.0
1	A	153	LEU	3.9
1	A	332	ASP	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	320	GLN	3.8
1	A	323	ASP	3.8
1	A	293	VAL	3.7
1	A	16	PHE	3.7
1	A	253	ARG	3.5
1	A	13	ALA	3.5
1	A	302	ASN	3.5
1	A	38	TYR	3.5
1	A	14	ASN	3.4
1	A	275	PHE	3.4
1	A	4	GLU	3.4
1	A	326	TRP	3.4
1	A	105	GLU	3.3
1	A	125	GLN	3.3
1	A	141	ASP	3.3
1	A	207	ILE	3.3
1	A	234	ASP	3.2
1	A	83	LEU	3.2
1	A	227	ILE	3.2
1	A	237	LEU	3.2
1	A	329	ILE	3.2
1	A	39	ASN	3.2
1	A	8	LEU	3.2
1	A	142	LEU	3.2
1	A	264	ALA	3.1
1	A	232	SER	3.1
1	A	34	ILE	3.1
1	A	67	ILE	3.1
1	A	126	ILE	3.1
1	A	71	ALA	3.1
1	A	171	ASN	3.1
1	A	230	TYR	3.0
1	A	311	TRP	3.0
1	A	41	VAL	3.0
1	A	190	ILE	3.0
1	A	282	ARG	3.0
1	A	281	TYR	3.0
1	A	152	LEU	3.0
1	A	201	GLY	3.0
1	A	285	ILE	2.9
1	A	78	VAL	2.9
1	A	163	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	24	THR	2.9
1	A	321	PRO	2.9
1	A	113	GLN	2.9
1	A	313	PHE	2.9
1	A	124	ASP	2.8
1	A	202	ALA	2.8
1	A	176	TYR	2.8
1	A	286	THR	2.8
1	A	150	LEU	2.7
1	A	300	LEU	2.7
1	A	283	SER	2.7
1	A	68	VAL	2.7
1	A	225	GLN	2.7
1	A	96	ASP	2.6
1	A	72	VAL	2.6
1	A	6	PRO	2.6
1	A	26	MET	2.6
1	A	21	ALA	2.6
1	A	192	ASN	2.6
1	A	200	GLN	2.6
1	A	62	GLU	2.6
1	A	224	ARG	2.6
1	A	271	TYR	2.6
1	A	318	GLU	2.6
1	A	111	LEU	2.6
1	A	198	LEU	2.6
1	A	256	LEU	2.6
1	A	317	THR	2.6
1	A	223	ILE	2.6
1	A	288	VAL	2.6
1	A	12	TYR	2.5
1	A	25	ARG	2.5
1	A	136	ILE	2.5
1	A	294	ALA	2.5
1	A	22	VAL	2.5
1	A	77	GLY	2.5
1	A	298	THR	2.5
1	A	277	LEU	2.5
1	A	115	ILE	2.5
1	A	213	TRP	2.5
1	A	63	ALA	2.4
1	A	193	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	239	VAL	2.4
1	A	64	ALA	2.4
1	A	248	ARG	2.4
1	A	140	THR	2.3
1	A	266	LEU	2.3
1	A	33	PHE	2.3
1	A	42	THR	2.3
1	A	127	TYR	2.3
1	A	101	THR	2.3
1	A	327	ARG	2.3
1	A	85	TRP	2.2
1	A	197	LEU	2.2
1	A	314	VAL	2.2
1	A	255	ASP	2.2
1	A	49	PHE	2.2
1	A	175	PHE	2.2
1	A	247	PHE	2.2
1	A	116	ASP	2.2
1	A	149	TRP	2.2
1	A	291	TRP	2.2
1	A	29	THR	2.2
1	A	129	TRP	2.1
1	A	122	TYR	2.1
1	A	159	VAL	2.1
1	A	9	SER	2.1
1	A	162	PHE	2.1
1	A	36	LYS	2.1
1	A	324	SER	2.1
1	A	191	TYR	2.1
1	A	280	GLU	2.0
1	A	216	HIS	2.0
1	A	17	LYS	2.0
1	A	92	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HXH	A	1000	24/24	0.81	0.21	-0.13	9,18,29,30	7

6.5 Other polymers [i](#)

There are no such residues in this entry.