



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 08:27 AM GMT

PDB ID : 3EPV
Title : X-ray Structure of the Metal-sensor CnrX in both the Apo- and Copper-bound Forms
Authors : Pompidor, G.; Maillard, A.P.; Girard, E.; Gambarelli, S.; Kahn, R.; Coves, J.
Deposited on : 2008-09-30
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2417 (1.76-1.72)
Clashscore	102246	2570 (1.76-1.72)
Ramachandran outliers	100387	2544 (1.76-1.72)
Sidechain outliers	100360	2544 (1.76-1.72)
RSRZ outliers	91569	2420 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CU	B	3	-	-	-	X

2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 3986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nickel and cobalt resistance protein cnrR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	109	Total	C 885	N 548	O 168	Se 167	2	0	3	0
1	B	109	Total	C 887	N 548	O 170	Se 167	2	0	3	0
1	C	106	Total	C 873	N 544	O 167	Se 159	3	0	3	0
1	D	109	Total	C 897	N 554	O 174	Se 167	2	0	4	0

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cu 1 1	0	0
2	A	1	Total	Cu 1 1	0	0
2	C	1	Total	Cu 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	109	Total	O 109 109	0	0
3	B	121	Total	O 121 121	0	0
3	C	127	Total	O 127 127	0	0
3	D	84	Total	O 84 84	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nickel and cobalt resistance protein cnrR




- Molecule 1: Nickel and cobalt resistance protein cnrR



- Molecule 1: Nickel and cobalt resistance protein cnrR



- Molecule 1: Nickel and cobalt resistance protein cnrR



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	31.94Å 77.89Å 93.20Å 90.00° 90.17° 90.00°	Depositor
Resolution (Å)	40.00 – 1.74 39.99 – 1.74	Depositor EDS
% Data completeness (in resolution range)	97.5 (40.00-1.74) 97.0 (39.99-1.74)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$< I/\sigma(I) >$ ¹	1.70 (at 1.74Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.196 , 0.259 0.197 , 0.257	Depositor DCC
R_{free} test set	2294 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 32.1	EDS
Estimated twinning fraction	0.217 for h,-k,-l	Xtriage
L-test for twinning ²	$< L > = 0.40$, $< L^2 > = 0.23$	Xtriage
Outliers	0 of 45536 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3986	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.50	4/906 (0.4%)	1.52	12/1220 (1.0%)
1	B	1.54	6/909 (0.7%)	1.50	8/1224 (0.7%)
1	C	1.48	8/897 (0.9%)	1.31	9/1208 (0.7%)
1	D	1.46	7/921 (0.8%)	1.34	7/1238 (0.6%)
All	All	1.49	25/3633 (0.7%)	1.42	36/4890 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	D	0	2
All	All	0	5

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	67	GLU	CG-CD	11.05	1.68	1.51
1	A	67	GLU	CD-OE2	9.56	1.36	1.25
1	D	104	ALA	CA-CB	-7.83	1.36	1.52
1	C	27	GLU	CD-OE2	-7.20	1.17	1.25
1	D	67	GLU	CB-CG	6.95	1.65	1.52
1	C	100	GLU	CG-CD	6.56	1.61	1.51
1	A	67	GLU	CB-CG	6.40	1.64	1.52
1	B	42	GLU	CB-CG	-6.33	1.40	1.52
1	C	42	GLU	CB-CG	6.30	1.64	1.52
1	B	78	ALA	CA-CB	6.30	1.65	1.52
1	B	33	GLU	CG-CD	6.11	1.61	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	92	GLU	CD-OE1	6.01	1.32	1.25
1	C	36	PHE	CE2-CZ	5.98	1.48	1.37
1	C	80	GLY	N-CA	5.89	1.54	1.46
1	D	67	GLU	CG-CD	5.69	1.60	1.51
1	D	69	GLU	CB-CG	-5.66	1.41	1.52
1	C	64	TRP	CE3-CZ3	5.43	1.47	1.38
1	D	114	ARG	CG-CD	5.40	1.65	1.51
1	B	30	GLU	N-CA	5.35	1.57	1.46
1	D	69	GLU	CD-OE1	5.27	1.31	1.25
1	A	108	VAL	CB-CG1	5.27	1.64	1.52
1	C	95	ALA	CA-CB	-5.25	1.41	1.52
1	C	27	GLU	CG-CD	5.20	1.59	1.51
1	D	63	ALA	CA-CB	5.07	1.63	1.52
1	B	36	PHE	CG-CD2	-5.02	1.31	1.38

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	84	ARG	NE-CZ-NH2	-21.47	109.56	120.30
1	B	84	ARG	NE-CZ-NH1	15.11	127.86	120.30
1	A	46	ARG	NE-CZ-NH1	14.45	127.53	120.30
1	A	107	ARG	NE-CZ-NH2	-14.40	113.10	120.30
1	D	46	ARG	NE-CZ-NH2	-12.99	113.81	120.30
1	A	107	ARG	NE-CZ-NH1	12.88	126.74	120.30
1	D	46	ARG	NE-CZ-NH1	12.19	126.40	120.30
1	B	82	LEU	CB-CG-CD1	-10.64	92.92	111.00
1	A	41	ARG	NE-CZ-NH1	10.58	125.59	120.30
1	A	46	ARG	NE-CZ-NH2	-10.46	115.07	120.30
1	B	46	ARG	NE-CZ-NH1	9.40	125.00	120.30
1	B	46	ARG	NE-CZ-NH2	-9.32	115.64	120.30
1	C	84[A]	ARG	NE-CZ-NH2	-8.94	115.83	120.30
1	C	84[B]	ARG	NE-CZ-NH2	-8.94	115.83	120.30
1	A	41	ARG	NE-CZ-NH2	-7.75	116.43	120.30
1	C	46	ARG	NE-CZ-NH1	7.50	124.05	120.30
1	C	46	ARG	NE-CZ-NH2	-7.21	116.69	120.30
1	D	56	ASP	CB-CG-OD1	6.87	124.48	118.30
1	C	56	ASP	CB-CG-OD2	-6.61	112.35	118.30
1	B	15	LEU	N-CA-C	6.56	128.72	111.00
1	A	81	ASP	CB-CG-OD2	-6.53	112.42	118.30
1	A	102	ARG	NE-CZ-NH2	-6.46	117.07	120.30
1	C	77	ARG	NE-CZ-NH1	6.19	123.40	120.30
1	C	77	ARG	NE-CZ-NH2	-6.18	117.21	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	22	ASP	CB-CG-OD2	5.93	123.64	118.30
1	D	77	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	C	40	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	B	84	ARG	CG-CD-NE	-5.76	99.71	111.80
1	C	106	ASP	CB-CG-OD1	5.65	123.38	118.30
1	A	81	ASP	CB-CG-OD1	5.60	123.34	118.30
1	A	26	ARG	NE-CZ-NH1	-5.37	117.61	120.30
1	D	102	ARG	NE-CZ-NH2	-5.35	117.63	120.30
1	D	34	ASP	N-CA-C	5.34	125.43	111.00
1	B	26	ARG	NE-CZ-NH2	5.34	122.97	120.30
1	A	41	ARG	CD-NE-CZ	5.17	130.83	123.60
1	D	19	VAL	CB-CA-C	-5.14	101.63	111.40

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	ASP	Peptide
1	B	15	LEU	Peptide
1	B	17	GLU	Peptide
1	D	19	VAL	Peptide
1	D	21	LEU	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	885	0	889	19	0
1	B	887	0	885	26	0
1	C	873	0	891	16	0
1	D	897	0	906	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	109	0	0	8	1
3	B	121	0	0	5	1
3	C	127	0	0	6	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	84	0	0	3	0
All	All	3986	0	3571	104	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:VAL:HG12	3:A:170:HOH:O	1.46	1.15
1:B:19:VAL:HG21	3:B:307:HOH:O	1.44	1.14
1:D:29:LEU:O	1:D:93[B]:MSE:HE1	1.60	1.01
1:A:29:LEU:HD13	1:A:93[B]:MSE:HE3	1.43	0.99
1:D:34:ASP:H	1:D:35:ALA:HB3	1.24	0.99
1:D:45:THR:HA	1:D:48:ARG:HD2	1.45	0.97
1:D:35:ALA:O	1:D:37:ALA:N	1.97	0.97
1:A:29:LEU:HD13	1:A:93[B]:MSE:CE	2.05	0.86
1:D:31:LEU:HD13	1:D:32:LYS:H	1.40	0.86
1:D:32:LYS:HB2	3:D:179:HOH:O	1.74	0.85
1:D:45:THR:HA	1:D:48:ARG:CD	2.11	0.81
1:B:14:ILE:O	1:B:18:ALA:HB2	1.82	0.79
1:B:30:GLU:OE2	3:B:306:HOH:O	2.01	0.77
1:A:16:HIS:CE1	1:A:33:GLU:OE2	2.38	0.77
1:C:115:ARG:HA	1:C:115:ARG:NE	1.99	0.77
1:C:115:ARG:HE	1:C:115:ARG:HA	1.51	0.76
1:B:39:ARG:O	1:B:42:GLU:HG2	1.87	0.75
1:D:19:VAL:HG23	1:D:21:LEU:HD12	1.70	0.74
1:C:16:HIS:HE1	1:C:93[B]:MSE:HE2	1.52	0.73
1:D:34:ASP:N	1:D:35:ALA:HB3	2.01	0.73
1:D:35:ALA:O	1:D:36:PHE:C	2.28	0.71
1:D:24:ASN:O	1:D:28:ILE:HD13	1.91	0.71
1:C:15:LEU:O	3:C:192:HOH:O	2.09	0.70
1:C:18:ALA:N	3:C:192:HOH:O	2.25	0.70
1:B:16:HIS:O	1:B:17:GLU:C	2.30	0.69
1:C:28[B]:ILE:HD12	3:C:248:HOH:O	1.93	0.69
1:B:111[B]:ASP:OD1	3:B:249:HOH:O	2.12	0.67
1:C:28[B]:ILE:CD1	3:C:248:HOH:O	2.43	0.67
1:D:31:LEU:HG	3:D:183:HOH:O	1.94	0.67
1:D:31:LEU:HD13	1:D:32:LYS:N	2.08	0.67
1:C:40:ARG:NH1	1:C:82:LEU:HD11	2.09	0.66
1:D:32:LYS:O	1:D:35:ALA:HB1	1.96	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:45:THR:O	1:D:48:ARG:HG2	1.98	0.64
1:B:43:ILE:HG21	1:B:82:LEU:HD23	1.80	0.63
1:D:48:ARG:HG2	1:D:49:ALA:N	2.14	0.62
1:D:31:LEU:HD22	1:D:31:LEU:C	2.20	0.61
1:A:19:VAL:CG1	3:A:170:HOH:O	2.22	0.61
1:D:19:VAL:CG2	1:D:21:LEU:HD12	2.31	0.61
1:A:29:LEU:HD22	1:A:93[B]:MSE:SE	2.51	0.59
1:C:114:ARG:O	1:C:115:ARG:HD2	2.03	0.58
1:D:28:ILE:N	1:D:28:ILE:HD12	2.18	0.58
1:D:29:LEU:O	1:D:33:GLU:HB2	2.03	0.57
1:D:28:ILE:N	1:D:28:ILE:CD1	2.66	0.57
1:A:22:ASP:OD1	1:A:98:LYS:NZ	2.24	0.57
1:B:114:ARG:NH2	3:B:311:HOH:O	2.34	0.57
1:B:87:LEU:O	1:B:90:VAL:HG22	2.05	0.57
1:D:10:ASP:HA	1:D:13:GLU:HG3	1.87	0.56
1:A:19:VAL:O	1:A:19:VAL:HG12	2.05	0.55
1:D:31:LEU:CD1	1:D:32:LYS:HG3	2.36	0.55
1:B:15:LEU:O	1:B:18:ALA:HB3	2.06	0.54
1:B:89:HIS:CE1	1:B:93[B]:MSE:SE	3.11	0.54
1:D:29:LEU:HD23	3:D:171:HOH:O	2.08	0.54
1:D:21:LEU:HD22	1:D:26:ARG:HB2	1.91	0.52
1:B:12[B]:HIS:CD2	1:B:86:THR:HG23	2.44	0.52
1:C:53:LYS:NZ	3:C:188:HOH:O	2.43	0.52
1:A:42:GLU:OE1	3:A:228:HOH:O	2.19	0.51
1:B:12[B]:HIS:CE1	1:B:90:VAL:HG13	2.46	0.51
1:D:31:LEU:HD13	1:D:32:LYS:HG3	1.93	0.51
1:D:19:VAL:CG2	1:D:21:LEU:CD1	2.89	0.51
1:A:114:ARG:HB3	1:A:115:ARG:HH21	1.75	0.50
1:B:79:ALA:HB2	3:B:299:HOH:O	2.12	0.50
1:A:29:LEU:HB3	1:A:93[B]:MSE:HE1	1.94	0.50
1:C:114:ARG:O	1:C:115:ARG:HB2	2.11	0.50
1:D:29:LEU:C	1:D:93[B]:MSE:HE1	2.32	0.49
1:A:87[A]:LEU:HG	3:A:154:HOH:O	2.10	0.49
1:D:30[B]:GLU:HA	1:D:33:GLU:HB3	1.95	0.49
1:D:48:ARG:HG2	1:D:49:ALA:H	1.77	0.49
1:B:33:GLU:OE1	1:B:93[B]:MSE:HE2	2.13	0.48
1:D:22:ASP:OD2	1:D:101:HIS:HE1	1.96	0.48
1:D:32:LYS:O	1:D:35:ALA:CB	2.61	0.47
1:D:31:LEU:O	1:D:34:ASP:OD2	2.33	0.47
1:A:21:LEU:HD11	1:A:93[B]:MSE:CE	2.46	0.46
1:B:19:VAL:CG2	1:B:20:PRO:HD2	2.45	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:ARG:NH1	3:A:187:HOH:O	2.49	0.46
1:C:38:GLN:O	1:C:42:GLU:CD	2.54	0.46
1:B:15:LEU:HA	1:B:18:ALA:CB	2.46	0.46
1:D:21:LEU:N	1:D:25:GLU:OE1	2.47	0.46
1:A:115:ARG:CA	1:A:115:ARG:HE	2.30	0.44
1:D:26:ARG:HG3	1:D:26:ARG:O	2.16	0.44
1:A:42:GLU:HB2	3:A:228:HOH:O	2.18	0.44
1:A:19:VAL:HG11	1:A:105:TYR:HD1	1.83	0.44
1:A:26:ARG:NH2	3:A:167:HOH:O	2.49	0.44
1:B:10:ASP:N	1:B:13:GLU:HG3	2.32	0.44
1:D:29:LEU:O	1:D:33:GLU:CB	2.66	0.43
1:C:89:HIS:CE1	1:C:93[B]:MSE:SE	3.21	0.43
1:A:41:ARG:HD2	3:A:132:HOH:O	2.17	0.43
1:D:19:VAL:HG21	1:D:21:LEU:HD11	2.00	0.43
1:B:12[A]:HIS:CE1	1:B:33:GLU:OE2	2.72	0.43
1:B:12[B]:HIS:NE2	1:B:86:THR:HA	2.34	0.42
1:B:19:VAL:HG23	1:B:20:PRO:HD2	2.01	0.42
1:B:17:GLU:OE2	1:B:17:GLU:HA	2.19	0.42
1:D:20:PRO:HA	1:D:21:LEU:HB2	2.00	0.42
1:C:40:ARG:HH11	1:C:82:LEU:HD11	1.80	0.42
1:D:21:LEU:HD22	1:D:26:ARG:HA	2.01	0.42
1:D:38:GLN:O	1:D:42:GLU:HG3	2.20	0.42
1:B:12[B]:HIS:CE1	1:B:89:HIS:HB3	2.55	0.41
1:D:31:LEU:HD22	1:D:31:LEU:O	2.20	0.41
1:C:16:HIS:C	3:C:192:HOH:O	2.57	0.41
1:D:48:ARG:CG	1:D:49:ALA:N	2.82	0.40
1:C:115:ARG:CA	1:C:115:ARG:NE	2.78	0.40
1:D:57:ALA:HA	1:D:60:LYS:HE2	2.04	0.40
1:B:22:ASP:OD2	1:B:98:LYS:NZ	2.37	0.40
1:B:51:ASN:HA	1:B:51:ASN:HD22	1.78	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:128:HOH:O	3:C:235:HOH:O[2_555]	1.88	0.32
3:B:209:HOH:O	3:C:241:HOH:O[2_555]	2.06	0.14

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	110/109 (101%)	109 (99%)	1 (1%)	0	100 100
1	B	110/109 (101%)	106 (96%)	3 (3%)	1 (1%)	21 6
1	C	108/109 (99%)	106 (98%)	2 (2%)	0	100 100
1	D	111/109 (102%)	102 (92%)	5 (4%)	4 (4%)	4 0
All	All	439/436 (101%)	423 (96%)	11 (2%)	5 (1%)	16 4

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	16	HIS
1	D	36	PHE
1	D	35	ALA
1	D	20	PRO
1	D	21	LEU

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	90/86 (105%)	85 (94%)	5 (6%)	26 6
1	B	90/86 (105%)	89 (99%)	1 (1%)	80 66
1	C	89/86 (104%)	86 (97%)	3 (3%)	44 18
1	D	91/86 (106%)	83 (91%)	8 (9%)	12 11
All	All	360/344 (105%)	343 (95%)	17 (5%)	33 9

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	67	GLU
1	A	77	ARG
1	A	115	ARG
1	A	118	GLN
1	B	17	GLU
1	C	42	GLU
1	C	60	LYS
1	C	115	ARG
1	D	10	ASP
1	D	22	ASP
1	D	29	LEU
1	D	30[A]	GLU
1	D	30[B]	GLU
1	D	31	LEU
1	D	33	GLU
1	D	48	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	GLN
1	B	51	ASN
1	C	16	HIS
1	C	83	GLN
1	C	89	HIS
1	D	16	HIS
1	D	51	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	108/109 (99%)	0.33	4 (3%) 45 51	13, 21, 37, 42	0
1	B	108/109 (99%)	0.13	4 (3%) 45 51	13, 21, 39, 48	0
1	C	105/109 (96%)	0.13	3 (2%) 55 61	12, 20, 40, 47	0
1	D	108/109 (99%)	0.80	18 (16%) 2 3	16, 24, 52, 56	0
All	All	429/436 (98%)	0.35	29 (6%) 20 25	12, 22, 45, 56	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	31	LEU	6.8
1	D	19	VAL	6.7
1	D	32	LYS	4.9
1	D	41	ARG	4.6
1	D	18	ALA	4.2
1	B	17	GLU	4.1
1	D	21	LEU	4.0
1	D	22	ASP	3.8
1	D	34	ASP	3.6
1	C	17	GLU	3.5
1	D	118	GLN	3.5
1	D	28	ILE	3.2
1	A	118	GLN	3.1
1	C	16	HIS	3.1
1	D	10	ASP	3.0
1	D	30[A]	GLU	3.0
1	D	16	HIS	3.0
1	A	115	ARG	3.0
1	D	20	PRO	2.9
1	A	26	ARG	2.8
1	D	17	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	10	ASP	2.6
1	B	14	ILE	2.5
1	D	48	ARG	2.3
1	B	16	HIS	2.3
1	D	13	GLU	2.2
1	A	20	PRO	2.1
1	D	29	LEU	2.1
1	B	11	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CU	B	3	1/1	0.84	0.27	2.90	33,33,33,33	1
2	CU	A	1	1/1	0.97	0.04	-	30,30,30,30	1
2	CU	C	2	1/1	0.96	0.08	-	35,35,35,35	1

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.