



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:25 PM GMT

PDB ID : 4ESA  
Title : X-ray structure of carbonmonoxy hemoglobin of Elegendops maclovinus  
Authors : Merlino, A.; Vitagliano, L.; Mazzarella, L.; Vergara, A.  
Deposited on : 2012-04-23  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1278 (1.48-1.44)
Clashscore	102246	1336 (1.48-1.44)
Ramachandran outliers	100387	1320 (1.48-1.44)
Sidechain outliers	100360	1320 (1.48-1.44)
RSRZ outliers	91569	1279 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	143	<div> <div>3%</div> <div>89%</div> <div>9%</div> <div>.</div> </div>
1	C	143	<div> <div>3%</div> <div>86%</div> <div>11%</div> <div>.</div> </div>
2	B	146	<div> <div>9%</div> <div>76%</div> <div>18%</div> <div>.</div> </div>
2	D	146	<div> <div>3%</div> <div>79%</div> <div>16%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	D	203	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hemoglobin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	143	Total	C	N	O	S	0	1	0
			1107	713	188	201	5			
1	C	143	Total	C	N	O	S	0	1	0
			1107	713	188	201	5			

- Molecule 2 is a protein called Hemoglobin beta chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	144	Total	C	N	O	S	0	3	0
			1128	725	193	206	4			
2	D	145	Total	C	N	O	S	0	1	0
			1131	727	192	208	4			

- Molecule 3 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 2	C 1	O 1	0	0
3	B	1	Total 2	C 1	O 1	0	0
3	C	1	Total 2	C 1	O 1	0	0
3	D	1	Total 2	C 1	O 1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
4	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
4	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
4	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			6	3	3		



- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	104	Total	O	0	0
			104	104		
6	B	72	Total	O	0	0
			72	72		
6	C	105	Total	O	0	0
			105	105		
6	D	127	Total	O	0	0
			127	127		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Hemoglobin alpha chain



- Molecule 1: Hemoglobin alpha chain




- Molecule 2: Hemoglobin beta chain



HIS

- Molecule 2: Hemoglobin beta chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.17Å 88.08Å 123.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.45 23.88 – 1.45	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.45) 91.2 (23.88-1.45)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.52 (at 1.45Å)	Xtriage
Refinement program	SHELXL-93	Depositor
R, $R_{free}$	0.212 , 0.253 0.189 , 0.194	Depositor DCC
$R_{free}$ test set	8771 reflections (9.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.9	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 53.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 109671 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5067	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, GOL, ACE, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/1134	1.05	3/1533 (0.2%)
1	C	0.47	0/1134	1.04	3/1533 (0.2%)
2	B	0.43	0/1165	1.07	3/1578 (0.2%)
2	D	0.48	0/1159	1.16	7/1570 (0.4%)
All	All	0.46	0/4592	1.08	16/6214 (0.3%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	40	ARG	NE-CZ-NH2	10.97	125.78	120.30
2	D	8	ARG	NE-CZ-NH2	-9.63	115.49	120.30
2	B	8	ARG	NE-CZ-NH1	-8.44	116.08	120.30
2	D	40	ARG	NE-CZ-NH1	-7.79	116.41	120.30
2	B	108	ASP	CB-CG-OD1	7.03	124.63	118.30
2	B	8	ARG	CD-NE-CZ	-6.92	113.91	123.60
2	D	45	PHE	CB-CG-CD2	6.00	125.00	120.80
1	C	31	ARG	NE-CZ-NH2	-5.89	117.35	120.30
2	D	72	ASP	CB-CG-OD1	5.80	123.52	118.30
1	C	141	TYR	CB-CG-CD1	-5.62	117.63	121.00
2	D	72	ASP	CB-CG-OD2	-5.50	113.35	118.30
1	C	141	TYR	CB-CG-CD2	5.31	124.19	121.00
1	A	31	ARG	NE-CZ-NH1	5.13	122.86	120.30
2	D	5	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	76	ASP	CB-CG-OD1	5.03	122.83	118.30
1	A	90	TYR	CB-CG-CD2	-5.01	117.99	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1107	0	1140	13	0
1	C	1107	0	1140	13	0
2	B	1128	0	1135	26	0
2	D	1131	0	1134	18	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	43	0	30	1	0
4	B	43	0	30	1	0
4	C	43	0	30	0	0
4	D	43	0	30	0	0
5	D	6	0	8	0	0
6	A	104	0	0	3	0
6	B	72	0	0	1	0
6	C	105	0	0	1	0
6	D	127	0	0	0	0
All	All	5067	0	4677	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:GLN:HE21	1:A:38:GLN:H	1.10	0.94
2:D:80:ASN:O	2:D:84:ILE:HG23	1.84	0.77
1:C:33:ILE:HG23	1:C:40:LYS:HG2	1.66	0.76
1:C:30:SER:O	1:C:34[A]:VAL:HG22	1.84	0.76
2:B:37:TRP:HE1	2:B:102:ASN:HD21	1.36	0.73
1:A:30:SER:O	1:A:34[A]:VAL:HG12	1.88	0.72
2:B:101:ASP:HB3	6:B:358:HOH:O	1.90	0.72
1:C:37:PRO:HB3	1:C:40:LYS:HE3	1.73	0.71
1:C:38:GLN:HE21	1:C:38:GLN:H	1.34	0.71
2:B:90:ILE:O	2:B:94:GLU:HB3	1.95	0.66

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:37:TRP:HE1	2:D:102:ASN:HD21	1.44	0.64
1:C:87:GLN:HG3	1:C:91:LYS:HE3	1.80	0.62
1:A:38:GLN:H	1:A:38:GLN:NE2	1.90	0.61
2:B:89:SER:O	2:B:93:SER:HB2	2.02	0.59
2:D:85:TYR:HD2	2:D:88:LEU:HD12	1.67	0.58
1:A:34[A]:VAL:HG13	1:A:35:VAL:HG23	1.86	0.58
2:D:85:TYR:CD2	2:D:88:LEU:HD12	2.39	0.58
2:B:143:LYS:HD3	2:B:144:GLN:N	2.18	0.58
2:D:139:SER:O	2:D:143:LYS:HD2	2.04	0.57
2:B:123:ASN:ND2	2:B:126:THR:H	2.03	0.56
2:B:123:ASN:HD22	2:B:126:THR:H	1.53	0.56
2:B:84:ILE:HD12	2:B:85:TYR:CD2	2.42	0.55
1:A:19:LYS:HG2	6:A:342:HOH:O	2.07	0.55
2:B:75:VAL:HG12	2:B:76:LYS:HE3	1.89	0.55
2:D:90:ILE:CG1	2:D:144:GLN:HG2	2.39	0.53
1:C:33:ILE:O	1:C:40:LYS:HE2	2.09	0.53
2:D:80:ASN:OD1	2:D:83:GLU:HB3	2.10	0.52
2:B:143:LYS:O	2:B:144:GLN:HG2	2.10	0.52
2:D:37:TRP:HE1	2:D:102:ASN:ND2	2.09	0.51
2:B:37:TRP:HE1	2:B:102:ASN:ND2	2.06	0.51
2:B:75:VAL:HB	2:B:76:LYS:NZ	2.25	0.51
2:D:130:PHE:CE2	2:D:134:LEU:HD11	2.47	0.50
2:B:59:LYS:HB3	2:B:59:LYS:NZ	2.27	0.49
2:D:143:LYS:HE3	2:D:143:LYS:HA	1.95	0.48
2:D:123:ASN:ND2	2:D:126:THR:H	2.12	0.47
2:B:80:ASN:HD22	2:B:83:GLU:CB	2.27	0.47
2:B:75:VAL:HB	2:B:76:LYS:HZ1	1.79	0.47
2:B:80:ASN:HD22	2:B:83:GLU:HB2	1.79	0.46
2:B:74:ALA:HB1	2:B:81:ILE:HG12	1.98	0.46
1:A:34[A]:VAL:HG11	2:B:124:PRO:HB3	1.96	0.46
1:A:116:LYS:HG2	6:A:321:HOH:O	2.16	0.46
1:C:82:LEU:HD23	1:C:82:LEU:HA	1.83	0.46
2:B:40:ARG:HH22	1:C:41:THR:HG23	1.81	0.46
1:C:116:LYS:HG2	1:C:116:LYS:O	2.12	0.45
2:D:123:ASN:HD22	2:D:126:THR:H	1.64	0.45
1:A:5:LYS:NZ	6:A:393:HOH:O	2.50	0.45
1:C:116:LYS:HD3	1:C:117:GLU:OE2	2.17	0.44
2:B:72:ASP:O	2:B:76:LYS:NZ	2.50	0.44
2:D:84:ILE:HD11	2:D:85:TYR:CZ	2.53	0.44
1:A:57:LYS:O	1:A:61:LYS:HG3	2.18	0.44
1:A:34[B]:VAL:HG23	1:A:35:VAL:HG23	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:57:ASN:ND2	2:D:60:VAL:H	2.15	0.43
1:A:22:ASP:CG	1:A:61:LYS:HE2	2.38	0.43
2:B:22:ASP:OD1	2:B:26:LYS:HE3	2.18	0.43
1:A:34[A]:VAL:HG13	1:A:35:VAL:N	2.34	0.43
1:C:19:LYS:HG2	6:C:378:HOH:O	2.18	0.42
2:D:90:ILE:HG12	2:D:144:GLN:HG2	2.00	0.42
2:B:95:LYS:HB3	2:B:96[B]:LEU:HD13	2.00	0.42
1:C:86:GLU:OE1	1:C:140:ARG:HD2	2.20	0.42
2:D:89:SER:OG	2:D:144:GLN:HB2	2.20	0.42
2:B:1:VAL:O	2:B:1:VAL:HG13	2.20	0.41
2:B:143:LYS:HD3	2:B:144:GLN:H	1.83	0.41
1:C:111:SER:OG	2:D:116:ALA:HB2	2.21	0.41
2:B:38:THR:HB	4:B:202:HEM:CBC	2.51	0.41
2:D:14:ILE:O	2:D:18:LEU:HG	2.21	0.40
1:A:88:HIS:CE1	4:A:202:HEM:NA	2.89	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/143 (99%)	141 (99%)	1 (1%)	0	100	100
1	C	142/143 (99%)	141 (99%)	1 (1%)	0	100	100
2	B	145/146 (99%)	142 (98%)	3 (2%)	0	100	100
2	D	144/146 (99%)	143 (99%)	1 (1%)	0	100	100
All	All	573/578 (99%)	567 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/122 (101%)	118 (96%)	5 (4%)	37	6
1	C	123/122 (101%)	117 (95%)	6 (5%)	31	4
2	B	122/121 (101%)	114 (93%)	8 (7%)	21	1
2	D	121/121 (100%)	113 (93%)	8 (7%)	21	1
All	All	489/486 (101%)	462 (94%)	27 (6%)	26	3

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LYS
1	A	29	LEU
1	A	38	GLN
1	A	49	LEU
1	A	116	LYS
2	B	30	ARG
2	B	76	LYS
2	B	80	ASN
2	B	83	GLU
2	B	95	LYS
2	B	120	SER
2	B	123	ASN
2	B	143	LYS
1	C	19	LYS
1	C	27	ASP
1	C	29	LEU
1	C	38	GLN
1	C	82	LEU
1	C	116	LYS
2	D	57	ASN
2	D	59	LYS
2	D	68	LEU
2	D	77	ASN
2	D	84	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	123	ASN
2	D	143	LYS
2	D	144	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	38	GLN
2	B	80	ASN
2	B	102	ASN
2	B	123	ASN
2	B	131	GLN
2	B	144	GLN
1	C	38	GLN
2	D	57	ASN
2	D	102	ASN
2	D	123	ASN
2	D	131	GLN
2	D	144	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CMO	A	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	A	202	1,3	30,50,50	2.58	7 (23%)	24,82,82	2.61	9 (37%)
3	CMO	B	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	B	202	3,2	30,50,50	2.41	7 (23%)	24,82,82	2.77	14 (58%)
3	CMO	C	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	C	202	1,3	30,50,50	2.59	8 (26%)	24,82,82	2.63	10 (41%)
3	CMO	D	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	D	202	3,2	30,50,50	2.39	6 (20%)	24,82,82	2.55	8 (33%)
5	GOL	D	203	-	5,5,5	0.11	0	5,5,5	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CMO	A	201	4	-	0/0/0/0	0/0/0/0
4	HEM	A	202	1,3	-	0/10/54/54	0/0/8/8
3	CMO	B	201	4	-	0/0/0/0	0/0/0/0
4	HEM	B	202	3,2	-	0/10/54/54	0/0/8/8
3	CMO	C	201	4	-	0/0/0/0	0/0/0/0
4	HEM	C	202	1,3	-	0/10/54/54	0/0/8/8
3	CMO	D	201	4	-	0/0/0/0	0/0/0/0
4	HEM	D	202	3,2	-	0/10/54/54	0/0/8/8
5	GOL	D	203	-	-	0/4/4/4	0/0/0/0

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	202	HEM	C3B-C4B	-8.33	1.44	1.51
4	C	202	HEM	C3B-C4B	-8.15	1.44	1.51
4	B	202	HEM	C3B-C4B	-7.06	1.45	1.51
4	C	202	HEM	C2D-C3D	-7.02	1.33	1.54
4	B	202	HEM	C2D-C3D	-6.80	1.34	1.54
4	D	202	HEM	C2D-C3D	-6.69	1.34	1.54
4	D	202	HEM	C3B-C4B	-6.63	1.45	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	202	HEM	C2D-C3D	-6.42	1.35	1.54
4	A	202	HEM	C3D-C4D	-6.03	1.43	1.51
4	D	202	HEM	C3D-C4D	-5.67	1.44	1.51
4	C	202	HEM	C3D-C4D	-5.44	1.44	1.51
4	B	202	HEM	C3D-C4D	-4.72	1.45	1.51
4	B	202	HEM	C2C-C1C	-4.01	1.45	1.52
4	A	202	HEM	C2C-C1C	-3.71	1.45	1.52
4	C	202	HEM	C2C-C1C	-3.58	1.45	1.52
4	D	202	HEM	C2C-C1C	-3.24	1.46	1.52
4	B	202	HEM	C2B-C1B	-2.35	1.44	1.51
4	D	202	HEM	C2D-C1D	-2.17	1.44	1.51
4	B	202	HEM	C2D-C1D	-2.16	1.44	1.51
4	A	202	HEM	C2D-C1D	-2.16	1.44	1.51
4	A	202	HEM	C2B-C1B	-2.04	1.45	1.51
4	C	202	HEM	C2B-C1B	-2.03	1.45	1.51
4	C	202	HEM	FE-NC	2.02	2.03	1.95
4	A	202	HEM	C4C-NC	2.22	1.38	1.36
4	B	202	HEM	C1C-NC	2.33	1.38	1.36
4	C	202	HEM	C3B-CAB	2.33	1.55	1.51
4	D	202	HEM	CAA-C2A	2.35	1.56	1.52
4	C	202	HEM	C4C-NC	2.39	1.38	1.36

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	202	HEM	CAA-C2A-C1A	-3.96	122.71	127.01
4	A	202	HEM	CAA-C2A-C1A	-3.92	122.75	127.01
4	B	202	HEM	CMA-C3A-C4A	-3.85	121.99	128.36
4	B	202	HEM	CAA-C2A-C1A	-3.85	122.83	127.01
4	C	202	HEM	CAA-C2A-C1A	-3.55	123.15	127.01
4	C	202	HEM	C3B-CAB-CBB	-3.16	119.60	124.46
4	D	202	HEM	CMA-C3A-C4A	-2.92	123.54	128.36
4	A	202	HEM	CMA-C3A-C4A	-2.90	123.56	128.36
4	C	202	HEM	CMA-C3A-C4A	-2.88	123.61	128.36
4	B	202	HEM	C3B-CAB-CBB	-2.38	120.81	124.46
4	B	202	HEM	CAA-CBA-CGA	2.06	116.53	112.75
4	B	202	HEM	CMA-C3A-C2A	2.08	129.58	125.24
4	B	202	HEM	C3B-C4B-CHC	2.23	126.31	123.16
4	C	202	HEM	CBA-CAA-C2A	2.47	116.96	112.53
4	B	202	HEM	CBA-CAA-C2A	2.64	117.26	112.53
4	B	202	HEM	CAD-C3D-C4D	2.64	121.78	112.47
4	B	202	HEM	CBD-CAD-C3D	2.77	121.62	113.55

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	202	HEM	C3B-C4B-CHC	2.85	127.18	123.16
4	D	202	HEM	CMD-C2D-C3D	2.94	127.37	114.35
4	B	202	HEM	CMD-C2D-C3D	2.95	127.39	114.35
4	B	202	HEM	C2D-C3D-C4D	3.16	106.86	101.50
4	A	202	HEM	CAD-C3D-C4D	3.18	123.69	112.47
4	C	202	HEM	CMD-C2D-C3D	3.23	128.65	114.35
4	D	202	HEM	CAD-C3D-C4D	3.25	123.92	112.47
4	C	202	HEM	CAD-C3D-C4D	3.28	124.05	112.47
4	A	202	HEM	CMD-C2D-C3D	3.32	129.04	114.35
4	D	202	HEM	C2D-C3D-C4D	3.69	107.76	101.50
4	A	202	HEM	C2D-C3D-C4D	3.79	107.93	101.50
4	C	202	HEM	C2D-C3D-C4D	3.93	108.16	101.50
4	A	202	HEM	CAD-C3D-C2D	4.98	127.54	113.22
4	A	202	HEM	CMB-C2B-C3B	4.99	128.98	116.53
4	B	202	HEM	CMB-C2B-C3B	5.02	129.05	116.53
4	C	202	HEM	CMC-C2C-C3C	5.06	129.16	116.53
4	C	202	HEM	CAD-C3D-C2D	5.06	127.77	113.22
4	D	202	HEM	CAD-C3D-C2D	5.11	127.90	113.22
4	B	202	HEM	CMC-C2C-C3C	5.13	129.33	116.53
4	D	202	HEM	CMB-C2B-C3B	5.21	129.53	116.53
4	C	202	HEM	CMB-C2B-C3B	5.21	129.54	116.53
4	A	202	HEM	CMC-C2C-C3C	5.64	130.61	116.53
4	D	202	HEM	CMC-C2C-C3C	5.97	131.44	116.53
4	B	202	HEM	CAD-C3D-C2D	6.06	130.65	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	202	HEM	1	0
4	B	202	HEM	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	142/143 (99%)	-0.11	4 (2%) 56 57	11, 17, 29, 46	0
1	C	142/143 (99%)	-0.02	4 (2%) 56 57	10, 16, 32, 42	0
2	B	144/146 (98%)	0.36	13 (9%) 12 11	13, 21, 43, 69	0
2	D	145/146 (99%)	-0.09	5 (3%) 49 49	9, 13, 29, 49	0
All	All	573/578 (99%)	0.04	26 (4%) 37 37	9, 17, 36, 69	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	1	VAL	4.6
2	B	144	GLN	4.3
2	D	145	TYR	4.1
2	D	144	GLN	4.0
1	A	51	PRO	4.0
1	A	52	GLY	3.9
2	D	143	LYS	3.5
2	B	84	ILE	3.4
1	C	51	PRO	3.1
1	C	52	GLY	2.9
2	D	2	GLU	2.9
1	A	50	SER	2.8
2	B	83	GLU	2.7
2	B	94	GLU	2.7
1	C	90	TYR	2.7
2	B	1	VAL	2.6
2	B	95	LYS	2.5
2	B	143	LYS	2.3
2	B	90	ILE	2.2
2	B	97[A]	HIS	2.2
1	C	46	TRP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	116	LYS	2.1
2	B	77	ASN	2.1
2	B	86	ALA	2.0
2	B	43	GLY	2.0
2	B	106	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	D	203	6/6	0.94	0.10	2.10	17,20,26,32	0
4	HEM	A	202	43/43	0.98	0.09	0.66	9,13,24,49	0
4	HEM	D	202	43/43	0.98	0.09	0.47	9,13,26,34	0
4	HEM	B	202	43/43	0.95	0.11	0.36	15,21,55,61	0
3	CMO	B	201	2/2	0.97	0.11	0.20	16,16,16,21	0
4	HEM	C	202	43/43	0.97	0.09	0.14	11,15,24,46	0
3	CMO	A	201	2/2	0.99	0.08	-0.16	12,12,12,15	0
3	CMO	D	201	2/2	0.99	0.07	-1.51	12,12,12,14	0
3	CMO	C	201	2/2	0.98	0.05	-1.67	14,14,14,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.