



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:55 AM GMT

PDB ID : 2EVS  
Title : Crystal structure of human Glycolipid Transfer Protein complexed with n-hexyl-beta-D-glucoside  
Authors : Malinina, L.; Malakhova, M.L.; Kanack, A.T.; Abagyan, R.; Brown, R.E.; Patel, D.J.  
Deposited on : 2005-10-31  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	<div> <div>4%</div> <div>80%</div> <div>15%</div> <div>• •</div> </div>
1	E	209	<div> <div>28%</div> <div>73%</div> <div>14%</div> <div>• 10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	HEX	A	301	-	-	-	X
3	HEX	A	311	-	-	-	X
3	HEX	E	312	-	-	-	X
4	D10	A	302	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3496 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycolipid transfer protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	205	Total	C	N	O	S	0	3	0
			1650	1074	270	297	9			
1	E	188	Total	C	N	O	S	0	3	0
			1525	995	245	275	10			

- Molecule 2 is n-hexyl-beta-D-glucoside (three-letter code: GLC) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	E	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is HEXANE (three-letter code: HEX) (formula:  $C_6H_{14}$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C	0	0
			6	6		
3	A	1	Total	C	0	0
			6	6		
3	E	1	Total	C	0	0
			6	6		

- Molecule 4 is DECANE (three-letter code: D10) (formula: C<sub>10</sub>H<sub>22</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C	0	0
			10	10		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	227	Total 227	O 227	0	0
5	E	42	Total 42	O 42	0	0



- Molecule 1: Glycolipid transfer protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.91Å 42.18Å 70.82Å 90.00° 104.57° 90.00°	Depositor
Resolution (Å)	15.00 – 2.20 17.14 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.3 (15.00-2.20) 98.3 (17.14-2.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.21Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.190 , 0.236 0.199 , 0.192	Depositor DCC
$R_{free}$ test set	1180 reflections (5.43%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.3	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 59.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 22940 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3496	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: D10, GLC, HEX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	3/1701 (0.2%)	0.79	1/2305 (0.0%)
1	E	0.51	0/1577	0.58	0/2135
All	All	0.68	3/3278 (0.1%)	0.70	1/4440 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	174	GLU	CG-CD	6.41	1.61	1.51
1	A	174	GLU	CB-CG	6.08	1.63	1.52
1	A	196	GLU	CG-CD	5.53	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	ARG	NE-CZ-NH2	-6.71	116.95	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1650	0	1667	38	2
1	E	1525	0	1539	23	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	12	0	11	0	0
2	E	12	0	11	2	0
3	A	12	0	24	3	0
3	E	6	0	14	2	0
4	A	10	0	22	0	0
5	A	227	0	0	15	3
5	E	42	0	0	1	0
All	All	3496	0	3288	59	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:LYS:HE3	5:A:533:HOH:O	1.29	1.26
1:A:71:GLN:NE2	1:A:201[B]:MET:SD	2.53	0.80
1:A:209:VAL:HG21	3:A:301:HEX:H12	1.66	0.77
1:A:95:MET:SD	5:A:515:HOH:O	2.44	0.75
1:A:174:GLU:HB3	5:A:534:HOH:O	1.89	0.72
1:A:174:GLU:HG3	5:A:344:HOH:O	1.90	0.71
1:A:74:LEU:HB3	1:A:201[B]:MET:HE2	1.73	0.71
1:A:19:GLU:OE1	5:A:503:HOH:O	2.11	0.68
3:A:311:HEX:H12	1:E:209:VAL:HG21	1.76	0.67
1:A:74:LEU:HB2	1:A:201[B]:MET:HE1	1.79	0.64
1:A:71:GLN:NE2	1:A:201[A]:MET:SD	2.73	0.62
1:E:125:ARG:HH21	1:E:153:TYR:HB3	1.66	0.60
1:A:44:PRO:O	3:A:301:HEX:H62	2.04	0.57
1:A:66:LYS:NZ	5:A:400:HOH:O	2.38	0.56
1:A:206:ASN:HB2	5:A:518:HOH:O	2.05	0.55
1:A:124:ILE:HA	1:A:127:ASN:HD22	1.72	0.55
1:A:200:GLN:HG3	5:A:468:HOH:O	2.07	0.54
1:A:128:ALA:HB1	1:A:152:LEU:HD22	1.89	0.54
1:E:41:VAL:O	1:E:42:PHE:HB2	2.08	0.53
1:A:74:LEU:CB	1:A:201[B]:MET:CE	2.86	0.53
1:E:41:VAL:O	1:E:42:PHE:CB	2.55	0.53
1:A:200:GLN:HG2	5:A:509:HOH:O	2.08	0.52
1:A:74:LEU:HB3	1:A:201[B]:MET:CE	2.39	0.52
1:E:124:ILE:HD11	1:E:156:PRO:HD2	1.93	0.50
1:E:18:ILE:HD13	1:E:190:THR:HG23	1.93	0.50
1:A:40:PRO:CB	2:E:310:GLC:H1	2.42	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:THR:N	1:A:44:PRO:CD	2.75	0.49
1:A:41:VAL:HG12	1:E:144:VAL:HG22	1.93	0.49
1:E:40:PRO:O	1:E:43:THR:HG23	2.11	0.49
1:A:74:LEU:CB	1:A:201[B]:MET:HE2	2.41	0.49
1:E:196:GLU:O	1:E:200:GLN:HB2	2.13	0.49
1:A:137:LYS:NZ	5:A:389:HOH:O	2.38	0.49
1:A:148:PHE:O	1:A:152:LEU:HG	2.14	0.48
1:E:41:VAL:O	1:E:42:PHE:CD1	2.67	0.47
1:A:74:LEU:HB2	1:A:201[B]:MET:CE	2.45	0.46
1:A:38:GLY:HA3	5:A:525:HOH:O	2.14	0.46
1:E:107:PHE:CD2	3:E:312:HEX:H52	2.51	0.46
1:E:68:ARG:CZ	5:E:337:HOH:O	2.64	0.46
1:A:71:GLN:NE2	1:A:201[B]:MET:CE	2.80	0.45
1:A:41:VAL:HG12	1:E:144:VAL:CG2	2.46	0.45
1:E:85:TRP:CG	1:E:86:PRO:HA	2.52	0.45
1:E:43:THR:N	1:E:44:PRO:CD	2.81	0.44
1:A:57:LYS:NZ	5:A:448:HOH:O	2.49	0.44
1:E:77:GLU:HA	1:E:80:MET:HE2	1.99	0.44
1:E:39:SER:O	1:E:41:VAL:O	2.36	0.43
1:A:145:GLN:O	1:A:149:GLN:CG	2.67	0.43
1:A:196:GLU:HG3	5:A:462:HOH:O	2.18	0.42
1:A:138:LYS:HD3	5:A:438:HOH:O	2.18	0.42
1:A:19:GLU:HG2	5:A:503:HOH:O	2.18	0.42
1:E:63:ASN:HD22	1:E:66:LYS:HE2	1.85	0.42
1:A:40:PRO:HB3	2:E:310:GLC:H1	2.02	0.42
1:E:107:PHE:CE2	3:E:312:HEX:H52	2.55	0.41
1:A:139:TYR:HE1	1:A:206:ASN:HD22	1.68	0.41
1:E:142:TRP:CE2	1:E:143:ILE:HD12	2.55	0.41
1:E:63:ASN:ND2	1:E:66:LYS:HB2	2.36	0.41
1:A:164:ALA:O	1:A:167:LYS:HD2	2.21	0.40
1:E:81:TYR:CD2	1:E:88:VAL:HG11	2.57	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:ALA:C	5:A:500:HOH:O[1_545]	1.97	0.23
1:A:151:ALA:N	5:A:500:HOH:O[1_545]	1.99	0.21
5:A:398:HOH:O	5:A:455:HOH:O[2_656]	2.17	0.03

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/209 (98%)	199 (97%)	6 (3%)	0	100	100
1	E	187/209 (90%)	175 (94%)	10 (5%)	2 (1%)	17	14
All	All	392/418 (94%)	374 (95%)	16 (4%)	2 (0%)	34	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	42	PHE
1	E	82	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/180 (99%)	167 (94%)	11 (6%)	23	25
1	E	166/180 (92%)	156 (94%)	10 (6%)	24	26
All	All	344/360 (96%)	323 (94%)	21 (6%)	28	26

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	50[A]	SER
1	A	50[B]	SER
1	A	57	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	115	GLU
1	A	130	LYS
1	A	138	LYS
1	A	167	LYS
1	A	171	VAL
1	A	201[A]	MET
1	A	201[B]	MET
1	E	41	VAL
1	E	71	GLN
1	E	84	GLU
1	E	113	ASP
1	E	153	TYR
1	E	174	GLU
1	E	182	LEU
1	E	196	GLU
1	E	201[A]	MET
1	E	201[B]	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	72	ASN
1	A	127	ASN
1	A	170	ASN
1	A	206	ASN
1	E	63	ASN
1	E	127	ASN
1	E	206	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	A	300	3	12,12,12	0.67	0	17,17,17	2.09	5 (29%)
3	HEX	A	301	2	5,5,5	0.33	0	4,4,4	0.31	0
4	D10	A	302	-	9,9,9	0.35	0	8,8,8	0.30	0
3	HEX	A	311	2	5,5,5	0.23	0	4,4,4	0.60	0
2	GLC	E	310	3	12,12,12	0.60	0	17,17,17	2.13	4 (23%)
3	HEX	E	312	-	5,5,5	0.28	0	4,4,4	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	300	3	-	0/2/22/22	0/1/1/1
3	HEX	A	301	2	-	0/3/3/3	0/0/0/0
4	D10	A	302	-	-	0/7/7/7	0/0/0/0
3	HEX	A	311	2	-	0/3/3/3	0/0/0/0
2	GLC	E	310	3	-	0/2/22/22	0/1/1/1
3	HEX	E	312	-	-	0/3/3/3	0/0/0/0

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	GLC	O1-C1-O5	-2.32	103.90	110.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	310	GLC	C1-C2-C3	2.20	113.70	110.43
2	A	300	GLC	O5-C5-C4	2.23	113.87	109.68
2	A	300	GLC	C1-C2-C3	2.36	113.93	110.43
2	E	310	GLC	O5-C5-C4	2.94	115.20	109.68
2	A	300	GLC	O5-C1-C2	4.68	117.27	109.80
2	E	310	GLC	C1-O5-C5	4.88	122.49	113.47
2	E	310	GLC	O5-C1-C2	4.95	117.69	109.80
2	A	300	GLC	C1-O5-C5	4.99	122.69	113.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	HEX	2	0
3	A	311	HEX	1	0
2	E	310	GLC	2	0
3	E	312	HEX	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	205/209 (98%)	0.27	8 (3%) 43 42	27, 35, 53, 62	0
1	E	188/209 (89%)	1.40	58 (30%) 1 0	28, 43, 49, 53	0
All	All	393/418 (94%)	0.81	66 (16%) 2 2	27, 40, 51, 62	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	182	LEU	7.1
1	E	142	TRP	6.2
1	E	164	ALA	5.7
1	E	29	HIS	5.1
1	A	38	GLY	5.0
1	E	200	GLN	4.8
1	E	202	ASN	4.8
1	A	168	GLY	4.7
1	E	74	LEU	4.7
1	E	22	PRO	4.6
1	E	171	VAL	4.4
1	A	153	TYR	4.3
1	E	73	ILE	4.3
1	E	172	THR	4.3
1	A	169	GLN	4.3
1	E	165	LEU	4.2
1	E	65	ALA	4.2
1	E	25	GLU	4.1
1	E	91	THR	3.9
1	A	170	ASN	3.7
1	E	68	ARG	3.5
1	E	84	GLU	3.4
1	E	112	CYS	3.4
1	E	62	THR	3.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	175	GLU	3.3
1	E	66	LYS	3.3
1	E	189	ALA	3.2
1	E	38	GLY	3.2
1	E	118	GLU	3.2
1	E	32	PRO	3.1
1	E	180	ILE	3.0
1	E	178	GLU	3.0
1	E	161	PHE	2.9
1	E	19	GLU	2.9
1	E	201[A]	MET	2.9
1	E	79	GLU	2.8
1	E	119	ASN	2.8
1	E	67	PHE	2.6
1	A	157	TYR	2.6
1	E	199	THR	2.6
1	E	85	TRP	2.5
1	E	83	ALA	2.4
1	E	205	LEU	2.4
1	E	198	TYR	2.4
1	E	157	TYR	2.4
1	E	176	CYS	2.4
1	E	206	ASN	2.4
1	E	193	VAL	2.4
1	E	203	ALA	2.3
1	E	153	TYR	2.3
1	E	187	TYR	2.3
1	A	174	GLU	2.3
1	E	40	PRO	2.3
1	E	134[A]	MET	2.3
1	E	39	SER	2.2
1	E	60	TYR	2.2
1	E	108	LEU	2.1
1	E	113	ASP	2.1
1	E	33	PHE	2.1
1	E	36	CYS	2.1
1	E	191	ILE	2.1
1	E	63	ASN	2.0
1	E	26	ALA	2.0
1	E	139	TYR	2.0
1	A	124	ILE	2.0
1	E	138	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	D10	A	302	10/10	0.83	0.29	3.18	45,46,48,48	0
3	HEX	A	301	6/6	0.86	0.27	2.88	51,52,53,53	0
3	HEX	E	312	6/6	0.75	0.37	2.24	52,53,53,53	0
3	HEX	A	311	6/6	0.89	0.21	2.08	41,42,44,45	0
2	GLC	E	310	12/12	0.92	0.14	-0.21	34,38,40,40	0
2	GLC	A	300	12/12	0.94	0.12	-0.77	40,45,48,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.