



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:56 AM GMT

PDB ID : 2EWW
Title : Crystal Structure of the Pilus Retraction Motor PilT and Bound ATP
Authors : Satyshur, K.A.; Forest, K.T.
Deposited on : 2005-11-07
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	<div> <div></div> <div>64%</div> <div>24%</div> <div>• 8%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called twitching motility protein PilT.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	343	2717	1731	471	505	1	9	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	68	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	136	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	156	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	218	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	318	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	327	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	330	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	349	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	361	MSE	MET	MODIFIED RESIDUE	GB 15606134
A	367	HIS	-	EXPRESSION TAG	GB 15606134
A	368	HIS	-	EXPRESSION TAG	GB 15606134
A	369	HIS	-	EXPRESSION TAG	GB 15606134
A	370	HIS	-	EXPRESSION TAG	GB 15606134
A	371	HIS	-	EXPRESSION TAG	GB 15606134
A	372	HIS	-	EXPRESSION TAG	GB 15606134

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total O 2 2	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: twitching motility protein PilT



4 Data and refinement statistics

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, α , β , γ	107.38 Å 107.38 Å 68.86 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.47 – 3.20 19.47 – 3.20	Depositor EDS
% Data completeness (in resolution range)	97.3 (19.47-3.20) 97.3 (19.47-3.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.22 (at 3.22 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.239 , 0.310 0.245 , 0.226	Depositor DCC
R_{free} test set	362 reflections (5.19%)	DCC
Wilson B-factor (Å ²)	79.4	Xtriage
Anisotropy	0.004	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 37.3	EDS
Estimated twinning fraction	0.049 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 7343 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2750	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2753	0.56	0/3702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2717	0	2798	51	0
2	A	31	0	12	0	0
3	A	2	0	0	0	0
All	All	2750	0	2810	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349:MSE:HA	1:A:349:MSE:HE2	1.68	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:ASN:HA	1:A:334:LEU:HD12	1.71	0.72
1:A:52:PHE:HB3	1:A:53:PRO:HD3	1.73	0.70
1:A:166:TYR:H	1:A:184:SER:HB3	1.60	0.66
1:A:137:GLY:HA2	1:A:272:ILE:HG13	1.80	0.64
1:A:358:LEU:C	1:A:360:ARG:H	2.03	0.62
1:A:69:SER:O	1:A:73:ARG:HG3	1.99	0.62
1:A:41:ARG:HD3	1:A:44:GLY:HA2	1.83	0.60
1:A:37:PRO:HG2	1:A:48:PHE:HB3	1.83	0.60
1:A:298:PRO:HB2	1:A:303:ARG:HE	1.66	0.58
1:A:170:THR:HG22	1:A:214:PHE:HB3	1.91	0.52
1:A:284:LYS:HG2	1:A:285:ILE:N	2.25	0.52
1:A:259:LEU:HA	1:A:262:GLN:HB2	1.92	0.51
1:A:142:THR:HG22	1:A:241:LEU:HB2	1.92	0.51
1:A:294:GLU:HB2	1:A:330:MSE:HG2	1.93	0.50
1:A:361:MSE:HE2	1:A:361:MSE:HA	1.93	0.50
1:A:270:SER:O	1:A:303:ARG:HD3	2.12	0.50
1:A:138:LEU:HD22	1:A:140:LEU:HG	1.93	0.49
1:A:220:ASP:OD2	1:A:222:GLU:HB3	2.13	0.48
1:A:64:ALA:O	1:A:68:MSE:HG3	2.14	0.48
1:A:259:LEU:HA	1:A:262:GLN:CB	2.43	0.48
1:A:34:ALA:O	1:A:56:THR:O	2.31	0.48
1:A:195:THR:HG21	1:A:201:ALA:HB2	1.96	0.47
1:A:293:TYR:H	1:A:333:THR:HG21	1.78	0.47
1:A:335:TYR:HB2	1:A:358:LEU:HD11	1.96	0.47
1:A:158:ASP:O	1:A:162:GLN:HG3	2.15	0.47
1:A:230:ALA:HA	1:A:233:THR:HG22	1.97	0.47
1:A:284:LYS:HG2	1:A:285:ILE:H	1.80	0.46
1:A:57:PRO:C	1:A:59:ASP:H	2.18	0.46
1:A:153:ILE:HG21	1:A:214:PHE:CZ	2.51	0.46
1:A:273:LEU:HD11	1:A:276:ILE:HG13	1.98	0.45
1:A:216:GLY:O	1:A:240:THR:OG1	2.34	0.45
1:A:65:TYR:C	1:A:67:VAL:H	2.20	0.45
1:A:57:PRO:C	1:A:59:ASP:N	2.70	0.45
1:A:91:VAL:HG13	1:A:92:GLY:H	1.82	0.44
1:A:358:LEU:C	1:A:360:ARG:N	2.71	0.44
1:A:166:TYR:N	1:A:184:SER:HB3	2.31	0.43
1:A:349:MSE:HA	1:A:349:MSE:CE	2.43	0.43
1:A:302:ILE:HG22	1:A:306:ILE:HD11	2.01	0.43
1:A:122:LEU:HD23	1:A:124:LEU:HD23	2.00	0.43
1:A:50:LYS:O	1:A:51:ASP:C	2.57	0.43
1:A:126:ASP:OD2	1:A:126:ASP:N	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:VAL:O	1:A:240:THR:HA	2.20	0.42
1:A:196:LYS:HD2	1:A:196:LYS:N	2.34	0.42
1:A:122:LEU:HD21	1:A:152:THR:HA	2.03	0.41
1:A:49:LEU:O	1:A:49:LEU:HD12	2.21	0.41
1:A:310:LYS:O	1:A:314:VAL:HG23	2.20	0.41
1:A:264:GLN:HA	1:A:267:ILE:HD12	2.02	0.41
1:A:153:ILE:HG21	1:A:214:PHE:CE1	2.56	0.40
1:A:59:ASP:HB3	1:A:62:LYS:HG3	2.02	0.40
1:A:128:VAL:HG13	1:A:277:ILE:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/372 (91%)	306 (90%)	26 (8%)	7 (2%)	9	46

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	51	ASP
1	A	13	LEU
1	A	52	PHE
1	A	58	GLU
1	A	122	LEU
1	A	176	GLU
1	A	286	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	297/311 (96%)	261 (88%)	36 (12%)	6 28

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ILE
1	A	30	ILE
1	A	51	ASP
1	A	56	THR
1	A	62	LYS
1	A	67	VAL
1	A	74	GLN
1	A	75	LYS
1	A	82	VAL
1	A	83	ASP
1	A	95	ARG
1	A	101	GLN
1	A	102	ARG
1	A	120	LYS
1	A	126	ASP
1	A	135	LYS
1	A	138	LEU
1	A	185	ILE
1	A	195	THR
1	A	196	LYS
1	A	202	LEU
1	A	215	VAL
1	A	219	ARG
1	A	220	ASP
1	A	236	LEU
1	A	251	HIS
1	A	252	ARG
1	A	259	LEU
1	A	261	GLN
1	A	262	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	266	ARG
1	A	290	VAL
1	A	313	GLN
1	A	315	TYR
1	A	330	MSE
1	A	359	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	GLN
1	A	133	HIS
1	A	187	ASN
1	A	244	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ATP	A	500	-	24,33,33	1.02	1 (4%)	31,52,52	1.74	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	500	-	-	0/18/38/38	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	ATP	C5-C4	3.32	1.48	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	ATP	N3-C2-N1	-6.24	124.12	128.89
2	A	500	ATP	PB-O3B-PG	-3.00	122.62	132.67
2	A	500	ATP	C4-C5-N7	-2.95	106.76	109.48
2	A	500	ATP	PA-O3A-PB	-2.88	124.64	132.73
2	A	500	ATP	C2'-C1'-N9	-2.32	110.74	114.29
2	A	500	ATP	C4'-O4'-C1'	2.15	112.08	109.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/372 (89%)	-0.06	4 (1%) 81 69	39, 64, 106, 123	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	51	ASP	2.2
1	A	317	LEU	2.2
1	A	332	GLN	2.2
1	A	261	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ATP	A	500	31/31	0.93	0.18	-0.49	61,62,68,69	0

6.5 Other polymers

There are no such residues in this entry.