



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 4, 2016 – 11:06 AM EDT

PDB ID : 5EWL
Title : CRYSTAL STRUCTURE OF AMINO TERMINAL DOMAINS OF THE
NMDA RECEPTOR SUBUNIT GLUN1 AND GLUN2B IN COMPLEX
WITH MK-22
Authors : Pandit, J.
Deposited on : 2015-11-20
Resolution : 2.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1992 (3.00-2.96)
Clashscore	102246	2349 (3.00-2.96)
Ramachandran outliers	100387	2274 (3.00-2.96)
Sidechain outliers	100360	2277 (3.00-2.96)
RSRZ outliers	91569	2007 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	<div> <div>0%</div> <div> <div>78%</div> <div>12%</div> <div>• 8%</div> </div> </div>
1	C	390	<div> <div>5%</div> <div> <div>80%</div> <div>10%</div> <div>• 9%</div> </div> </div>
2	B	364	<div> <div>2%</div> <div> <div>81%</div> <div>13%</div> <div>• •</div> </div> </div>
2	D	364	<div> <div>6%</div> <div> <div>77%</div> <div>17%</div> <div>• •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NA	A	501	-	-	-	X
3	NA	C	501	-	-	-	X
4	NAG	A	502	-	-	-	X
4	NAG	C	502	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NMDA glutamate receptor subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	357	Total	C	N	O	S	0	0	0
			2752	1753	477	511	11			
1	C	356	Total	C	N	O	S	0	0	0
			2737	1741	479	507	10			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	GLN	ASN	engineered mutation	UNP Q91977
A	371	GLN	ASN	engineered mutation	UNP Q91977
A	409	LEU	-	expression tag	UNP Q91977
A	410	VAL	-	expression tag	UNP Q91977
A	411	PRO	-	expression tag	UNP Q91977
A	412	ARG	-	expression tag	UNP Q91977
C	61	GLN	ASN	engineered mutation	UNP Q91977
C	371	GLN	ASN	engineered mutation	UNP Q91977
C	409	LEU	-	expression tag	UNP Q91977
C	410	VAL	-	expression tag	UNP Q91977
C	411	PRO	-	expression tag	UNP Q91977
C	412	ARG	-	expression tag	UNP Q91977

- Molecule 2 is a protein called Glutamate receptor ionotropic, NMDA 2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	352	Total	C	N	O	S	0	0	0
			2739	1766	432	526	15			
2	D	355	Total	C	N	O	S	0	0	0
			2789	1797	441	535	16			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	348	ASP	ASN	engineered mutation	UNP Q13224
D	348	ASP	ASN	engineered mutation	UNP Q13224

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Na 2 2	0	0
3	A	3	Total Na 3 3	0	0
3	C	1	Total Na 1 1	0	0

- Molecule 4 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 14 8 1 5	0	0
4	A	1	Total C N O 14 8 1 5	0	0
4	A	1	Total C N O 14 8 1 5	0	0
4	B	1	Total C N O 14 8 1 5	0	0
4	B	1	Total C N O 14 8 1 5	0	0
4	C	1	Total C N O 14 8 1 5	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			11	6	5		
6	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 7 is {N}-(1 {S},3 {S})-3-[3-[(4-methylphenyl)methyl]-1,2,4-oxadiazol-5-yl]cyclopentyl]-1 {H}-pyrazolo[3,4-d]pyrimidin-4-amine (three-letter code: 5SL) (formula: C₂₀H₂₁N₇O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			28	20	7	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	1	Total	C	N	O	0	0
			28	20	7	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	99	Total	O	0	0
			99	99		
8	B	83	Total	O	0	0
			83	83		
8	C	39	Total	O	0	0
			39	39		
8	D	48	Total	O	0	0
			48	48		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NMDA glutamate receptor subunit



• Molecule 1: NMDA glutamate receptor subunit





• Molecule 2: Glutamate receptor ionotropic, NMDA 2B





● Molecule 2: Glutamate receptor ionotropic, NMDA 2B



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	268.88Å 59.94Å 144.45Å 90.00° 116.51° 90.00°	Depositor
Resolution (Å)	30.04 – 2.98 29.97 – 2.98	Depositor EDS
% Data completeness (in resolution range)	95.8 (30.04-2.98) 95.8 (29.97-2.98)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 3.00Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.162 , 0.217 0.166 , 0.219	Depositor DCC
R_{free} test set	2068 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	68.8	Xtriage
Anisotropy	0.215	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 80.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 40941 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11493	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, BMA, NAG, 5SL, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/2809	0.72	0/3819
1	C	0.49	0/2792	0.69	0/3794
2	B	0.53	0/2801	0.73	0/3814
2	D	0.50	0/2853	0.72	0/3879
All	All	0.52	0/11255	0.72	0/15306

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2752	0	2717	24	0
1	C	2737	0	2705	17	0
2	B	2739	0	2649	27	0
2	D	2789	0	2707	34	0
3	A	3	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
4	A	42	0	37	0	0
4	B	28	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	14	0	13	0	0
4	D	28	0	26	0	0
5	A	11	0	8	0	0
6	A	22	0	20	0	0
7	A	28	0	0	1	0
7	C	28	0	0	0	0
8	A	99	0	0	1	0
8	B	83	0	0	0	0
8	C	39	0	0	0	0
8	D	48	0	0	0	0
All	All	11493	0	10908	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:173:THR:HG22	2:D:231:TYR:HB3	1.59	0.84
2:D:344:PHE:HB2	2:D:349:LEU:HD11	1.60	0.82
2:B:173:THR:HG22	2:B:231:TYR:HB3	1.61	0.80
2:D:85:ILE:HG22	2:D:89:MET:HE2	1.64	0.78
2:B:33:PRO:HG2	2:B:64:VAL:HG23	1.70	0.72
2:B:109:ALA:HB1	2:B:134:MET:HG3	1.76	0.67
1:C:27:VAL:HG23	1:C:60:LEU:HD23	1.76	0.67
1:A:376:ASN:HD22	1:A:378:GLN:HE21	1.43	0.66
2:D:33:PRO:HG2	2:D:64:VAL:HG23	1.77	0.66
1:A:57:LYS:HB2	8:A:670:HOH:O	1.95	0.66
1:C:167:VAL:HG12	1:C:243:SER:HB3	1.78	0.65
1:A:167:VAL:HG12	1:A:243:SER:HB3	1.78	0.64
2:B:158:LEU:HA	2:B:161:MET:HE3	1.83	0.61
2:D:149:SER:H	2:D:152:GLN:HE21	1.47	0.60
2:D:158:LEU:HA	2:D:161:MET:HE3	1.83	0.60
1:A:376:ASN:HD21	1:A:401:ILE:H	1.50	0.60
2:B:327:LYS:O	2:B:331:GLN:HG3	2.02	0.60
1:C:144:TYR:CE2	1:C:273:ARG:HD2	2.37	0.60
2:B:149:SER:H	2:B:152:GLN:HE21	1.49	0.59
2:D:169:PHE:HZ	2:D:190:ILE:HD11	1.65	0.59
1:C:287:ILE:HG22	1:C:377:LEU:HB2	1.84	0.58
2:D:234:LYS:O	2:D:238:THR:HG23	2.04	0.57
1:C:262:THR:HB	1:C:284:PRO:HB3	1.86	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:258:VAL:HG13	2:B:262:VAL:HB	1.87	0.56
2:B:234:LYS:O	2:B:238:THR:HG23	2.05	0.56
2:D:258:VAL:HG13	2:D:262:VAL:HB	1.89	0.54
1:C:40:GLN:HG3	1:C:43:ARG:HH22	1.73	0.54
2:D:230:LEU:O	2:D:259:PRO:HD3	2.08	0.54
2:D:362:LEU:HB2	2:D:379:TRP:HB3	1.90	0.54
2:B:230:LEU:O	2:B:259:PRO:HD3	2.08	0.53
2:D:85:ILE:HG22	2:D:89:MET:CE	2.37	0.53
2:D:283:ASP:HA	2:D:285:TRP:HE3	1.73	0.53
1:A:402:TRP:HD1	1:A:406:GLU:HA	1.73	0.53
2:B:250:THR:HB	2:B:274:PRO:HB3	1.92	0.52
1:A:322:ILE:HG23	1:A:338:THR:HG21	1.92	0.51
2:B:283:ASP:OD1	2:B:285:TRP:HB2	2.10	0.51
1:C:309:VAL:O	1:C:313:ILE:HG12	2.11	0.51
2:D:250:THR:HB	2:D:274:PRO:HB3	1.91	0.51
1:A:262:THR:HB	1:A:284:PRO:HB3	1.92	0.50
2:D:149:SER:H	2:D:152:GLN:NE2	2.08	0.50
2:B:193:SER:C	2:B:195:VAL:H	2.13	0.50
1:A:135:LEU:HD22	7:A:508:5SL:C15	2.42	0.50
2:D:318:LYS:H	2:D:318:LYS:CD	2.25	0.50
2:D:169:PHE:CZ	2:D:190:ILE:HD11	2.45	0.50
2:B:129:GLY:HA2	2:B:132:MET:SD	2.51	0.49
2:D:299:ILE:HA	2:D:342:VAL:HG11	1.94	0.49
1:A:229:LEU:HD12	1:A:255:SER:HB2	1.93	0.49
1:C:144:TYR:CZ	1:C:273:ARG:HD2	2.48	0.48
2:B:149:SER:H	2:B:152:GLN:NE2	2.10	0.48
2:B:329:ILE:HG22	2:D:308:LEU:HD11	1.96	0.48
2:D:325:HIS:CD2	2:D:326:GLU:HG3	2.48	0.48
1:C:157:LEU:HD23	1:C:158:PHE:CE1	2.49	0.47
2:D:283:ASP:HA	2:D:285:TRP:CE3	2.49	0.47
1:A:226:THR:HG22	1:A:255:SER:HB3	1.96	0.47
1:A:33:LEU:O	1:A:66:THR:HA	2.15	0.47
2:B:318:LYS:H	2:B:318:LYS:CD	2.27	0.47
2:D:150:ILE:HD13	2:D:178:GLY:HA3	1.96	0.47
1:C:147:GLN:HE22	1:C:272:GLU:H	1.63	0.47
2:D:89:MET:HE3	2:D:119:THR:HG21	1.97	0.46
1:C:33:LEU:O	1:C:66:THR:HA	2.16	0.46
1:A:27:VAL:HG13	1:A:88:TYR:CD1	2.51	0.46
1:A:157:LEU:HD23	1:A:158:PHE:CE1	2.50	0.46
2:D:283:ASP:HB3	2:D:363:VAL:HG23	1.97	0.46
2:D:105:GLN:HB3	2:D:108:ILE:HD12	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:308:VAL:HG21	1:C:359:ILE:HG21	1.98	0.45
1:A:242:LEU:HB3	1:A:270:VAL:HG23	1.98	0.45
1:C:242:LEU:HB3	1:C:270:VAL:HG23	1.99	0.45
1:A:135:LEU:H	1:A:135:LEU:HD23	1.81	0.45
2:B:216:ILE:HG22	2:B:243:VAL:HG11	1.99	0.45
2:D:92:ARG:O	2:D:92:ARG:HD3	2.16	0.45
1:A:308:VAL:HG21	1:A:359:ILE:HG21	1.98	0.45
2:D:144:PHE:HE1	2:D:340:ILE:HD12	1.81	0.45
1:A:133:ILE:HG22	1:A:134:HIS:CD2	2.52	0.44
2:B:344:PHE:HB3	2:B:349:LEU:HD11	1.99	0.44
2:B:220:LEU:HD12	2:B:243:VAL:HG12	2.00	0.44
2:B:109:ALA:CB	2:B:134:MET:HG3	2.45	0.43
1:A:41:ILE:HG22	1:A:302:ILE:HG21	2.01	0.43
2:B:50:ILE:HD11	2:B:289:LEU:HD23	2.00	0.43
2:D:247:VAL:O	2:D:247:VAL:HG12	2.18	0.42
2:D:318:LYS:H	2:D:318:LYS:HD3	1.83	0.42
2:B:161:MET:HE2	2:B:229:LEU:HD21	2.01	0.42
1:A:309:VAL:O	1:A:313:ILE:HG12	2.20	0.42
1:C:167:VAL:O	1:C:217:GLN:HA	2.20	0.42
2:D:124:LEU:HD23	2:D:144:PHE:HB2	2.01	0.42
1:A:382:LEU:HD23	1:A:382:LEU:HA	1.95	0.42
1:C:47:ASN:HA	1:C:50:ASN:HD22	1.85	0.42
1:C:147:GLN:HE22	1:C:272:GLU:N	2.18	0.41
1:A:167:VAL:O	1:A:217:GLN:HA	2.21	0.41
2:B:299:ILE:HA	2:B:342:VAL:HG11	2.03	0.41
2:D:216:ILE:HG22	2:D:243:VAL:HG11	2.02	0.41
2:D:95:GLN:NE2	2:D:314:ILE:HD12	2.35	0.41
2:B:269:VAL:HG11	2:B:367:LEU:HD11	2.03	0.41
1:A:229:LEU:CD1	1:A:255:SER:HB2	2.51	0.40
1:A:42:PHE:CE2	1:A:62:ALA:HB1	2.56	0.40
2:B:121:THR:OG1	2:B:318:LYS:HE3	2.21	0.40
2:D:121:THR:OG1	2:D:318:LYS:HE3	2.20	0.40
1:A:350:LYS:HG2	1:A:360:GLU:HG2	2.03	0.40
1:C:30:GLY:HA2	1:C:63:THR:O	2.21	0.40
2:B:318:LYS:H	2:B:318:LYS:HD3	1.86	0.40
2:B:92:ARG:O	2:B:92:ARG:HD3	2.22	0.40
2:D:260:SER:HA	2:D:280:VAL:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/390 (90%)	339 (97%)	12 (3%)	0	100	100
1	C	350/390 (90%)	338 (97%)	12 (3%)	0	100	100
2	B	348/364 (96%)	320 (92%)	27 (8%)	1 (0%)	46	82
2	D	349/364 (96%)	316 (90%)	30 (9%)	3 (1%)	21	63
All	All	1398/1508 (93%)	1313 (94%)	81 (6%)	4 (0%)	46	82

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	283	ASP
2	D	283	ASP
2	D	62	LEU
2	D	327	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/336 (87%)	274 (94%)	18 (6%)	23	59
1	C	289/336 (86%)	271 (94%)	18 (6%)	23	59
2	B	297/326 (91%)	276 (93%)	21 (7%)	18	53
2	D	307/326 (94%)	275 (90%)	32 (10%)	9	31
All	All	1185/1324 (90%)	1096 (92%)	89 (8%)	17	50

All (89) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	ILE
1	A	136	SER
1	A	182	THR
1	A	223	LYS
1	A	226	THR
1	A	230	LEU
1	A	255	SER
1	A	270	VAL
1	A	285	ASP
1	A	320	GLU
1	A	335	ILE
1	A	353	ASP
1	A	367	ARG
1	A	380	ARG
1	A	382	LEU
1	A	387	ILE
1	A	392	TYR
1	A	399	LYS
2	B	64	VAL
2	B	65	VAL
2	B	69	GLU
2	B	90	SER
2	B	91	ASP
2	B	92	ARG
2	B	152	GLN
2	B	190	ILE
2	B	246	SER
2	B	258	VAL
2	B	260	SER
2	B	308	LEU
2	B	310	GLU
2	B	318	LYS
2	B	329	ILE
2	B	335	LEU
2	B	349	LEU
2	B	357	GLN
2	B	378	LYS
2	B	380	LYS
2	B	381	ASP
1	C	27	VAL
1	C	54	PHE
1	C	55	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	58	ILE
1	C	127	ILE
1	C	135	LEU
1	C	136	SER
1	C	229	LEU
1	C	270	VAL
1	C	273	ARG
1	C	285	ASP
1	C	335	ILE
1	C	353	ASP
1	C	364	ASP
1	C	367	ARG
1	C	377	LEU
1	C	378	GLN
1	C	387	ILE
2	D	34	SER
2	D	59	PHE
2	D	64	VAL
2	D	65	VAL
2	D	69	GLU
2	D	90	SER
2	D	92	ARG
2	D	103	THR
2	D	145	GLN
2	D	152	GLN
2	D	180	GLN
2	D	193	SER
2	D	194	PHE
2	D	245	ASN
2	D	246	SER
2	D	258	VAL
2	D	260	SER
2	D	280	VAL
2	D	310	GLU
2	D	318	LYS
2	D	326	GLU
2	D	329	ILE
2	D	335	LEU
2	D	337	ARG
2	D	340	ILE
2	D	349	LEU
2	D	352	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	357	GLN
2	D	369	LYS
2	D	381	ASP
2	D	382	LYS
2	D	389	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	376	ASN
1	A	384	GLN
2	B	110	GLN
2	B	118	GLN
2	B	152	GLN
2	B	311	HIS
1	C	28	ASN
1	C	147	GLN
1	C	376	ASN
1	C	384	GLN
2	D	95	GLN
2	D	118	GLN
2	D	152	GLN
2	D	311	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 6 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	502	1	14,14,15	0.22	0	15,19,21	1.31	2 (13%)
4	NAG	A	503	1,4	14,14,15	0.26	0	15,19,21	0.33	0
4	NAG	A	504	5,4	14,14,15	0.26	0	15,19,21	0.51	0
5	BMA	A	505	4,6	11,11,12	0.31	0	15,15,17	0.79	0
6	MAN	A	506	5	11,11,12	0.42	0	15,15,17	0.88	1 (6%)
6	MAN	A	507	5	11,11,12	0.41	0	15,15,17	1.10	0
7	5SL	A	508	-	26,32,32	0.71	0	24,45,45	0.87	1 (4%)
4	NAG	B	501	2	14,14,15	0.28	0	15,19,21	0.47	0
4	NAG	B	502	2	14,14,15	0.30	0	15,19,21	0.71	1 (6%)
4	NAG	C	502	1	14,14,15	0.39	0	15,19,21	0.49	0
7	5SL	C	503	-	26,32,32	0.76	0	24,45,45	0.87	1 (4%)
4	NAG	D	501	2	14,14,15	0.37	0	15,19,21	0.47	0
4	NAG	D	502	2	14,14,15	0.31	0	15,19,21	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	502	1	-	0/6/23/26	0/1/1/1
4	NAG	A	503	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	504	5,4	-	0/6/23/26	0/1/1/1
5	BMA	A	505	4,6	-	0/2/19/22	0/1/1/1
6	MAN	A	506	5	-	0/2/19/22	0/1/1/1
6	MAN	A	507	5	-	0/2/19/22	0/1/1/1
7	5SL	A	508	-	-	0/8/21/21	0/4/5/5
4	NAG	B	501	2	-	0/6/23/26	0/1/1/1
4	NAG	B	502	2	-	0/6/23/26	0/1/1/1
4	NAG	C	502	1	-	0/6/23/26	0/1/1/1
7	5SL	C	503	-	-	0/8/21/21	0/4/5/5
4	NAG	D	501	2	-	0/6/23/26	0/1/1/1
4	NAG	D	502	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	B	502	NAG	C1-O5-C5	2.41	115.69	112.14
6	A	506	MAN	C1-O5-C5	2.44	115.72	112.14
4	A	502	NAG	C1-O5-C5	2.49	115.80	112.14
7	C	503	5SL	C11-C10-N1	2.90	129.81	124.27
7	A	508	5SL	C11-C10-N1	2.96	129.94	124.27
4	A	502	NAG	C3-C4-C5	3.21	115.95	110.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	508	5SL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/390 (91%)	-0.27	3 (0%) 87 72	35, 52, 84, 109	0
1	C	356/390 (91%)	0.11	20 (5%) 28 14	56, 78, 110, 137	0
2	B	352/364 (96%)	-0.03	6 (1%) 73 51	36, 66, 98, 119	0
2	D	355/364 (97%)	0.11	22 (6%) 24 12	53, 78, 119, 136	0
All	All	1420/1508 (94%)	-0.02	51 (3%) 46 27	35, 71, 108, 137	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	407	THR	5.4
2	D	193	SER	4.8
1	C	56	ARG	4.1
2	B	32	PRO	4.0
1	C	397	ASP	4.0
2	D	381	ASP	4.0
2	D	50	ILE	3.7
1	C	406	GLU	3.6
2	D	192	ASN	3.4
1	C	57	LYS	3.3
1	C	405	GLY	3.2
2	D	195	VAL	3.2
1	C	353	ASP	3.2
1	C	220	PRO	3.1
1	C	54	PHE	3.1
1	C	394	ILE	3.1
2	B	380	LYS	3.0
2	D	44	THR	3.0
2	D	380	LYS	3.0
2	D	344	PHE	3.0
2	D	213	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	405	GLY	2.7
1	C	321	ASN	2.7
2	D	325	HIS	2.7
1	C	404	GLY	2.7
2	B	382	LYS	2.7
1	C	264	ALA	2.7
2	D	194	PHE	2.6
2	D	372	LYS	2.5
1	C	259	LEU	2.5
1	A	407	THR	2.4
1	C	23	ASP	2.4
2	B	49	ALA	2.4
1	C	210	PRO	2.4
1	C	209	GLY	2.4
2	D	382	LYS	2.3
1	C	169	ASP	2.3
2	D	323	ASN	2.3
2	D	60	HIS	2.2
2	D	32	PRO	2.2
1	A	320	GLU	2.2
2	D	279	SER	2.2
1	C	69	PRO	2.2
2	D	285	TRP	2.2
2	D	258	VAL	2.2
2	B	285	TRP	2.1
2	D	180	GLN	2.1
2	D	191	GLU	2.1
2	B	325	HIS	2.0
2	D	370	GLU	2.0
1	C	174	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NA	A	501	1/1	0.89	0.40	6.60	66,66,66,66	0
4	NAG	C	502	14/15	0.84	0.39	3.91	72,80,89,94	0
3	NA	C	501	1/1	0.96	0.29	3.90	70,70,70,70	0
4	NAG	A	502	14/15	0.95	0.28	2.58	60,68,75,79	0
4	NAG	A	503	14/15	0.97	0.23	1.10	60,70,78,80	0
6	MAN	A	506	11/12	0.96	0.21	0.98	59,62,68,69	0
3	NA	A	509	1/1	0.88	0.18	0.98	49,49,49,49	0
7	5SL	A	508	28/28	0.98	0.14	-0.64	33,42,55,58	0
7	5SL	C	503	28/28	0.98	0.14	-0.85	52,60,64,65	0
3	NA	B	504	1/1	0.94	0.10	-1.47	60,60,60,60	0
3	NA	A	510	1/1	0.95	0.06	-1.65	43,43,43,43	0
3	NA	B	503	1/1	0.98	0.07	-2.73	67,67,67,67	0
4	NAG	B	502	14/15	0.44	0.41	-	146,150,157,160	0
4	NAG	D	501	14/15	0.64	0.46	-	145,147,152,154	0
6	MAN	A	507	11/12	0.89	0.21	-	76,80,86,88	0
4	NAG	D	502	14/15	0.65	0.47	-	165,171,173,175	0
4	NAG	B	501	14/15	0.74	0.50	-	127,135,141,141	0
4	NAG	A	504	14/15	0.97	0.29	-	64,68,75,75	0
5	BMA	A	505	11/12	0.96	0.22	-	68,73,82,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.