



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 11:07 PM GMT

PDB ID : 5EXE
Title : Crystal structure of oxalate oxidoreductase from Moorella thermoacetica bound with carboxy-TPP adduct
Authors : Gibson, M.I.; Chen, P.Y.-T.; Drennan, C.L.
Deposited on : 2015-11-23
Resolution : 1.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6965 (1.90-1.86)
Clashscore	102246	7778 (1.90-1.86)
Ramachandran outliers	100387	7691 (1.90-1.86)
Sidechain outliers	100360	7692 (1.90-1.86)
RSRZ outliers	91569	6979 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	<div> <div>4%</div> <div>86%</div> <div>13%</div> <div>•</div> </div>
1	D	395	<div> <div>3%</div> <div>87%</div> <div>12%</div> <div>•</div> </div>
2	B	315	<div> <div>21%</div> <div>76%</div> <div>22%</div> <div>•</div> </div>
2	E	315	<div> <div>26%</div> <div>78%</div> <div>14%</div> <div>8%</div> <div>•</div> </div>
3	C	314	<div> <div>%</div> <div>93%</div> <div>6%</div> <div>•</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	314	 90% 10%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 18353 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxalate oxidoreductase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	23	0
			3237	2059	552	611	15			
1	D	394	Total	C	N	O	S	0	24	0
			3245	2064	555	611	15			

- Molecule 2 is a protein called Oxalate oxidoreductase subunit delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	309	Total	C	N	O	S	0	0	0
			2331	1468	394	453	16			
2	E	290	Total	C	N	O	S	0	0	0
			2194	1383	371	425	15			

- Molecule 3 is a protein called Oxalate oxidoreductase subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	314	Total	C	N	O	S	0	0	0
			2406	1542	412	435	17			
3	F	313	Total	C	N	O	S	0	0	0
			2398	1538	410	433	17			

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	Fe	S	0	0
			8	4	4		
4	B	1	Total	Fe	S	0	0
			8	4	4		
4	C	1	Total	Fe	S	0	0
			8	4	4		
4	E	1	Total	Fe	S	0	0
			8	4	4		
4	E	1	Total	Fe	S	0	0
			8	4	4		
4	F	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 5 is [2-[3-[(4-azanyl-2-methyl-pyrimidin-5-yl)methyl]-2-carboxy-4-methyl-1,3-thiazol-3-ium-5-yl]ethoxy-oxidanyl-phosphoryl] hydrogen phosphate (three-letter code: 5SR) (formula: C₁₃H₁₈N₄O₉P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	P	S	
			29	13	4	9	2	1	0
5	F	1	Total	C	N	O	P	S	
			29	13	4	9	2	1	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	2	Total	Mg		
			2	2	0	0
6	F	1	Total	Mg		
			1	1	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	F	1	Total	Na		
			1	1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	406	Total	O		
			406	406	0	0
8	B	447	Total	O		
			447	447	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	380	Total 380	O 380	0	0
8	D	471	Total 471	O 471	0	0
8	E	296	Total 296	O 296	0	0
8	F	432	Total 432	O 432	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Oxalate oxidoreductase subunit alpha




- Molecule 1: Oxalate oxidoreductase subunit alpha




- Molecule 2: Oxalate oxidoreductase subunit delta




- Molecule 2: Oxalate oxidoreductase subunit delta



- Molecule 3: Oxalate oxidoreductase subunit beta



- Molecule 3: Oxalate oxidoreductase subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.65Å 144.13Å 161.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.70 – 1.88 48.70 – 1.88	Depositor EDS
% Data completeness (in resolution range)	92.9 (48.70-1.88) 92.9 (48.70-1.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 1.88Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.181 , 0.213 0.188 , 0.217	Depositor DCC
R_{free} test set	10068 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	22.5	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 200283 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18353	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SF4, MG, 5SR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3316	0.48	0/4500
1	D	0.31	0/3327	0.50	0/4515
2	B	0.29	0/2372	0.49	0/3233
2	E	0.25	0/2234	0.45	0/3049
3	C	0.30	0/2471	0.50	0/3357
3	F	0.31	0/2463	0.50	0/3346
All	All	0.29	0/16183	0.49	0/22000

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3237	0	3160	39	0
1	D	3245	0	3173	46	0
2	B	2331	0	2332	56	0
2	E	2194	0	2163	32	0
3	C	2406	0	2405	18	0
3	F	2398	0	2396	20	0
4	B	16	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	8	0	0	1	0
4	E	16	0	0	0	0
4	F	8	0	0	1	0
5	C	29	0	15	3	0
5	F	29	0	15	2	0
6	C	2	0	0	0	0
6	F	1	0	0	0	0
7	F	1	0	0	0	0
8	A	406	0	0	4	0
8	B	447	0	0	26	1
8	C	380	0	0	8	1
8	D	471	0	0	11	1
8	E	296	0	0	11	0
8	F	432	0	0	8	1
All	All	18353	0	15659	196	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 196 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:273:MET:SD	8:D:652:HOH:O	1.96	1.21
1:D:263:GLU:OE1	8:D:401:HOH:O	1.76	1.04
3:F:257:GLY:O	8:F:501:HOH:O	1.81	0.97
3:F:169:GLU:OE1	8:F:502:HOH:O	1.87	0.91
2:B:47:GLN:O	8:B:501:HOH:O	1.90	0.87

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:C:860:HOH:O	8:D:838:HOH:O[2_455]	1.95	0.25
8:B:529:HOH:O	8:F:773:HOH:O[3_555]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/395 (105%)	405 (98%)	10 (2%)	0	100	100
1	D	416/395 (105%)	402 (97%)	14 (3%)	0	100	100
2	B	305/315 (97%)	297 (97%)	8 (3%)	0	100	100
2	E	284/315 (90%)	274 (96%)	10 (4%)	0	100	100
3	C	312/314 (99%)	301 (96%)	10 (3%)	1 (0%)	46	33
3	F	311/314 (99%)	298 (96%)	12 (4%)	1 (0%)	46	33
All	All	2043/2048 (100%)	1977 (97%)	64 (3%)	2 (0%)	56	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	146	ILE
3	F	146	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/322 (105%)	331 (98%)	8 (2%)	57	46
1	D	340/322 (106%)	332 (98%)	8 (2%)	57	46
2	B	251/256 (98%)	248 (99%)	3 (1%)	78	74
2	E	236/256 (92%)	234 (99%)	2 (1%)	86	85
3	C	250/250 (100%)	245 (98%)	5 (2%)	63	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	F	249/250 (100%)	244 (98%)	5 (2%)	63	54
All	All	1665/1656 (100%)	1634 (98%)	31 (2%)	68	56

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	179	PRO
1	D	31[B]	ARG
3	F	172	LYS
3	C	264	LYS
1	D	33	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SF4	B	401	2	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	B	402	2	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	C	401	3	0,12,12	0.00	-	0,24,24	0.00	-
5	5SR	C	402	6	22,30,30	2.23	4 (18%)	30,45,45	1.78	9 (30%)
4	SF4	E	401	2	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	E	402	2	0,12,12	0.00	-	0,24,24	0.00	-
4	SF4	F	401	3	0,12,12	0.00	-	0,24,24	0.00	-
5	5SR	F	402	6	22,30,30	2.13	4 (18%)	30,45,45	1.87	9 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	B	401	2	-	0/0/48/48	0/6/5/5
4	SF4	B	402	2	-	0/0/48/48	0/6/5/5
4	SF4	C	401	3	-	0/0/48/48	0/6/5/5
5	5SR	C	402	6	-	0/16/21/21	0/2/2/2
4	SF4	E	401	2	-	0/0/48/48	0/6/5/5
4	SF4	E	402	2	-	0/0/48/48	0/6/5/5
4	SF4	F	401	3	-	0/0/48/48	0/6/5/5
5	5SR	F	402	6	-	0/16/21/21	0/2/2/2

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	402	5SR	C3-C8	-2.80	1.38	1.42
5	F	402	5SR	C3-C8	-2.74	1.38	1.42
5	F	402	5SR	C14-S13	5.14	1.83	1.74
5	C	402	5SR	C8-N9	5.42	1.48	1.34
5	F	402	5SR	C8-N9	5.43	1.48	1.34

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	402	5SR	N5-C6-N7	-2.84	120.16	125.50
5	F	402	5SR	N5-C6-N7	-2.77	120.31	125.50
5	C	402	5SR	C3-C4-N5	-2.45	119.57	123.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	402	5SR	C3-C4-N5	-2.39	119.69	123.86
5	F	402	5SR	O30-P24-O23	-2.24	99.26	106.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	401	SF4	1	0
5	C	402	5SR	3	0
4	F	401	SF4	1	0
5	F	402	5SR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	394/395 (99%)	-0.02	15 (3%)	44	46	15, 22, 38, 50	0
1	D	394/395 (99%)	-0.08	13 (3%)	50	52	14, 22, 39, 53	0
2	B	309/315 (98%)	0.92	66 (21%)	1	1	15, 38, 60, 93	0
2	E	290/315 (92%)	1.22	81 (27%)	1	0	19, 48, 74, 105	0
3	C	314/314 (100%)	-0.30	4 (1%)	79	81	14, 20, 40, 72	0
3	F	313/314 (99%)	-0.30	1 (0%)	94	94	15, 21, 37, 49	0
All	All	2014/2048 (98%)	0.20	180 (8%)	12	13	14, 24, 60, 105	0

The worst 5 of 180 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	219	VAL	9.3
2	B	213	ALA	8.9
2	B	218	ALA	7.9
2	E	186	LEU	7.5
2	E	213	ALA	7.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	MG	F	403	1/1	0.99	0.13	1.68	15,15,15,15	0
6	MG	C	403	1/1	0.99	0.13	1.22	13,13,13,13	0
6	MG	C	404	1/1	0.97	0.15	1.08	33,33,33,33	0
5	5SR	C	402	29/29	0.98	0.13	-0.47	12,18,24,28	3
5	5SR	F	402	29/29	0.98	0.13	-0.51	14,18,23,32	3
7	NA	F	404	1/1	0.96	0.08	-0.64	28,28,28,28	0
4	SF4	B	401	8/8	1.00	0.07	-0.93	17,18,19,20	0
4	SF4	E	402	8/8	0.99	0.05	-1.66	22,22,23,27	0
4	SF4	E	401	8/8	0.99	0.05	-1.83	19,22,23,23	0
4	SF4	C	401	8/8	1.00	0.07	-1.88	15,16,17,18	0
4	SF4	B	402	8/8	0.99	0.06	-2.19	17,18,19,22	0
4	SF4	F	401	8/8	0.99	0.06	-2.78	18,19,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.