



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2016 – 05:31 PM EDT

PDB ID : 5EYK  
Title : CRYSTAL STRUCTURE OF AURORA B IN COMPLEX WITH BI 847325  
Authors : Bader, G.; Zoepfel, A.  
Deposited on : 2015-11-25  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2910 (1.96-1.92)
Clashscore	102246	3095 (1.96-1.92)
Ramachandran outliers	100387	3062 (1.96-1.92)
Sidechain outliers	100360	3062 (1.96-1.92)
RSRZ outliers	91569	2915 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 3% (Poor fit), 87% (0-1 outliers), 8% (2 outliers), 3% (3+ outliers), 2% (Not modelled)
1	B	276	 3% (Poor fit), 91% (0-1 outliers), 9% (2 outliers), 7% (3+ outliers), 2% (Not modelled)
2	C	59	 2% (Poor fit), 47% (0-1 outliers), 8% (2 outliers), 41% (3+ outliers), 2% (Not modelled)
2	D	59	 12% (Poor fit), 31% (0-1 outliers), 7% (2 outliers), 63% (3+ outliers), 7% (Not modelled)

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aurora kinase B-A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	264	Total	C	N	O	P	S	40	0	0
			2203	1415	393	381	1	13			
1	B	276	Total	C	N	O	P	S	40	1	0
			2309	1479	415	400	1	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	VAL	GLY	engineered mutation	UNP Q6DE08
B	96	VAL	GLY	engineered mutation	UNP Q6DE08


- Molecule 2 is a protein called Inner centromere protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	35	Total	C	N	O	S	16	0	0
			288	184	47	56	1			
2	D	22	Total	C	N	O	S	21	0	0
			188	120	33	34	1			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	789	MET	-	initiating methionine	UNP O13024
D	789	MET	-	initiating methionine	UNP O13024

- Molecule 3 is 3-[(3 {Z})-3-[[[4-[(dimethylamino)methyl]phenyl]amino]-phenyl-methylidene]-2-oxidanylidene-1 {H}-indol-6-yl]- {N}-ethyl-prop-2-ynamide (three-letter code: 5U5) (formula: C<sub>29</sub>H<sub>28</sub>N<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			35	29	4	2		
3	B	1	Total	C	N	O	0	0
			35	29	4	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	178	Total	O	0	0
			178	178		
4	B	184	Total	O	0	0
			184	184		
4	C	14	Total	O	0	0
			14	14		
4	D	5	Total	O	0	0
			5	5		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aurora kinase B-A




- Molecule 1: Aurora kinase B-A



- Molecule 2: Inner centromere protein A



- Molecule 2: Inner centromere protein A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.14Å 67.45Å 116.76Å 90.00° 96.63° 90.00°	Depositor
Resolution (Å)	19.46 – 1.93 37.91 – 1.93	Depositor EDS
% Data completeness (in resolution range)	96.5 (19.46-1.93) 96.5 (37.91-1.93)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 1.94Å)	Xtrriage
Refinement program	BUSTER 2.9.2	Depositor
R, $R_{free}$	0.199 , 0.222 0.199 , 0.226	Depositor DCC
$R_{free}$ test set	2590 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.7	Xtrriage
Anisotropy	0.618	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 52.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5439	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, 5U5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	1/2251 (0.0%)	0.64	0/3029
1	B	0.53	0/2360	0.64	0/3174
2	C	0.57	0/293	0.69	0/395
2	D	0.48	0/191	0.56	0/257
All	All	0.54	1/5095 (0.0%)	0.64	0/6855

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	259	GLU	CD-OE1	-5.11	1.20	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2203	0	2202	10	0
1	B	2309	0	2317	19	0
2	C	288	0	285	12	0
2	D	188	0	187	2	0
3	A	35	0	0	0	0
3	B	35	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	178	0	0	0	0
4	B	184	0	0	0	0
4	C	14	0	0	0	0
4	D	5	0	0	0	0
All	All	5439	0	4991	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:ILE:H	2:C:814:GLN:HE22	1.21	0.86
1:A:118:ILE:H	2:D:814:GLN:HE22	1.25	0.85
2:C:803:SER:C	2:C:807:LEU:HD21	1.96	0.83
1:B:97:ARG:HH12	2:C:807:LEU:HD11	1.44	0.82
1:A:150:HIS:HD2	1:A:152:ASN:H	1.29	0.81
1:B:150:HIS:HD2	1:B:152:ASN:H	1.30	0.80
1:B:97:ARG:NH1	2:C:807:LEU:HD11	1.98	0.78
1:A:150:HIS:CD2	1:A:152:ASN:H	2.12	0.68
1:B:150:HIS:CD2	1:B:152:ASN:H	2.12	0.65
1:B:146:SER:HB3	1:B:156:MET:CE	2.28	0.62
1:B:97:ARG:NH1	2:C:803:SER:O	2.33	0.61
1:B:277:VAL:HG13	1:B:288:PRO:HD2	1.82	0.61
2:C:803:SER:CA	2:C:807:LEU:HD21	2.35	0.56
1:A:277:VAL:HG13	1:A:288:PRO:HD2	1.87	0.56
1:B:118:ILE:H	2:C:814:GLN:NE2	1.98	0.55
1:B:146:SER:HB3	1:B:156:MET:HE1	1.91	0.52
1:B:89:THR:HG22	1:B:91:ASP:H	1.76	0.51
2:C:804:GLY:O	2:C:807:LEU:HG	2.10	0.51
1:A:152:ASN:HD21	1:A:349:ARG:HH21	1.58	0.51
1:B:261:ILE:CG2	1:B:298:HIS:CD2	2.94	0.51
1:A:237:TRP:CD2	1:A:250:CYS:HB2	2.46	0.50
1:B:257:PRO:HD2	1:B:260:MET:HG3	1.92	0.50
1:B:152:ASN:HD21	1:B:349:ARG:HH21	1.58	0.50
1:B:261:ILE:HG23	1:B:298:HIS:CD2	2.48	0.49
1:A:145:GLN:O	1:A:145:GLN:HG3	2.12	0.48
1:B:239:VAL:HG11	1:B:246:ARG:HG3	1.96	0.46
1:A:158:ASN:ND2	2:D:825:TYR:O	2.50	0.43
2:C:803:SER:CA	2:C:807:LEU:CD2	2.97	0.43
1:B:123:VAL:HG11	1:B:165:ARG:HH11	1.84	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:ARG:HD3	1:A:248:TPO:O2P	2.20	0.42
1:A:90:ILE:H	1:A:90:ILE:HG13	1.70	0.41
2:C:803:SER:HA	2:C:807:LEU:CD2	2.51	0.41
2:C:803:SER:O	2:C:807:LEU:HD21	2.20	0.41
1:B:256:LEU:HA	1:B:257:PRO:HD3	1.99	0.40
1:B:353:PRO:HD3	2:C:815:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/276 (94%)	251 (97%)	8 (3%)	0	100	100
1	B	274/276 (99%)	268 (98%)	6 (2%)	0	100	100
2	C	33/59 (56%)	32 (97%)	1 (3%)	0	100	100
2	D	20/59 (34%)	20 (100%)	0	0	100	100
All	All	586/670 (88%)	571 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/249 (96%)	226 (95%)	12 (5%)	30	14
1	B	250/249 (100%)	245 (98%)	5 (2%)	63	53
2	C	32/54 (59%)	28 (88%)	4 (12%)	6	1
2	D	20/54 (37%)	18 (90%)	2 (10%)	9	2
All	All	540/606 (89%)	517 (96%)	23 (4%)	35	19

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	ILE
1	A	101	LYS
1	A	103	LYS
1	A	112	GLU
1	A	137	GLN
1	A	148	LEU
1	A	179	TYR
1	A	192	ARG
1	A	246	ARG
1	A	249	MET
1	A	294	HIS
1	A	298	HIS
1	B	101	LYS
1	B	139	ARG
1	B	161	HIS
1	B	163	ARG
1	B	294	HIS
2	C	803	SER
2	C	807	LEU
2	C	822	ASP
2	C	823	ARG
2	D	805	ASN
2	D	822	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	106	ASN
1	A	147	HIS
1	A	150	HIS
1	A	152	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	161	HIS
1	A	185	HIS
1	B	106	ASN
1	B	147	HIS
1	B	150	HIS
1	B	152	ASN
1	B	183	GLN
1	B	185	HIS
2	C	814	GLN
2	D	814	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	TPO	A	248	1	7,10,11	0.83	0	10,14,16	1.55	3 (30%)
1	TPO	B	248	1	7,10,11	1.25	1 (14%)	10,14,16	1.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	248	1	-	0/8/11/13	0/0/0/0
1	TPO	B	248	1	-	0/8/11/13	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	248	TPO	CG2-CB	2.23	1.56	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	248	TPO	O-C-CA	-2.13	119.86	125.69
1	A	248	TPO	P-OG1-CB	-2.04	112.52	121.42
1	A	248	TPO	O2P-P-OG1	2.64	114.51	106.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	248	TPO	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	5U5	A	401	-	38,38,38	1.36	4 (10%)	49,52,52	2.18	14 (28%)
3	5U5	B	401	-	38,38,38	1.61	6 (15%)	49,52,52	1.71	12 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5U5	A	401	-	-	0/21/36/36	0/4/4/4
3	5U5	B	401	-	-	0/21/36/36	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	5U5	C26-C27	-4.76	1.42	1.45
3	B	401	5U5	C8-N7	-3.06	1.32	1.36
3	A	401	5U5	C9-C8	-3.02	1.43	1.50
3	A	401	5U5	C8-N7	-2.79	1.33	1.36
3	B	401	5U5	C26-C25	-2.63	1.15	1.20
3	B	401	5U5	C9-C8	-2.23	1.44	1.50
3	B	401	5U5	C1-C2	2.01	1.43	1.39
3	A	401	5U5	C14-C13	3.28	1.44	1.39
3	B	401	5U5	C18-C13	3.66	1.45	1.39
3	A	401	5U5	C18-C13	3.76	1.45	1.39

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	5U5	C9-C11-N12	-5.74	113.82	118.12
3	A	401	5U5	C18-C13-C11	-4.62	115.36	120.76
3	B	401	5U5	C3-C6-C5	-4.19	118.33	122.42
3	B	401	5U5	C9-C11-N12	-4.19	114.98	118.12
3	A	401	5U5	C6-N7-C8	-3.88	109.10	111.34
3	B	401	5U5	C4-C1-C2	-2.95	116.66	120.36
3	A	401	5U5	C3-C6-C5	-2.89	119.60	122.42
3	B	401	5U5	O10-C8-C9	-2.73	124.16	128.16
3	A	401	5U5	C35-N29-C27	-2.70	118.60	122.53
3	B	401	5U5	C18-C13-C11	-2.62	117.69	120.76
3	B	401	5U5	C35-N29-C27	-2.54	118.82	122.53
3	B	401	5U5	C1-C2-C25	-2.46	116.34	120.79
3	A	401	5U5	C15-C14-C13	-2.36	117.53	120.35
3	B	401	5U5	O28-C27-C26	-2.29	117.79	121.21
3	A	401	5U5	O10-C8-C9	-2.21	124.93	128.16
3	A	401	5U5	O28-C27-N29	-2.14	118.04	122.52
3	A	401	5U5	C20-C19-N12	-2.04	113.59	120.40
3	A	401	5U5	C21-C22-C23	2.00	121.46	118.15
3	A	401	5U5	C11-C9-C8	2.13	126.75	122.70
3	B	401	5U5	C33-N31-C32	2.16	115.41	109.71
3	B	401	5U5	C19-N12-C11	2.28	135.38	127.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	5U5	C13-C11-N12	2.71	123.09	119.33
3	B	401	5U5	C9-C8-N7	3.30	109.02	106.82
3	A	401	5U5	C13-C11-N12	4.14	125.08	119.33
3	A	401	5U5	C14-C13-C11	4.50	126.01	120.76
3	A	401	5U5	C9-C8-N7	6.51	111.16	106.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/276 (95%)	0.20	9 (3%) 49 58	18, 33, 59, 72	10 (3%)
1	B	275/276 (99%)	-0.17	7 (2%) 61 69	15, 30, 56, 71	10 (3%)
2	C	35/59 (59%)	0.65	1 (2%) 55 63	34, 47, 66, 70	4 (11%)
2	D	22/59 (37%)	1.33	7 (31%) 1 0	43, 63, 76, 85	5 (22%)
All	All	595/670 (88%)	0.10	24 (4%) 42 51	15, 34, 63, 85	29 (4%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	807	LEU	5.0
1	B	81	ALA	5.0
1	A	355	TYR	4.6
2	D	819	ILE	4.4
2	D	822	ASP	4.2
1	A	159	TYR	3.7
1	B	356	GLN	3.4
2	D	824	MET	3.4
2	C	837	PHE	3.2
1	A	249	MET	2.9
1	A	250	CYS	2.9
1	B	104	PHE	2.8
1	A	96	VAL	2.8
1	A	240	HIS	2.8
2	D	806	LEU	2.8
1	A	354	VAL	2.7
1	A	88	PHE	2.4
1	B	129	LEU	2.2
1	B	138	LEU	2.2
1	A	237	TRP	2.1
1	B	179	TYR	2.1

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	355	TYR	2.1
2	D	825	TYR	2.0
2	D	823	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	B	248	11/12	0.96	0.08	-	32,33,36,37	0
1	TPO	A	248	11/12	0.88	0.24	-	60,64,67,67	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	5U5	B	401	35/35	0.94	0.11	0.46	24,33,48,49	0
3	5U5	A	401	35/35	0.96	0.09	-0.47	22,29,45,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.