



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:27 PM GMT

PDB ID : 4F0I
Title : Crystal structure of apo TrkA
Authors : Liu, J.
Deposited on : 2012-05-04
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	<div> <div>3%</div> <div>77%</div> <div>15%</div> <div>5%</div> </div>
1	B	300	<div> <div>%</div> <div>78%</div> <div>17%</div> <div>.</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called High affinity nerve growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	0	0	0
			2269	1452	411	390	16			
1	B	294	Total	C	N	O	S	0	0	0
			2351	1502	422	411	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	496	GLY	-	EXPRESSION TAG	UNP P04629
B	496	GLY	-	EXPRESSION TAG	UNP P04629


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	113	Total	O	0	0
			113	113		
2	B	199	Total	O	0	0
			199	199		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: High affinity nerve growth factor receptor



- Molecule 1: High affinity nerve growth factor receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.59 Å 92.59 Å 181.50 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.43 – 2.30 44.43 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.9 (44.43-2.30) 99.4 (44.43-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.93 (at 2.29 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.182 , 0.241 0.186 , 0.242	Depositor DCC
R_{free} test set	1789 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 35700 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4932	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/2325	0.51	0/3144
1	B	0.44	0/2410	0.57	0/3263
All	All	0.41	0/4735	0.54	0/6407

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2269	0	2271	38	0
1	B	2351	0	2342	34	0
2	A	113	0	0	1	0
2	B	199	0	0	3	0
All	All	4932	0	4613	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:553:ARG:NH2	1:A:579:THR:HB	1.97	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:729:GLN:H	1:B:729:GLN:HE21	1.29	0.78
1:A:553:ARG:HG2	1:A:553:ARG:HH11	1.51	0.76
1:B:603:HIS:HD2	1:B:619:GLY:O	1.70	0.74
1:B:736:ILE:O	1:B:740:THR:HB	1.88	0.74
1:A:557:GLN:O	1:A:561:GLU:HG3	1.89	0.73
1:A:553:ARG:CG	1:A:553:ARG:HH11	2.02	0.71
1:A:572:VAL:HG21	1:A:656:LEU:HD12	1.73	0.71
1:B:559:GLU:HG2	1:B:563:LEU:HD22	1.74	0.69
1:A:529:HIS:NE2	1:A:537:LYS:HE3	2.09	0.68
1:A:707:SER:HB3	2:A:849:HOH:O	1.93	0.68
1:A:704:THR:H	1:A:707:SER:HB2	1.59	0.67
1:A:774:LYS:HE2	1:A:774:LYS:HA	1.77	0.66
1:A:647:HIS:O	1:A:648:ARG:HB2	1.96	0.64
1:B:572:VAL:HG21	1:B:656:LEU:HD12	1.79	0.64
1:B:553:ARG:HH21	1:B:582:ARG:HH11	1.46	0.64
1:B:603:HIS:CD2	1:B:619:GLY:O	2.51	0.64
1:B:577:VAL:HG13	1:B:579:THR:HG23	1.82	0.62
1:A:603:HIS:HD2	1:A:619:GLY:O	1.82	0.62
1:B:507:ARG:HD2	2:B:919:HOH:O	2.03	0.58
1:A:553:ARG:HH22	1:A:579:THR:HB	1.69	0.57
1:A:569:GLN:HG3	1:A:570:HIS:CD2	2.40	0.57
1:B:581:GLY:C	1:B:582:ARG:HE	2.11	0.54
1:A:559:GLU:O	1:A:563:LEU:HD22	2.06	0.54
1:A:527:GLU:HA	1:A:538:MET:O	2.08	0.54
1:A:687:MET:HE3	1:A:687:MET:HA	1.90	0.54
1:B:604:GLY:C	1:B:606:ASP:H	2.12	0.53
1:B:510:VAL:HG13	1:B:527:GLU:HB2	1.90	0.53
1:B:567:GLN:HG2	2:B:859:HOH:O	2.09	0.53
1:B:715:VAL:O	1:B:719:ILE:HG13	2.09	0.53
1:A:529:HIS:CE1	1:A:537:LYS:HE3	2.44	0.52
1:B:556:PHE:CZ	1:B:577:VAL:HG22	2.44	0.52
1:A:527:GLU:HB3	1:A:537:LYS:HE2	1.92	0.52
1:A:580:GLU:HG2	1:A:580:GLU:O	2.10	0.51
1:A:553:ARG:CG	1:A:553:ARG:NH1	2.68	0.50
1:B:694:PRO:HB3	1:B:710:TRP:CD2	2.46	0.50
1:A:603:HIS:CD2	1:A:619:GLY:O	2.63	0.50
1:B:558:ARG:NE	2:B:900:HOH:O	2.43	0.50
1:A:531:LEU:HD21	1:A:575:PHE:CG	2.48	0.49
1:A:527:GLU:CB	1:A:537:LYS:HE2	2.43	0.49
1:B:740:THR:O	1:B:765:ARG:NH2	2.45	0.48
1:A:747:ARG:HG3	1:A:756:TYR:HB2	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:505:LYS:HD3	1:A:507:ARG:NH1	2.29	0.48
1:A:511:LEU:HA	1:A:526:ALA:HB2	1.97	0.47
1:B:729:GLN:NE2	1:B:729:GLN:H	2.04	0.47
1:A:536:ASP:O	1:A:538:MET:HB2	2.14	0.47
1:B:682:VAL:HG21	1:B:688:LEU:HD12	1.96	0.47
1:A:510:VAL:HG13	1:A:527:GLU:HB2	1.97	0.47
1:B:681:ARG:HD2	1:B:683:GLY:O	2.16	0.46
1:B:690:ILE:HD13	1:B:736:ILE:HG13	1.95	0.46
1:B:699:LEU:HB3	1:B:700:TYR:CD2	2.50	0.46
1:B:756:TYR:O	1:B:760:ARG:HG3	2.16	0.46
1:B:553:ARG:HE	1:B:582:ARG:NH1	2.12	0.46
1:A:610:LEU:HG	1:A:618:PRO:HB3	1.98	0.45
1:A:536:ASP:O	1:A:537:LYS:C	2.55	0.45
1:A:527:GLU:OE2	1:A:537:LYS:HE2	2.17	0.44
1:A:568:HIS:HB3	1:A:571:ILE:HG12	1.98	0.44
1:B:537:LYS:HE3	1:B:537:LYS:O	2.17	0.44
1:A:680:TYR:HA	1:A:687:MET:HE1	2.00	0.43
1:B:610:LEU:C	1:B:610:LEU:HD12	2.39	0.43
1:A:680:TYR:CA	1:A:687:MET:HE1	2.49	0.43
1:B:557:GLN:O	1:B:561:GLU:HG2	2.19	0.43
1:B:608:LYS:H	1:B:608:LYS:HG2	1.63	0.43
1:B:699:LEU:HB3	1:B:700:TYR:CE2	2.54	0.42
1:A:754:GLU:CD	1:A:754:GLU:H	2.23	0.42
1:A:787:PRO:HA	1:A:788:PRO:HD2	1.85	0.41
1:A:569:GLN:O	1:A:664:LYS:HE2	2.20	0.41
1:B:646:VAL:HG12	1:B:648:ARG:HG3	2.02	0.41
1:B:580:GLU:HG3	1:B:580:GLU:O	2.20	0.41
1:A:679:TYR:HB3	1:A:687:MET:HE2	2.02	0.41
1:B:647:HIS:O	1:B:648:ARG:HB2	2.20	0.41
1:B:553:ARG:NH2	1:B:582:ARG:HH11	2.16	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/300 (93%)	270 (97%)	8 (3%)	0	100	100
1	B	292/300 (97%)	284 (97%)	7 (2%)	1 (0%)	46	57
All	All	570/600 (95%)	554 (97%)	15 (3%)	1 (0%)	52	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	605	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/250 (95%)	221 (93%)	17 (7%)	18	23
1	B	247/250 (99%)	223 (90%)	24 (10%)	10	12
All	All	485/500 (97%)	444 (92%)	41 (8%)	13	16

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	511	LEU
1	A	515	LEU
1	A	553	ARG
1	A	562	LEU
1	A	563	LEU
1	A	580	GLU
1	A	598	ARG
1	A	623	LEU
1	A	687	MET
1	A	699	LEU
1	A	707	SER
1	A	709	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	716	LEU
1	A	729	GLN
1	A	754	GLU
1	A	769	GLN
1	A	774	LYS
1	B	500	CYS
1	B	510	VAL
1	B	515	LEU
1	B	536	ASP
1	B	562	LEU
1	B	563	LEU
1	B	577	VAL
1	B	582	ARG
1	B	608	LYS
1	B	609	LEU
1	B	610	LEU
1	B	614	GLU
1	B	623	LEU
1	B	661	LEU
1	B	685	ARG
1	B	699	LEU
1	B	709	VAL
1	B	729	GLN
1	B	738	CYS
1	B	740	THR
1	B	765	ARG
1	B	769	GLN
1	B	791	LEU
1	B	793	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	557	GLN
1	A	567	GLN
1	A	569	GLN
1	A	570	HIS
1	A	603	HIS
1	A	768	GLN
1	A	771	HIS
1	A	785	GLN
1	B	529	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	557	GLN
1	B	603	HIS
1	B	729	GLN
1	B	768	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	284/300 (94%)	-0.15	8 (2%) 56 66	22, 44, 86, 133	0
1	B	294/300 (98%)	-0.35	4 (1%) 78 83	17, 30, 71, 108	0
All	All	578/600 (96%)	-0.25	12 (2%) 67 74	17, 37, 84, 133	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	609	LEU	5.8
1	A	501	VAL	3.0
1	A	500	CYS	3.0
1	A	551	SER	2.9
1	B	606	ASP	2.7
1	A	783	LEU	2.6
1	A	536	ASP	2.4
1	A	537	LYS	2.3
1	B	608	LYS	2.2
1	A	532	LEU	2.2
1	A	533	PRO	2.1
1	B	790	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.