wwPDB X-ray Structure Validation Summary Report (i)

Oct 4, 2016 – 05:31 AM EDT

PDB ID	:	5F42
Title	:	Activity and Crystal Structure of Francisella novicida UDP-N-Acetylglucosamine Acyltransferase
Authors	:	Joo, S.H.; Raetz, C.R.H.
Deposited on	:	2015-12-03
Resolution	:	2.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1799 (2.08-2.04)
Clashscore	102246	1910 (2.08-2.04)
Ramachandran outliers	100387	1893 (2.08-2.04)
Sidechain outliers	100360	1893 (2.08-2.04)
RSRZ outliers	91569	1802 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	6% 88% 11% •
1	B	280	6% 89% 9% •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4579 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	2100	1328	369	390	13	0	0	0
1	B	276	2100	1328	369	390	13	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	260	ASP	-	expression tag	UNP A0Q7Y0
A	261	PRO	-	expression tag	UNP A0Q7Y0
A	262	ASN	-	expression tag	UNP A0Q7Y0
A	263	SER	-	expression tag	UNP A0Q7Y0
A	264	SER	-	expression tag	UNP A0Q7Y0
A	265	SER	-	expression tag	UNP A0Q7Y0
A	266	VAL	-	expression tag	UNP A0Q7Y0
A	267	ASP	-	expression tag	UNP A0Q7Y0
A	268	LYS	-	expression tag	UNP A0Q7Y0
A	269	LEU	-	expression tag	UNP A0Q7Y0
A	270	ALA	-	expression tag	UNP A0Q7Y0
A	271	ALA	-	expression tag	UNP A0Q7Y0
A	272	ALA	-	expression tag	UNP A0Q7Y0
A	273	LEU	-	expression tag	UNP A0Q7Y0
A	274	GLU	-	expression tag	UNP A0Q7Y0
A	275	HIS	-	expression tag	UNP A0Q7Y0
A	276	HIS	-	expression tag	UNP A0Q7Y0
A	277	HIS	-	expression tag	UNP A0Q7Y0
A	278	HIS	-	expression tag	UNP A0Q7Y0
A	279	HIS	-	expression tag	UNP A0Q7Y0
A	280	HIS	-	expression tag	UNP A0Q7Y0
B	260	ASP	-	expression tag	UNP A0Q7Y0
B	261	PRO	-	expression tag	UNP A0Q7Y0
B	262	ASN	-	expression tag	UNP A0Q7Y0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	263	SER	-	expression tag	UNP A0Q7Y0
B	264	SER	-	expression tag	UNP A0Q7Y0
B	265	SER	-	expression tag	UNP A0Q7Y0
B	266	VAL	-	expression tag	UNP A0Q7Y0
B	267	ASP	-	expression tag	UNP A0Q7Y0
B	268	LYS	-	expression tag	UNP A0Q7Y0
B	269	LEU	-	expression tag	UNP A0Q7Y0
B	270	ALA	-	expression tag	UNP A0Q7Y0
B	271	ALA	-	expression tag	UNP A0Q7Y0
B	272	ALA	-	expression tag	UNP A0Q7Y0
B	273	LEU	-	expression tag	UNP A0Q7Y0
B	274	GLU	-	expression tag	UNP A0Q7Y0
B	275	HIS	-	expression tag	UNP A0Q7Y0
B	276	HIS	-	expression tag	UNP A0Q7Y0
B	277	HIS	-	expression tag	UNP A0Q7Y0
B	278	HIS	-	expression tag	UNP A0Q7Y0
B	279	HIS	-	expression tag	UNP A0Q7Y0
B	280	HIS	-	expression tag	UNP A0Q7Y0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	208	208	208	0	0
			Total	O		
2	B	171	171	171	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase

Chain A:





- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α, β, γ	105.38Å 105.38Å 285.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	18.95 – 2.06 18.95 – 2.06	Depositor EDS
% Data completeness (in resolution range)	100.0 (18.95-2.06) 99.4 (18.95-2.06)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.41 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.210 , 0.260 0.209 , 0.257	Depositor DCC
R_{free} test set	1894 reflections (5.26%)	DCC
Wilson B-factor (Å^2)	25.5	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 47.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	4579	wwPDB-VP
Average B, all atoms (Å^2)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.02% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.40	0/2137	0.55	0/2886
1	B	0.37	0/2137	0.54	0/2886
All	All	0.39	0/4274	0.54	0/5772

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2100	0	2129	17	0
1	B	2100	0	2129	16	0
2	A	208	0	0	0	0
2	B	171	0	0	1	0
All	All	4579	0	4258	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 33 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:PRO:HB3	1:B:120:HIS:HB3	1.76	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:SER:OG	1:B:34:GLU:HG3	1.96	0.65
1:B:250:ILE:HA	1:B:257:ILE:HD11	1.80	0.64
1:A:143:ASP:HB3	1:A:162:LYS:HG2	1.85	0.58
1:A:19:ILE:HB	1:A:37:GLU:HG3	1.85	0.57

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/280 (98%)	264 (96%)	9 (3%)	1 (0%)	39	28
1	B	274/280 (98%)	262 (96%)	11 (4%)	1 (0%)	39	28
All	All	548/560 (98%)	526 (96%)	20 (4%)	2 (0%)	39	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	185	VAL
1	B	185	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	228/232 (98%)	227 (100%)	1 (0%)	93 94
1	B	228/232 (98%)	228 (100%)	0	100 100
All	All	456/464 (98%)	455 (100%)	1 (0%)	95 96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	276/280 (98%)	0.20	18 (6%)	22	24	15, 25, 43, 51	0
1	B	276/280 (98%)	0.31	16 (5%)	26	29	16, 31, 46, 54	0
All	All	552/560 (98%)	0.25	34 (6%)	24	26	15, 28, 45, 54	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	186	ASN	11.2
1	A	186	ASN	8.2
1	A	74	GLY	7.2
1	B	185	VAL	6.5
1	A	187	ALA	6.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.