



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:21 PM GMT

PDB ID : 1F7U
Title : CRYSTAL STRUCTURE OF THE ARGINYL-TRNA SYNTHETASE COM-
PLEXED WITH THE TRNA(ARG) AND L-ARG
Authors : Delagoutte, B.; Moras, D.; Cavarelli, J.
Deposited on : 2000-06-28
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)
RNA backbone	2183	1062 (2.80-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	76	<div> <div>4%</div> <div>71%</div> <div>26%</div> <div>.</div> </div>
2	A	607	<div> <div>16%</div> <div>74%</div> <div>24%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called TRNA(ARG).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	76	Total	C	N	O	P	0	0	0
			1629	730	290	533	76			

- Molecule 2 is a protein called ARGINYL-TRNA SYNTHETASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	606	Total	C	N	O	S	0	0	0
			4892	3138	828	905	21			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ARGinine (three-letter code: ARG) (formula: C₆H₁₅N₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	6	4	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	348	Total	O	0	0
			348	348		
5	B	240	Total	O	0	0
			240	240		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRNA(ARG)



• Molecule 2: ARGINYL-TRNA SYNTHETASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	129.65Å 107.47Å 71.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.98 – 2.20 24.82 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (14.98-2.20) 99.8 (24.82-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.71 (at 2.19Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.190 , 0.233 0.190 , 0.233	Depositor DCC
R_{free} test set	2558 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	20.7	Xtriage
Anisotropy	0.725	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 66.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 51242 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7126	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5MU, H2U, SO4, 2MG, 5MC, 1MA, M2G, 1MG, PSU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.30	0/1532	0.69	0/2387
2	A	0.30	0/4996	0.55	0/6736
All	All	0.30	0/6528	0.59	0/9123

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1629	0	834	18	0
2	A	4892	0	4906	119	0
3	B	5	0	0	0	0
4	A	12	0	12	1	0
5	A	348	0	0	3	0
5	B	240	0	0	2	0
All	All	7126	0	5752	133	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 133 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:299:LYS:H	2:A:299:LYS:HE3	1.11	1.15
2:A:331:LYS:H	2:A:331:LYS:HD2	1.24	1.00
2:A:323:LEU:HB3	2:A:337:ILE:HD13	1.53	0.90
2:A:299:LYS:N	2:A:299:LYS:HE3	1.94	0.83
2:A:331:LYS:HD2	2:A:331:LYS:N	1.97	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	604/607 (100%)	576 (95%)	23 (4%)	5 (1%)	24 22

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	237	SER
2	A	259	ASP
2	A	240	LEU
2	A	150	SER
2	A	216	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	525/526 (100%)	507 (97%)	18 (3%)	44 54

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	299	LYS
2	A	349	THR
2	A	526	LEU
2	A	253	LYS
2	A	260	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	A	245	ASN
2	A	508	GLN
2	A	406	GLN
2	A	62	ASN
2	A	442	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	B	75/76 (98%)	2 (2%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	B	909	1MG
1	B	974	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PSU	B	901	1	18,22,22	1.83	4 (22%)	24,33,33	5.15	4 (16%)
1	1MG	B	909	1	16,26,27	0.87	0	19,39,42	1.57	2 (10%)
1	2MG	B	910	1	17,26,27	1.20	2 (11%)	21,38,41	3.93	5 (23%)
1	H2U	B	916	1	17,21,22	0.72	1 (5%)	23,30,33	1.14	2 (8%)
1	H2U	B	919	1	17,21,22	0.66	0	23,30,33	1.28	1 (4%)
1	M2G	B	926	1	17,27,28	1.25	2 (11%)	22,40,43	3.71	6 (27%)
1	PSU	B	927	1	13,21,22	2.21	3 (23%)	18,30,33	5.93	3 (16%)
1	H2U	B	947	1	17,21,22	0.81	1 (5%)	23,30,33	1.36	1 (4%)
1	5MC	B	949	1	13,22,23	1.05	1 (7%)	15,32,35	0.57	0
1	5MU	B	954	1	12,22,23	1.27	3 (25%)	14,32,35	4.42	3 (21%)
1	PSU	B	955	1	13,21,22	2.16	4 (30%)	18,30,33	5.94	3 (16%)
1	1MA	B	958	1	14,25,26	0.98	1 (7%)	15,37,40	1.18	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PSU	B	901	1	-	0/10/26/26	0/2/2/2
1	1MG	B	909	1	-	0/3/25/26	0/3/3/3
1	2MG	B	910	1	-	0/5/27/28	0/3/3/3
1	H2U	B	916	1	-	0/7/38/39	0/2/2/2
1	H2U	B	919	1	-	0/7/38/39	0/2/2/2
1	M2G	B	926	1	-	0/7/29/30	0/3/3/3
1	PSU	B	927	1	-	0/7/25/26	0/2/2/2
1	H2U	B	947	1	-	0/7/38/39	0/2/2/2
1	5MC	B	949	1	-	0/3/25/26	0/2/2/2
1	5MU	B	954	1	-	0/3/25/26	0/2/2/2
1	PSU	B	955	1	-	0/7/25/26	0/2/2/2
1	1MA	B	958	1	-	0/3/25/26	0/3/3/3

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	927	PSU	C5-C1'	-5.16	1.47	1.52
1	B	927	PSU	C6-N1	-4.75	1.24	1.34
1	B	955	PSU	C5-C1'	-4.73	1.48	1.52
1	B	955	PSU	C6-N1	-4.71	1.24	1.34
1	B	901	PSU	C6-N1	-4.64	1.24	1.34

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	955	PSU	N1-C2-N3	-21.63	114.53	128.33
1	B	927	PSU	N1-C2-N3	-21.55	114.58	128.33
1	B	901	PSU	N1-C2-N3	-21.47	114.63	128.33
1	B	910	2MG	CM2-N2-C2	-13.19	108.19	123.07
1	B	926	M2G	CM1-N2-C2	-9.51	111.56	121.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	901	PSU	1	0
1	B	909	1MG	1	0
1	B	919	H2U	1	0
1	B	926	M2G	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ARG	A	800	-	5,11,11	0.32	0	3,13,13	0.17	0
3	SO4	B	900	-	4,4,4	0.14	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ARG	A	800	-	-	0/5/11/11	0/0/0/0
3	SO4	B	900	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	800	ARG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	63/76 (82%)	-0.14	3 (4%) 34 34	7, 17, 43, 75	0
2	A	606/607 (99%)	0.84	95 (15%) 3 2	5, 26, 83, 120	0
All	All	669/683 (97%)	0.75	98 (14%) 3 3	5, 25, 82, 120	0

The worst 5 of 98 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	410	THR	14.5
2	A	238	ILE	13.8
2	A	237	SER	12.7
2	A	236	ASP	10.9
2	A	235	GLY	10.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	PSU	B	927	20/21	0.95	0.12	-	14,21,25,30	0
1	1MA	B	958	23/24	0.96	0.11	-	8,11,15,15	0
1	M2G	B	926	25/26	0.96	0.11	-	10,15,20,22	0
1	PSU	B	955	20/21	0.93	0.14	-	15,19,22,24	0
1	H2U	B	916	20/21	0.96	0.09	-	11,14,18,19	0
1	2MG	B	910	24/25	0.96	0.10	-	4,11,17,18	0
1	H2U	B	947	20/21	0.83	0.24	-	34,44,57,57	0
1	5MU	B	954	21/22	0.96	0.11	-	8,14,17,21	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	PSU	B	901	21/21	0.78	0.32	-	56,58,68,69	0
1	1MG	B	909	24/25	0.96	0.11	-	8,12,17,21	0
1	H2U	B	919	20/21	0.91	0.13	-	22,25,30,38	0
1	5MC	B	949	21/22	0.98	0.10	-	3,12,14,14	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ARG	A	800	12/12	0.90	0.16	0.88	23,25,30,32	0
3	SO4	B	900	5/5	0.99	0.14	-	30,33,35,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.