



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 14, 2016 – 07:40 PM EDT

PDB ID : 5FI4  
Title : Discovery of imidazo[1,2-a]-pyridine inhibitors of pan-PI3 kinases that are efficacious in a mouse xenograft model  
Authors : Elling, R.A.; Knapp, M.S.; Han, W.; Daniel, L.M.; Xy, Y.; Burger, M.T.; Ni, Z.; Smith, A.; Lan, J.; Williams, T.; Verhagen, J.; Huh, K.; Merritt, H.; Chan, J.; Kaufman, S.; Voliva, C.F.; Pecchi, S.  
Deposited on : 2015-12-22  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1074	<div> <div>3%</div> <div>84%</div> <div>10%</div> <div>6%</div> </div>
2	B	323	<div> <div>7%</div> <div>70%</div> <div>7%</div> <div>23%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10499 atoms, of which 23 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1009	Total	C	N	O	S	0	5	0
			8126	5203	1381	1477	65			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	LYS	MET	engineered mutation	UNP P42336
A	233	LYS	LEU	engineered mutation	UNP P42336
A	1069	HIS	-	expression tag	UNP P42336
A	1070	HIS	-	expression tag	UNP P42336
A	1071	HIS	-	expression tag	UNP P42336
A	1072	HIS	-	expression tag	UNP P42336
A	1073	HIS	-	expression tag	UNP P42336
A	1074	HIS	-	expression tag	UNP P42336

- Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	249	Total	C	N	O	S	0	0	0
			2054	1288	361	400	5			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	295	MET	-	initiating methionine	UNP P27986
B	296	GLU	-	expression tag	UNP P27986
B	297	TYR	-	expression tag	UNP P27986
B	298	MET	-	expression tag	UNP P27986
B	299	PRO	-	expression tag	UNP P27986
B	300	MET	-	expression tag	UNP P27986
B	301	GLU	-	expression tag	UNP P27986

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	306	TYR	THR	engineered mutation	UNP P27986

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	H	O	
			14	3	8	3	

- Molecule 4 is {N}-[6-[5-azanyl-6-[(2 {S})-1,1,1-tris(fluoranyl)propan-2-yl]oxy-pyrazin-2-yl]imidazo[1,2-a]pyridin-2-yl]ethanamide (three-letter code: 5XV) (formula:  $C_{16}H_{15}F_3N_6O_2$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	F	H	N	O	0	0
			42	16	3	15	6	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	237	Total	O	0	0
			237	237		
5	B	26	Total	O	0	0
			26	26		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.20Å 105.32Å 134.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.56 – 2.50 38.38 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (38.56-2.50) 100.0 (38.38-2.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.65 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, $R_{free}$	0.194 , 0.245 0.199 , 0.253	Depositor DCC
$R_{free}$ test set	2669 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	44.0	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 57.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10499	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5XV, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/8323	0.69	0/11279
2	B	0.47	0/2085	0.67	0/2799
All	All	0.50	0/10408	0.68	0/14078

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8126	0	7969	58	0
2	B	2054	0	1957	13	0
3	A	6	8	8	1	0
4	A	27	15	15	0	0
5	A	237	0	0	0	0
5	B	26	0	0	1	0
All	All	10476	23	9949	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:ILE:HD12	1:A:278:MET:HG2	1.71	0.72
1:A:1053:THR:HG23	1:A:1054:LYS:H	1.54	0.71
1:A:873:ASN:HD22	1:A:875:HIS:HB2	1.56	0.70
1:A:25:LEU:HD13	2:B:497:GLN:HG3	1.77	0.67
1:A:492:ILE:HG21	1:A:584:TYR:HD2	1.61	0.66
1:A:5:PRO:HB3	2:B:479:MET:HG2	1.79	0.64
2:B:494:PHE:HB3	2:B:535:ILE:HG12	1.83	0.60
1:A:873:ASN:ND2	1:A:875:HIS:HB2	2.15	0.60
1:A:214:ASP:HA	1:A:266:PRO:HB3	1.84	0.59
1:A:807:LEU:HD12	1:A:846:GLY:HA3	1.85	0.58
2:B:454:THR:O	2:B:458:GLU:HG2	2.03	0.58
1:A:640:LYS:HG2	1:A:680:VAL:HG11	1.88	0.56
1:A:408:SER:HB3	1:A:422:LEU:HD21	1.89	0.55
1:A:552:TRP:HZ3	1:A:583:MET:HE2	1.72	0.54
1:A:42:LEU:HD21	1:A:92:LEU:HD11	1.89	0.54
1:A:744:PHE:CZ	1:A:748:LEU:CD1	2.91	0.54
1:A:956:LEU:HD11	1:A:980:PHE:CZ	2.43	0.54
1:A:562:PRO:HB3	1:A:591:PRO:HG2	1.91	0.52
1:A:897:PHE:O	1:A:901:CYS:HB2	2.09	0.52
1:A:58:LEU:HB3	1:A:61:LEU:HD12	1.91	0.52
1:A:736:VAL:O	1:A:740:ARG:HG3	2.10	0.52
1:A:181:ILE:CD1	1:A:278:MET:HG2	2.40	0.52
1:A:267:LEU:HG	1:A:273:ILE:HG13	1.92	0.52
2:B:581:LEU:O	2:B:585:THR:HG23	2.10	0.51
1:A:157:ASN:HB3	1:A:161:SER:HB3	1.92	0.51
1:A:328:TRP:HA	1:A:394:PRO:HB3	1.93	0.51
1:A:323:SER:O	1:A:482:SER:HB2	2.11	0.51
1:A:962:ILE:HG23	1:A:967:GLY:HA2	1.92	0.50
3:A:1101:GOL:H2	2:B:481:ARG:HH21	1.76	0.50
1:A:744:PHE:CZ	1:A:748:LEU:HD12	2.47	0.49
1:A:120:ALA:O	1:A:672:LYS:HE2	2.11	0.49
1:A:492:ILE:HG21	1:A:584:TYR:CD2	2.46	0.49
1:A:542:GLU:HG3	2:B:340:ARG:HH21	1.77	0.49
1:A:27:PRO:HD3	1:A:101:VAL:HB	1.95	0.48
1:A:638:VAL:HG23	1:A:649:LEU:HD21	1.95	0.48
1:A:175:PRO:HB3	1:A:266:PRO:HG2	1.94	0.48
1:A:279:LEU:HD13	1:A:281:ARG:HE	1.79	0.47
1:A:363:GLY:N	1:A:607:PRO:HG3	2.29	0.47
2:B:398:PHE:HZ	2:B:407:HIS:CD2	2.32	0.47
1:A:1027:ALA:HB1	1:A:1030:LYS:HD2	1.96	0.47
1:A:294:TYR:HA	1:A:297:LEU:HD12	1.97	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:410:ASN:HB2	5:B:723:HOH:O	2.14	0.47
2:B:453:ASN:O	2:B:457:GLN:HG2	2.14	0.47
1:A:916:ARG:HB3	1:A:921:ILE:HD11	1.97	0.47
1:A:209:LEU:HD22	1:A:223:GLU:HG3	1.97	0.47
2:B:354:THR:HA	2:B:426:TYR:O	2.14	0.46
1:A:488:ASP:O	1:A:492:ILE:HG12	2.14	0.46
1:A:1002:PHE:HD1	1:A:1013:LEU:HD11	1.81	0.46
2:B:387:ASP:HB2	2:B:389:LYS:H	1.81	0.45
1:A:402:LEU:HB3	1:A:427:ILE:HG23	1.99	0.45
1:A:95:PHE:CD2	1:A:96:GLN:HG2	2.52	0.45
1:A:543:ILE:HD11	1:A:567:LYS:HD3	2.00	0.44
1:A:71:VAL:HG22	1:A:81:GLU:HA	1.99	0.44
1:A:240:LYS:O	1:A:244:LEU:HG	2.17	0.44
1:A:824:TRP:HB3	1:A:829:LEU:HB2	2.00	0.43
1:A:999:ILE:HD11	1:A:1028:LEU:CD1	2.48	0.43
1:A:559:VAL:HG13	1:A:591:PRO:HD3	2.01	0.43
1:A:194:ILE:HD13	1:A:209:LEU:HD12	2.01	0.42
1:A:785:ASN:HB2	1:A:797[B]:ASN:HD21	1.84	0.42
1:A:193:VAL:HG22	1:A:208:THR:HG22	2.01	0.42
1:A:356:VAL:HG22	1:A:406:ILE:HG12	2.03	0.41
1:A:360:ILE:HG22	1:A:367:LEU:HD12	2.02	0.41
1:A:440:LYS:HG3	1:A:476:GLU:HG3	2.01	0.41
1:A:146:VAL:HG21	1:A:651:ARG:HG2	2.01	0.41
2:B:559:ILE:O	2:B:563:MET:HG3	2.21	0.41
1:A:324:THR:HG22	1:A:483:VAL:HB	2.03	0.40
1:A:180:HIS:ND1	1:A:828:GLY:HA2	2.35	0.40
1:A:959:ASP:O	1:A:963:VAL:HG23	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1004/1074 (94%)	959 (96%)	43 (4%)	2 (0%)	52	75
2	B	237/323 (73%)	232 (98%)	5 (2%)	0	100	100
All	All	1241/1397 (89%)	1191 (96%)	48 (4%)	2 (0%)	52	75

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	946	GLY
1	A	412	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	888/980 (91%)	880 (99%)	8 (1%)	84	95
2	B	215/301 (71%)	211 (98%)	4 (2%)	65	87
All	All	1103/1281 (86%)	1091 (99%)	12 (1%)	80	94

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	PHE
1	A	206	LYS
1	A	236	SER
1	A	427	ILE
1	A	526	ASN
1	A	901	CYS
1	A	937	PHE
1	A	939	ASP
2	B	387	ASP
2	B	416	TYR
2	B	509	ILE
2	B	562	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	873	ASN
2	B	415	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	1101	-	5,5,5	0.10	0	5,5,5	0.40	0
4	5XV	A	1102	-	25,29,29	2.85	5 (20%)	25,43,43	1.90	7 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1101	-	-	0/4/4/4	0/0/0/0
4	5XV	A	1102	-	-	0/16/18/18	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1102	5XV	C5-N9	-10.86	1.23	1.33
4	A	1102	5XV	C3-N4	-3.43	1.31	1.37
4	A	1102	5XV	C8-N10	-2.72	1.35	1.40
4	A	1102	5XV	C6-C5	4.33	1.47	1.40
4	A	1102	5XV	C1-C2	4.59	1.49	1.39

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1102	5XV	C1-C6-C5	-3.86	114.98	119.70
4	A	1102	5XV	F26-C22-C21	-3.47	105.64	112.19
4	A	1102	5XV	O14-C12-C13	-2.95	116.63	122.07
4	A	1102	5XV	C19-C11-N15	-2.43	117.31	120.02
4	A	1102	5XV	O14-C12-N10	2.48	126.11	123.08
4	A	1102	5XV	C2-C11-N15	2.90	120.10	116.00
4	A	1102	5XV	C11-N15-C16	3.00	121.82	117.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1101	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1009/1074 (93%)	-0.06	37 (3%) 45 50	25, 47, 82, 116	0
2	B	249/323 (77%)	0.44	21 (8%) 14 14	33, 62, 93, 107	0
All	All	1258/1397 (90%)	0.04	58 (4%) 36 41	25, 50, 88, 116	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	242	CYS	6.1
2	B	400	SER	5.5
1	A	231	SER	5.0
1	A	1057	TRP	4.6
1	A	872	PHE	4.4
2	B	426	TYR	4.3
1	A	947	TYR	4.2
2	B	440	ASP	4.2
1	A	236	SER	4.2
1	A	244	LEU	4.1
2	B	422	VAL	4.1
1	A	157	ASN	4.0
2	B	431	TYR	3.7
2	B	583	TRP	3.7
2	B	399	SER	3.5
1	A	238	GLN	3.4
2	B	401	VAL	3.4
1	A	526	ASN	3.4
1	A	879	GLN	3.3
2	B	398	PHE	3.2
1	A	561	ILE	3.1
1	A	181	ILE	3.1
1	A	207	TYR	3.0
1	A	239	LEU	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	246	TYR	2.9
1	A	557	TYR	2.9
2	B	406	ASN	2.8
1	A	1060	HIS	2.8
2	B	588	GLY	2.7
2	B	390	TYR	2.6
2	B	397	THR	2.6
1	A	247	GLN	2.6
1	A	871	GLN	2.6
1	A	411	GLY	2.6
1	A	229	THR	2.6
2	B	415	GLN	2.6
1	A	527	ASP	2.4
2	B	404	LEU	2.4
1	A	479	TRP	2.4
1	A	793	LEU	2.3
1	A	1059	PHE	2.3
1	A	945	PHE	2.3
1	A	237	GLU	2.2
2	B	384	PHE	2.2
1	A	230	ARG	2.2
1	A	241	LEU	2.2
2	B	408	TYR	2.2
2	B	403	GLU	2.2
2	B	590	ARG	2.2
1	A	883	ASP	2.1
1	A	560	THR	2.1
1	A	328	TRP	2.1
1	A	228	LYS	2.1
1	A	529	GLU	2.1
2	B	402	VAL	2.1
1	A	54	ARG	2.1
1	A	187	LYS	2.0
2	B	442	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	1101	6/6	0.88	0.18	0.68	65,68,70,70	0
4	5XV	A	1102	27/27	0.97	0.14	-0.31	24,33,53,56	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.