



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 17, 2016 – 02:29 PM EDT

PDB ID : 5FL7  
Title : Structure of the F1c10 complex from *Yarrowia lipolytica* ATP synthase  
Authors : Parey, K.; Bublitz, M.; Meier, T.  
Deposited on : 2015-10-22  
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1051 (3.60-3.40)
Clashscore	102246	1157 (3.60-3.40)
Ramachandran outliers	100387	1120 (3.60-3.40)
Sidechain outliers	100360	1121 (3.60-3.40)
RSRZ outliers	91569	1058 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	536	<div> <div>3%</div> <div>73%</div> <div>17%</div> <div>10%</div> </div>
1	B	536	<div> <div>6%</div> <div>73%</div> <div>16%</div> <div>11%</div> </div>
1	C	536	<div> <div>5%</div> <div>76%</div> <div>14%</div> <div>10%</div> </div>
2	D	509	<div> <div>5%</div> <div>76%</div> <div>16%</div> <div>8%</div> </div>
2	E	509	<div> <div>8%</div> <div>74%</div> <div>18%</div> <div>8%</div> </div>
2	F	509	<div> <div>3%</div> <div>78%</div> <div>15%</div> <div>8%</div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	G	293	
4	H	137	
5	I	16	
6	K	76	
6	L	76	
6	M	76	
6	N	76	
6	O	76	
6	P	76	
6	Q	76	
6	R	76	
6	S	76	
6	T	76	

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 30119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP SYNTHASE SUBUNIT ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	0	0
			3699	2341	645	704	9			
1	B	479	Total	C	N	O	S	0	0	0
			3653	2310	638	696	9			
1	C	485	Total	C	N	O	S	0	0	0
			3699	2341	645	704	9			

- Molecule 2 is a protein called ATP SYNTHASE SUBUNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	470	Total	C	N	O	S	0	0	0
			3535	2226	605	696	8			
2	E	468	Total	C	N	O	S	0	0	0
			3524	2219	603	694	8			
2	F	470	Total	C	N	O	S	0	0	0
			3535	2226	605	696	8			

- Molecule 3 is a protein called ATP SYNTHASE SUBUNIT GAMMA CHAIN, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	267	Total	C	N	O	S	0	0	0
			2068	1291	357	411	9			

- Molecule 4 is a protein called ATP SYNTHASE DELTA CHAIN, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	113	Total	C	N	O	0	0	0
			843	525	135	183			


- Molecule 5 is a protein called ATP SYNTHASE EPSILON CHAIN, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	I	16	Total	C	N	O	0	0	0
			80	48	16	16			

- Molecule 6 is a protein called ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	K	73	Total	C	N	O	S	0	0	0
			524	353	82	88	1			
6	L	72	Total	C	N	O	S	0	0	0
			512	344	81	86	1			
6	M	75	Total	C	N	O	S	0	0	0
			536	361	84	90	1			
6	N	75	Total	C	N	O	S	0	0	0
			537	361	84	90	2			
6	O	75	Total	C	N	O	S	0	0	0
			537	361	84	91	1			
6	P	75	Total	C	N	O	S	0	0	0
			537	361	84	90	2			
6	Q	76	Total	C	N	O	S	0	0	0
			544	366	85	91	2			
6	R	74	Total	C	N	O	S	0	0	0
			529	356	83	89	1			
6	S	74	Total	C	N	O	S	0	0	0
			529	356	83	89	1			
6	T	74	Total	C	N	O	S	0	0	0
			529	356	83	89	1			

- Molecule 7 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
7	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
7	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Mg	0	0
			1	1		
8	A	1	Total	Mg	0	0
			1	1		
8	D	1	Total	Mg	0	0
			1	1		
8	C	1	Total	Mg	0	0
			1	1		
8	F	1	Total	Mg	0	0
			1	1		

- Molecule 9 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
9	F	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	3	Total	O	0	0
			3	3		
10	B	3	Total	O	0	0
			3	3		
10	C	3	Total	O	0	0
			3	3		
10	D	4	Total	O	0	0
			4	4		
10	F	4	Total	O	0	0
			4	4		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ATP SYNTHASE SUBUNIT ALPHA




#### • Molecule 1: ATP SYNTHASE SUBUNIT ALPHA




#### • Molecule 1: ATP SYNTHASE SUBUNIT ALPHA







• Molecule 2: ATP SYNTHASE SUBUNIT BETA




• Molecule 2: ATP SYNTHASE SUBUNIT BETA







• Molecule 2: ATP SYNTHASE SUBUNIT BETA



• Molecule 3: ATP SYNTHASE SUBUNIT GAMMA CHAIN, MITOCHONDRIAL



• Molecule 4: ATP SYNTHASE DELTA CHAIN, MITOCHONDRIAL




• Molecule 5: ATP SYNTHASE EPSILON CHAIN, MITOCHONDRIAL




There are no outlier residues recorded for this chain.


• Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL




- Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL



● Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL



● Molecule 6: ATP SYNTHASE SUBUNIT 9, MITOCHONDRIAL



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	169.50 Å   182.20 Å   193.00 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	49.37 – 3.50 49.37 – 3.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.37-3.50) 100.0 (49.37-3.50)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.07 (at 3.48 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.274   ,   0.305 0.273   ,   0.305	Depositor DCC
$R_{free}$ test set	1894 reflections (2.50%)	DCC
Wilson B-factor (Å <sup>2</sup> )	157.5	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 94.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	30119	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	166.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.24	0/3755	0.41	0/5076
1	B	0.24	0/3706	0.41	0/5009
1	C	0.24	0/3755	0.40	0/5076
2	D	0.24	0/3587	0.40	0/4868
2	E	0.24	0/3576	0.41	0/4853
2	F	0.24	0/3587	0.40	0/4868
3	G	0.23	0/2090	0.38	0/2812
4	H	0.24	0/851	0.43	0/1157
6	K	0.24	0/532	0.36	0/720
6	L	0.24	0/519	0.35	0/702
6	M	0.24	0/544	0.36	0/737
6	N	0.24	0/545	0.35	0/737
6	O	0.25	0/545	0.38	0/737
6	P	0.24	0/545	0.33	0/737
6	Q	0.25	0/552	0.38	0/747
6	R	0.24	0/537	0.35	0/727
6	S	0.25	0/537	0.33	0/727
6	T	0.25	0/537	0.38	0/727
All	All	0.24	0/30300	0.40	0/41017

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3699	0	3781	59	0
1	B	3653	0	3741	52	0
1	C	3699	0	3781	49	0
2	D	3535	0	3562	60	0
2	E	3524	0	3551	58	0
2	F	3535	0	3562	49	0
3	G	2068	0	2099	31	0
4	H	843	0	824	18	0
5	I	80	0	18	0	0
6	K	524	0	565	20	0
6	L	512	0	556	15	0
6	M	536	0	579	15	0
6	N	537	0	582	8	0
6	O	537	0	579	10	0
6	P	537	0	582	15	0
6	Q	544	0	591	15	0
6	R	529	0	570	15	0
6	S	529	0	570	22	0
6	T	529	0	570	22	0
7	A	31	0	12	0	0
7	B	31	0	12	0	0
7	C	31	0	12	2	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
8	F	1	0	0	0	0
9	D	27	0	12	3	0
9	F	27	0	12	3	0
10	A	3	0	0	0	0
10	B	3	0	0	0	0
10	C	3	0	0	0	0
10	D	4	0	0	0	0
10	F	4	0	0	2	0
All	All	30119	0	30723	432	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 432 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:329:SER:HB2	2:F:253:MET:HB3	1.66	0.77
3:G:208:ALA:HB3	3:G:211:PHE:HB2	1.71	0.73
2:D:189:GLY:O	2:D:194:LYS:NZ	2.22	0.73
1:C:370:SER:HB3	2:D:291:ARG:HH22	1.57	0.70
1:B:450:LEU:HD21	1:B:474:GLY:HA3	1.73	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	483/536 (90%)	456 (94%)	26 (5%)	1 (0%)	52	88
1	B	475/536 (89%)	454 (96%)	21 (4%)	0	100	100
1	C	483/536 (90%)	456 (94%)	27 (6%)	0	100	100
2	D	468/509 (92%)	448 (96%)	20 (4%)	0	100	100
2	E	466/509 (92%)	443 (95%)	23 (5%)	0	100	100
2	F	468/509 (92%)	439 (94%)	28 (6%)	1 (0%)	52	88
3	G	263/293 (90%)	242 (92%)	21 (8%)	0	100	100
4	H	109/137 (80%)	92 (84%)	13 (12%)	4 (4%)	4	36
6	K	71/76 (93%)	71 (100%)	0	0	100	100
6	L	70/76 (92%)	67 (96%)	3 (4%)	0	100	100
6	M	73/76 (96%)	71 (97%)	2 (3%)	0	100	100
6	N	73/76 (96%)	71 (97%)	2 (3%)	0	100	100
6	O	73/76 (96%)	73 (100%)	0	0	100	100
6	P	73/76 (96%)	72 (99%)	1 (1%)	0	100	100
6	Q	74/76 (97%)	71 (96%)	3 (4%)	0	100	100
6	R	72/76 (95%)	72 (100%)	0	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
6	S	72/76 (95%)	70 (97%)	2 (3%)	0	100	100
6	T	72/76 (95%)	71 (99%)	1 (1%)	0	100	100
All	All	3938/4325 (91%)	3739 (95%)	193 (5%)	6 (0%)	52	88

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	75	ASN
4	H	120	ALA
4	H	45	PRO
1	A	400	VAL
4	H	78	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/429 (91%)	389 (100%)	2 (0%)	92	97
1	B	387/429 (90%)	383 (99%)	4 (1%)	82	93
1	C	391/429 (91%)	391 (100%)	0	100	100
2	D	377/409 (92%)	377 (100%)	0	100	100
2	E	376/409 (92%)	376 (100%)	0	100	100
2	F	377/409 (92%)	377 (100%)	0	100	100
3	G	224/242 (93%)	221 (99%)	3 (1%)	76	91
4	H	92/114 (81%)	92 (100%)	0	100	100
6	K	50/52 (96%)	50 (100%)	0	100	100
6	L	49/52 (94%)	48 (98%)	1 (2%)	63	87
6	M	51/52 (98%)	51 (100%)	0	100	100
6	N	51/52 (98%)	50 (98%)	1 (2%)	63	87
6	O	51/52 (98%)	49 (96%)	2 (4%)	39	76
6	P	51/52 (98%)	50 (98%)	1 (2%)	63	87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	Q	52/52 (100%)	52 (100%)	0	100	100
6	R	50/52 (96%)	49 (98%)	1 (2%)	63	87
6	S	50/52 (96%)	49 (98%)	1 (2%)	63	87
6	T	50/52 (96%)	49 (98%)	1 (2%)	63	87
All	All	3120/3390 (92%)	3103 (100%)	17 (0%)	92	97

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	149	PHE
3	G	176	TYR
6	P	1	MET
3	G	78	THR
6	R	55	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	203	GLN
6	S	46	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	ATP	A	600	8	26,33,33	0.97	1 (3%)	26,52,52	1.66	1 (3%)
7	ATP	B	600	8	26,33,33	0.97	1 (3%)	26,52,52	1.63	1 (3%)
7	ATP	C	600	8	26,33,33	0.96	1 (3%)	26,52,52	1.65	1 (3%)
9	ADP	D	600	8	24,29,29	0.99	1 (4%)	23,45,45	1.74	1 (4%)
9	ADP	F	600	8	24,29,29	0.99	1 (4%)	23,45,45	1.71	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ATP	A	600	8	-	0/18/38/38	0/3/3/3
7	ATP	B	600	8	-	0/18/38/38	0/3/3/3
7	ATP	C	600	8	-	0/18/38/38	0/3/3/3
9	ADP	D	600	8	-	0/12/32/32	0/3/3/3
9	ADP	F	600	8	-	0/12/32/32	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	D	600	ADP	C5-C4	3.11	1.47	1.40
7	C	600	ATP	C5-C4	3.11	1.47	1.40
9	F	600	ADP	C5-C4	3.12	1.47	1.40
7	B	600	ATP	C5-C4	3.15	1.47	1.40
7	A	600	ATP	C5-C4	3.16	1.47	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	D	600	ADP	N3-C2-N1	-6.63	123.66	128.87
7	A	600	ATP	N3-C2-N1	-6.59	123.70	128.87
7	B	600	ATP	N3-C2-N1	-6.55	123.72	128.87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	600	ATP	N3-C2-N1	-6.55	123.73	128.87
9	F	600	ADP	N3-C2-N1	-6.51	123.76	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	600	ATP	2	0
9	D	600	ADP	3	0
9	F	600	ADP	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	485/536 (90%)	0.11	17 (3%)	48	38	116, 153, 209, 224	0
1	B	479/536 (89%)	0.26	30 (6%)	23	18	104, 156, 234, 261	0
1	C	485/536 (90%)	0.21	28 (5%)	26	21	108, 150, 222, 280	0
2	D	470/509 (92%)	0.28	26 (5%)	29	22	108, 158, 195, 215	0
2	E	468/509 (91%)	0.34	42 (8%)	12	10	117, 160, 210, 248	0
2	F	470/509 (92%)	0.07	15 (3%)	51	42	113, 152, 192, 213	0
3	G	267/293 (91%)	0.70	44 (16%)	2	3	109, 185, 224, 241	0
4	H	113/137 (82%)	0.92	24 (21%)	1	1	161, 207, 255, 277	0
5	I	0/16	-	-			-	-
6	K	73/76 (96%)	0.45	12 (16%)	2	3	148, 184, 206, 218	0
6	L	72/76 (94%)	0.10	6 (8%)	14	12	154, 180, 213, 224	0
6	M	75/76 (98%)	0.17	10 (13%)	4	5	152, 179, 200, 213	0
6	N	75/76 (98%)	-0.31	0	100	100	140, 157, 188, 194	0
6	O	75/76 (98%)	0.15	3 (4%)	42	33	132, 158, 186, 192	0
6	P	75/76 (98%)	0.13	6 (8%)	15	12	132, 160, 200, 205	0
6	Q	76/76 (100%)	0.18	4 (5%)	30	23	150, 177, 198, 204	0
6	R	74/76 (97%)	-0.17	1 (1%)	78	68	162, 182, 197, 213	0
6	S	74/76 (97%)	0.15	7 (9%)	10	10	164, 179, 203, 217	0
6	T	74/76 (97%)	0.10	4 (5%)	29	23	163, 182, 205, 210	0
All	All	3980/4341 (91%)	0.24	279 (7%)	19	15	104, 163, 216, 280	0

The worst 5 of 279 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	GLN	9.8
3	G	116	ASP	8.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
4	H	61	ASN	8.4
1	C	515	LEU	7.3
2	E	396	GLN	6.9

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
9	ADP	F	600	27/27	0.92	0.30	-0.11	159,176,182,186	0
7	ATP	B	600	31/31	0.89	0.29	-0.23	149,182,193,201	0
7	ATP	C	600	31/31	0.92	0.26	-0.23	149,159,174,183	0
7	ATP	A	600	31/31	0.92	0.24	-0.48	146,160,172,197	0
9	ADP	D	600	27/27	0.96	0.26	-0.56	159,164,170,173	0
8	MG	F	601	1/1	0.93	0.13	-3.14	189,189,189,189	0
8	MG	D	601	1/1	0.83	0.17	-	183,183,183,183	0
8	MG	A	601	1/1	0.96	0.20	-	173,173,173,173	0
8	MG	B	601	1/1	0.98	0.18	-	154,154,154,154	0
8	MG	C	601	1/1	0.90	0.22	-	171,171,171,171	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.