



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:26 PM GMT

PDB ID : 1FM9  
Title : THE 2.1 ANGSTROM RESOLUTION CRYSTAL STRUCTURE OF THE HETERODIMER OF THE HUMAN RXRALPHA AND PPARGAMMA LIGAND BINDING DOMAINS RESPECTIVELY BOUND WITH 9-CIS RETINOIC ACID AND GI262570 AND CO-ACTIVATOR PEPTIDES.  
Authors : Gampe Jr., R.T.; Montana, V.G.; Lambert, M.H.; Miller, A.B.; Bledsoe, R.K.; Milburn, M.V.; Kliewer, S.A.; Willson, T.M.; Xu, H.E.  
Deposited on : 2000-08-16  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	<div> <div>25%</div> <div>62%</div> <div>30%</div> <div>5%</div> <div>• •</div> </div>
2	D	272	<div> <div>16%</div> <div>67%</div> <div>28%</div> <div>5%</div> </div>
3	B	25	<div> <div>12%</div> <div>20%</div> <div>16%</div> <div>•</div> <div>60%</div> </div>
3	E	25	<div> <div>48%</div> <div>24%</div> <div>32%</div> <div>8%</div> <div>36%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RETINOIC ACID RECEPTOR RXR-ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	0	0	0
			1754	1117	306	321	10			


- Molecule 2 is a protein called PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR GAMMA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	272	Total	C	N	O	S	0	0	0
			2179	1407	355	407	10			

- Molecule 3 is a protein called STEROID RECEPTOR COACTIVATOR.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	B	10	Total	C	N	O	0	0	0
			86	55	18	13			
3	E	16	Total	C	N	O	0	0	0
			129	79	27	23			

- Molecule 4 is (9CIS)-RETINOIC ACID (three-letter code: 9CR) (formula: C<sub>20</sub>H<sub>28</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			22	20	2		

- Molecule 5 is 2-(2-BENZOYL-PHENYLAMINO)-3-{4-[2-(5-METHYL-2-PHENYL-OXAZOL-4-YL)-ETHOXY]-PHENYL}-PROPIONIC ACID (three-letter code: 570) (formula:  $C_{34}H_{30}N_2O_5$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			41	34	2	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	36	Total 36	O 36	0	0
6	D	89	Total 89	O 89	0	0
6	E	6	Total 6	O 6	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: RETINOIC ACID RECEPTOR RXR-ALPHA




#### • Molecule 2: PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR GAMMA



#### • Molecule 3: STEROID RECEPTOR COACTIVATOR



#### • Molecule 3: STEROID RECEPTOR COACTIVATOR



CYS	PRO	SER	SER	HIS	SER	SER	LEU	THR	E685	E686	H687	K688	I689	L690	H691	R692	L693	L694	Q695	E696	G697	S698	P699	S700
-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.14Å 54.05Å 211.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.10 19.69 – 2.10	Depositor EDS
% Data completeness (in resolution range)	91.8 (20.00-2.10) 91.9 (19.69-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.14 (at 2.11Å)	Xtriage
Refinement program	CNS 0.5	Depositor
R, $R_{free}$	0.239 , 0.268 0.239 , 0.268	Depositor DCC
$R_{free}$ test set	2895 reflections (9.95%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.3	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 57.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 29087 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4342	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9CR, 570

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/1789	0.64	2/2414 (0.1%)
2	D	0.42	0/2217	0.71	1/2987 (0.0%)
3	B	0.38	0/87	0.72	0/116
3	E	0.28	0/131	0.52	0/175
All	All	0.40	0/4224	0.68	3/5692 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	GLU	C-N-CD	-10.66	97.14	120.60
1	A	243	GLU	C-N-CA	6.33	148.58	122.00
2	D	206	PRO	N-CA-CB	5.39	109.77	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1754	0	1733	80	0
2	D	2179	0	2237	99	0
3	B	86	0	86	7	0
3	E	129	0	125	15	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	22	0	27	4	0
5	D	41	0	29	5	0
6	A	36	0	0	3	0
6	D	89	0	0	2	0
6	E	6	0	0	1	0
All	All	4342	0	4237	187	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 187 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:268:THR:H	2:D:269:PRO:HD2	1.25	0.99
2:D:293:VAL:HG22	2:D:322:VAL:HG11	1.57	0.86
1:A:306:ASN:HD22	1:A:306:ASN:H	1.24	0.85
1:A:230:MET:CE	1:A:235:ILE:HD11	2.07	0.84
1:A:236:LEU:HD22	1:A:240:LEU:HD22	1.58	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/238 (97%)	196 (85%)	20 (9%)	14 (6%)	2	0
2	D	270/272 (99%)	255 (94%)	7 (3%)	8 (3%)	5	2
3	B	8/25 (32%)	7 (88%)	1 (12%)	0	100	100
3	E	14/25 (56%)	10 (71%)	3 (21%)	1 (7%)	1	0
All	All	522/560 (93%)	468 (90%)	31 (6%)	23 (4%)	3	1

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	GLU
1	A	244	PRO
1	A	257	ASN
1	A	259	SER
1	A	260	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/205 (89%)	167 (92%)	15 (8%)	14	10
2	D	243/245 (99%)	224 (92%)	19 (8%)	16	11
3	B	9/24 (38%)	8 (89%)	1 (11%)	8	4
3	E	14/24 (58%)	13 (93%)	1 (7%)	18	14
All	All	448/498 (90%)	412 (92%)	36 (8%)	15	11

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	228	LEU
2	D	242	THR
2	D	477	TYR
2	D	237	LEU
2	D	244	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	375	ASN
2	D	424	ASN
3	B	634	HIS
2	D	335	ASN
3	B	638	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	9CR	A	201	-	19,22,22	3.81	9 (47%)	26,30,30	2.55	11 (42%)
5	570	D	200	-	38,45,45	2.39	18 (47%)	48,61,61	1.77	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	9CR	A	201	-	-	0/13/32/32	0/1/1/1
5	570	D	200	-	-	0/23/30/30	0/4/5/5

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	200	570	CB-CA	2.15	1.56	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	200	570	C3J-C3I	2.16	1.43	1.39
5	D	200	570	OH-CZ	2.33	1.43	1.37
5	D	200	570	CD1-CG	2.35	1.43	1.38
5	D	200	570	CD2-CG	2.37	1.43	1.38

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	201	9CR	C16-C1-C6	-4.91	102.61	110.30
5	D	200	570	O1G-C1G-C1H	-4.61	112.75	120.12
4	A	201	9CR	C19-C9-C10	-4.14	116.79	122.90
4	A	201	9CR	C11-C10-C9	-3.51	122.12	127.20
4	A	201	9CR	C1-C6-C5	-2.83	118.51	122.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	201	9CR	4	0
5	D	200	570	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/238 (97%)	2.02	59 (25%) <b>1</b> <b>1</b>	23, 42, 109, 124	0
2	D	272/272 (100%)	1.42	43 (15%) <b>3</b> <b>4</b>	26, 39, 115, 131	0
3	B	10/25 (40%)	1.71	3 (30%) <b>1</b> <b>1</b>	50, 57, 69, 76	0
3	E	16/25 (64%)	3.84	12 (75%) <b>0</b> <b>0</b>	60, 68, 96, 98	0
All	All	530/560 (94%)	1.76	117 (22%) <b>1</b> <b>1</b>	23, 41, 110, 131	0

The worst 5 of 117 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	267	ILE	33.0
1	A	252	ALA	18.8
1	A	256	LEU	18.7
2	D	274	SER	18.6
2	D	266	HIS	17.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	9CR	A	201	22/22	0.88	0.25	1.80	43,48,50,50	0
5	570	D	200	41/41	0.93	0.17	0.30	33,38,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.