



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:26 PM GMT

PDB ID : 1FNQ
Title : CRYSTAL STRUCTURE ANALYSIS OF THE MUTANT REACTION CENTER PRO L209-> GLU FROM THE PHOTOSYNTHETIC PURPLE BACTERIUM RHODOBACTER SPHAEROIDES
Authors : Kuglstatter, A.; Ermler, U.; Michel, H.; Baciou, L.; Fritzsche, G.
Deposited on : 2000-08-23
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	281	
2	M	307	
3	H	260	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	BCL	L	304	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	BCL	M	801	X	-	-	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 7248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called REACTION CENTER PROTEIN L CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	281	Total	C	N	O	S	0	0	0
			2234	1507	355	364	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	209	GLU	PRO	ENGINEERED	UNP P02954

- Molecule 2 is a protein called REACTION CENTER PROTEIN M CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	302	Total	C	N	O	S	0	0	0
			2404	1603	394	397	10			


- Molecule 3 is a protein called REACTION CENTER PROTEIN H CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	239	Total	C	N	O	S	0	0	0
			1823	1166	313	335	9			

- Molecule 4 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	M	1	Total	Fe	0	0
			1	1		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	M	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula: $\text{C}_{55}\text{H}_{74}\text{MgN}_4\text{O}_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	M	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
6	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
6	M	1	Total 66	C 55	Mg 1	N 4	O 6	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 7 is BACTERIOPHEOPHYTIN A (three-letter code: BPH) (formula: $C_{55}H_{76}N_4O_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	M	1	Total	C	N	O	0	0
			65	55	4	6		
7	L	1	Total	C	N	O	0	0
			65	55	4	6		

- Molecule 8 is UBIQUINONE-10 (three-letter code: U10) (formula: $C_{59}H_{90}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	M	1	Total	C	O	0	0
			48	44	4		
8	L	1	Total	C	O	0	0
			48	44	4		


- Molecule 9 is SPHEROIDENE (three-letter code: SPO) (formula: $C_{41}H_{60}O$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	M	1	Total	C	O	0	0
			42	41	1		

- Molecule 10 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula:

C₁₄H₃₁NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	M	1	Total	C	N	O	0	0
			16	14	1	1		
10	H	1	Total	C	N	O	0	0
			16	14	1	1		
10	M	1	Total	C	N	O	0	0
			16	14	1	1		
10	M	1	Total	C	N	O	0	0
			16	14	1	1		
10	L	1	Total	C	N	O	0	0
			16	14	1	1		
10	H	1	Total	C	N	O	0	0
			16	14	1	1		
10	M	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 11 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	H	64	Total	O	0	0
			64	64		
11	L	34	Total	O	0	0
			34	34		
11	M	39	Total	O	0	0
			39	39		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: REACTION CENTER PROTEIN L CHAIN



• Molecule 2: REACTION CENTER PROTEIN M CHAIN



• Molecule 3: REACTION CENTER PROTEIN H CHAIN



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	141.53Å 141.53Å 187.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.60	Depositor
% Data completeness (in resolution range)	93.1 (50.00-2.60)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 0.3	Depositor
R, R_{free}	0.217 , 0.247	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7248	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BCL, LDA, BPH, PO4, FE, SPO, U10

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.44	0/2321	0.49	0/3175
2	M	0.44	0/2496	0.49	1/3407 (0.0%)
3	H	0.41	0/1871	0.51	0/2545
All	All	0.43	0/6688	0.50	1/9127 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	M	0	2
3	H	0	1
All	All	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	214	LEU	CA-CB-CG	5.86	128.78	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	H	118	ARG	Sidechain
2	M	198	TYR	Sidechain
2	M	48	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	2234	0	2186	39	0
2	M	2404	0	2311	47	0
3	H	1823	0	1831	27	0
4	M	1	0	0	0	0
5	M	5	0	0	0	0
6	L	132	0	148	12	0
6	M	132	0	148	17	0
7	L	65	0	76	8	0
7	M	65	0	76	11	0
8	L	48	0	63	2	0
8	M	48	0	63	2	0
9	M	42	0	60	2	0
10	H	32	0	62	3	0
10	L	16	0	31	0	0
10	M	64	0	124	4	0
11	H	64	0	0	5	0
11	L	34	0	0	1	0
11	M	39	0	0	1	0
All	All	7248	0	7179	135	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 135 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:L:304:BCL:HBB2	6:L:304:BCL:HHC	1.43	1.01
6:M:801:BCL:HHC	6:M:801:BCL:HBB2	1.43	0.98
7:L:402:BPH:HHC	7:L:402:BPH:HBB3	1.51	0.92
1:L:202:LYS:HG3	1:L:203:GLY:H	1.36	0.90
1:L:241:VAL:HG21	7:L:402:BPH:HAC2	1.57	0.85

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	279/281 (99%)	266 (95%)	12 (4%)	1 (0%)	39	65
2	M	300/307 (98%)	281 (94%)	17 (6%)	2 (1%)	26	51
3	H	237/260 (91%)	227 (96%)	9 (4%)	1 (0%)	39	65
All	All	816/848 (96%)	774 (95%)	38 (5%)	4 (0%)	34	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	M	301	HIS
3	H	116	ALA
1	L	202	LYS
2	M	48	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	220/220 (100%)	203 (92%)	17 (8%)	16	31
2	M	235/240 (98%)	212 (90%)	23 (10%)	10	19
3	H	194/208 (93%)	182 (94%)	12 (6%)	23	45
All	All	649/668 (97%)	597 (92%)	52 (8%)	15	29

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	133	THR
2	M	182	HIS
3	H	221	SER
2	M	136	ARG
2	M	156	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	M	44	ASN
3	H	206	ASN
2	M	199	ASN
1	L	183	ASN
2	M	193	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 1 is monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	LDA	H	702	-	15,15,15	4.69	3 (20%)	16,17,17	0.60	0
10	LDA	H	706	-	15,15,15	4.36	3 (20%)	16,17,17	0.55	0
6	BCL	L	302	1	53,74,74	1.22	5 (9%)	57,115,115	1.61	9 (15%)
6	BCL	L	304	1	53,74,74	1.35	4 (7%)	57,115,115	2.08	7 (12%)
7	BPH	L	402	-	64,70,70	1.22	5 (7%)	73,101,101	1.77	10 (13%)
8	U10	L	502	-	48,48,63	1.87	13 (27%)	58,61,79	1.04	5 (8%)
10	LDA	L	705	-	15,15,15	4.36	1 (6%)	16,17,17	0.49	0
7	BPH	M	401	-	64,70,70	1.19	6 (9%)	73,101,101	1.80	12 (16%)
8	U10	M	501	-	48,48,63	2.07	14 (29%)	58,61,79	1.01	3 (5%)
9	SPO	M	600	-	40,41,41	3.16	21 (52%)	45,50,50	2.11	12 (26%)
10	LDA	M	701	-	15,15,15	4.45	2 (13%)	16,17,17	0.52	0
10	LDA	M	703	-	15,15,15	4.25	1 (6%)	16,17,17	0.56	0
10	LDA	M	704	-	15,15,15	4.43	4 (26%)	16,17,17	0.72	0
10	LDA	M	707	-	15,15,15	3.44	1 (6%)	16,17,17	0.68	0
5	PO4	M	800	-	4,4,4	1.41	0	6,6,6	0.30	0
6	BCL	M	801	2	53,74,74	1.23	4 (7%)	57,115,115	2.00	7 (12%)
6	BCL	M	802	2	53,74,74	1.18	4 (7%)	57,115,115	1.92	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	LDA	H	702	-	-	0/13/13/13	0/0/0/0
10	LDA	H	706	-	-	0/13/13/13	0/0/0/0
6	BCL	L	302	1	-	0/37/137/137	0/0/9/9
6	BCL	L	304	1	1/1/21/25	0/37/137/137	0/0/9/9
7	BPH	L	402	-	-	0/54/105/105	0/1/6/6
8	U10	L	502	-	-	0/45/69/87	0/1/1/1
10	LDA	L	705	-	-	0/13/13/13	0/0/0/0
7	BPH	M	401	-	-	0/54/105/105	0/1/6/6
8	U10	M	501	-	-	0/45/69/87	0/1/1/1
9	SPO	M	600	-	-	0/47/47/47	0/0/0/0
10	LDA	M	701	-	-	0/13/13/13	0/0/0/0
10	LDA	M	703	-	-	0/13/13/13	0/0/0/0
10	LDA	M	704	-	-	0/13/13/13	0/0/0/0
10	LDA	M	707	-	-	0/13/13/13	0/0/0/0
5	PO4	M	800	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BCL	M	801	2	1/1/21/25	0/37/137/137	0/0/9/9
6	BCL	M	802	2	-	0/37/137/137	0/0/9/9

The worst 5 of 91 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	H	702	LDA	O1-N1	-17.61	1.22	1.39
10	M	701	LDA	O1-N1	-16.87	1.23	1.39
10	L	705	LDA	O1-N1	-16.60	1.23	1.39
10	H	706	LDA	O1-N1	-16.47	1.23	1.39
10	M	703	LDA	O1-N1	-16.34	1.24	1.39

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	L	304	BCL	O1D-CGD-CBD	-7.99	113.17	124.62
6	M	801	BCL	O1D-CGD-CBD	-7.48	113.90	124.62
7	M	401	BPH	O1D-CGD-CBD	-7.30	114.16	124.62
7	L	402	BPH	O1D-CGD-CBD	-7.29	114.17	124.62
6	M	802	BCL	O1D-CGD-CBD	-7.22	114.28	124.62

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	M	801	BCL	C8
6	L	304	BCL	C13

There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 55 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	H	706	LDA	3	0
6	L	302	BCL	7	0
6	L	304	BCL	5	0
7	L	402	BPH	8	0
8	L	502	U10	2	0
7	M	401	BPH	11	0
8	M	501	U10	2	0
9	M	600	SPO	2	0
10	M	701	LDA	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	M	704	LDA	3	0
6	M	801	BCL	7	0
6	M	802	BCL	12	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.