



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:41 AM GMT

PDB ID : 3FN1
Title : E2-RING expansion of the NEDD8 cascade confers specificity to cullin modification.
Authors : Huang, D.T.; Ayrault, O.; Hunt, H.W.; Taherbhoy, A.M.; Duda, D.M.; Scott, D.C.; Borg, L.A.; Neale, G; Murray, P.J.; Roussel, M.F.; Schulman, B.A.
Deposited on : 2008-12-22
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	98	
2	B	167	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2127 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEDD8-activating enzyme E1 catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	92	715	451	120	142	2	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	345	GLY	-	EXPRESSION TAG	UNP Q8TBC4
A	346	SER	-	EXPRESSION TAG	UNP Q8TBC4
A	394	MSE	LEU	ENGINEERED	UNP Q8TBC4

- Molecule 2 is a protein called NEDD8-conjugating enzyme UBE2F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	160	1277	814	217	240	6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	19	GLY	-	EXPRESSION TAG	UNP Q969M7
B	20	SER	-	EXPRESSION TAG	UNP Q969M7

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	32	Total	O	0	0
			32	32		
3	B	103	Total	O	0	0
			103	103		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NEDD8-activating enzyme E1 catalytic subunit

Chain A: 



- Molecule 2: NEDD8-conjugating enzyme UBE2F

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	81.17Å 81.17Å 212.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.88 – 2.50 24.88 – 2.50	Depositor EDS
% Data completeness (in resolution range)	92.5 (24.88-2.50) 89.4 (24.88-2.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 2.50Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.224 , 0.265 0.213 , 0.256	Depositor DCC
R_{free} test set	728 reflections (5.67%)	DCC
Wilson B-factor (Å ²)	55.7	Xtrriage
Anisotropy	0.531	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.7	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	1 of 14601 reflections (0.007%)	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2127	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/723	0.51	0/977
2	B	0.49	2/1309 (0.2%)	0.58	0/1780
All	All	0.48	2/2032 (0.1%)	0.56	0/2757

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	50	CYS	CB-SG	-5.39	1.73	1.81
2	B	167	ARG	CZ-NH1	5.07	1.39	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	715	0	736	14	0
2	B	1277	0	1232	18	0
3	A	32	0	0	1	0
3	B	103	0	0	2	0
All	All	2127	0	1968	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:376:MSE:HE3	1:A:379:PRO:HB3	1.42	1.00
2:B:106:HIS:HD2	2:B:108:ASN:H	1.20	0.87
1:A:368:LEU:HB3	1:A:376:MSE:HG3	1.67	0.77
1:A:376:MSE:HE3	1:A:379:PRO:CB	2.15	0.76
1:A:396:SER:HB2	3:A:14:HOH:O	1.97	0.64
2:B:106:HIS:CD2	2:B:108:ASN:H	2.08	0.63
2:B:167:ARG:NH1	3:B:271:HOH:O	2.36	0.59
2:B:126:ASP:C	2:B:128:THR:H	2.07	0.58
2:B:167:ARG:HH11	2:B:167:ARG:HG2	1.69	0.57
1:A:379:PRO:HD2	1:A:395:GLN:HE21	1.71	0.55
2:B:85:THR:HB	2:B:98:VAL:HG22	1.89	0.55
2:B:33:ARG:HD2	2:B:87:VAL:HG23	1.89	0.55
1:A:382:THR:HB	1:A:389:ASN:HD21	1.72	0.54
1:A:368:LEU:HD13	1:A:376:MSE:HE2	1.88	0.54
2:B:105:TRP:CE2	2:B:175:LYS:HE3	2.43	0.54
1:A:376:MSE:HE1	1:A:425:VAL:HG23	1.89	0.53
1:A:378:SER:OG	1:A:395:GLN:HG3	2.10	0.52
2:B:167:ARG:NH1	2:B:167:ARG:HG2	2.24	0.52
1:A:393:TYR:CE2	1:A:395:GLN:HB3	2.44	0.52
2:B:33:ARG:HG3	2:B:91:TYR:HB3	1.94	0.49
2:B:87:VAL:HG12	2:B:96:PRO:HB3	1.94	0.49
1:A:376:MSE:HE1	1:A:425:VAL:CG2	2.43	0.49
1:A:405:ARG:N	1:A:406:PRO:HD2	2.28	0.48
1:A:371:SER:O	1:A:375:GLN:HG3	2.16	0.46
2:B:69:VAL:HG13	2:B:144:ASN:HD22	1.81	0.46
2:B:34:ASP:O	2:B:38:VAL:HG23	2.16	0.45
2:B:106:HIS:HD2	2:B:108:ASN:N	2.02	0.43
2:B:126:ASP:O	2:B:128:THR:N	2.51	0.42
1:A:355:PHE:O	1:A:440:PHE:HA	2.21	0.41
2:B:85:THR:HG22	2:B:98:VAL:HG13	2.01	0.41
2:B:185:ARG:NH1	3:B:187:HOH:O	2.54	0.40
2:B:164:HIS:CE1	2:B:172:PHE:HA	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/98 (92%)	89 (99%)	1 (1%)	0	100	100
2	B	158/167 (95%)	149 (94%)	8 (5%)	1 (1%)	30	50
All	All	248/265 (94%)	238 (96%)	9 (4%)	1 (0%)	39	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	127	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/86 (96%)	78 (94%)	5 (6%)	24	43
2	B	139/150 (93%)	131 (94%)	8 (6%)	25	45
All	All	222/236 (94%)	209 (94%)	13 (6%)	24	44

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	360	LYS
1	A	403	ARG
1	A	413	LYS
1	A	414	GLU
1	A	425	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	54	VAL
2	B	64	CYS
2	B	85	THR
2	B	92	ASN
2	B	119	LEU
2	B	152	ASN
2	B	167	ARG
2	B	185	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	362	GLN
1	A	389	ASN
1	A	395	GLN
1	A	432	GLN
2	B	106	HIS
2	B	144	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	90/98 (91%)	0.78	13 (14%) 3 3	50, 63, 70, 79	0
2	B	160/167 (95%)	0.49	12 (7%) 17 19	40, 53, 70, 82	0
All	All	250/265 (94%)	0.60	25 (10%) 9 10	40, 57, 70, 82	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	125	ILE	9.9
2	B	124	SER	6.0
2	B	126	ASP	3.8
1	A	350	PRO	3.6
2	B	185	ARG	3.3
2	B	136	LEU	3.2
1	A	440	PHE	3.0
2	B	128	THR	2.8
1	A	358	SER	2.7
1	A	418	VAL	2.7
1	A	372	ALA	2.5
2	B	41	VAL	2.5
2	B	123	HIS	2.5
1	A	377	LYS	2.4
1	A	381	ILE	2.4
1	A	393	TYR	2.4
1	A	419	ASP	2.3
1	A	441	THR	2.3
1	A	382	THR	2.3
2	B	169	LYS	2.2
2	B	143	LEU	2.1
1	A	375	GLN	2.1
2	B	167	ARG	2.1
1	A	392	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	83	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.