



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:41 AM GMT

PDB ID : 3FNJ
Title : Crystal structure of the full-length lp_1913 protein from Lactobacillus plantarum, Northeast Structural Genomics Consortium Target LpR140
Authors : Seetharaman, J.; Chen, Y.; Forouhar, F.; Sahdev, S.; Janjua, H.; Xiao, R.; Ciccocanti, C.; Foote, E.L.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2008-12-25
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	130	<div> <div>6%</div> <div> <div></div> <div>64%</div> <div>27%</div> <div>• 8%</div> </div> </div>
1	B	130	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>18%</div> <div>• • 7%</div> </div> </div>
1	C	130	<div> <div>2%</div> <div> <div></div> <div>68%</div> <div>24%</div> <div>• 8%</div> </div> </div>
1	D	130	<div> <div>6%</div> <div> <div></div> <div>68%</div> <div>23%</div> <div>• 6%</div> </div> </div>
1	E	130	<div> <div>2%</div> <div> <div></div> <div>68%</div> <div>21%</div> <div>• 8%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	130	<div><div></div><div>6%</div><div>65%</div><div>23%</div><div>•</div><div>8%</div></div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5614 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called protein lp_1913.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	120	Total	C	N	O	Se	0	0	0
			919	592	149	174	4			
1	B	121	Total	C	N	O	Se	0	0	0
			924	595	150	175	4			
1	C	120	Total	C	N	O	Se	0	0	0
			919	592	149	174	4			
1	D	122	Total	C	N	O	Se	0	0	0
			932	600	151	178	3			
1	E	120	Total	C	N	O	Se	0	0	0
			919	592	149	174	4			
1	F	120	Total	C	N	O	Se	0	0	0
			919	592	149	174	4			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	LEU	-	expression tag	UNP Q88VW8
A	124	GLU	-	expression tag	UNP Q88VW8
A	125	HIS	-	expression tag	UNP Q88VW8
A	126	HIS	-	expression tag	UNP Q88VW8
A	127	HIS	-	expression tag	UNP Q88VW8
A	128	HIS	-	expression tag	UNP Q88VW8
A	129	HIS	-	expression tag	UNP Q88VW8
A	130	HIS	-	expression tag	UNP Q88VW8
B	123	LEU	-	expression tag	UNP Q88VW8
B	124	GLU	-	expression tag	UNP Q88VW8
B	125	HIS	-	expression tag	UNP Q88VW8
B	126	HIS	-	expression tag	UNP Q88VW8
B	127	HIS	-	expression tag	UNP Q88VW8
B	128	HIS	-	expression tag	UNP Q88VW8
B	129	HIS	-	expression tag	UNP Q88VW8
B	130	HIS	-	expression tag	UNP Q88VW8
C	123	LEU	-	expression tag	UNP Q88VW8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	124	GLU	-	expression tag	UNP Q88VW8
C	125	HIS	-	expression tag	UNP Q88VW8
C	126	HIS	-	expression tag	UNP Q88VW8
C	127	HIS	-	expression tag	UNP Q88VW8
C	128	HIS	-	expression tag	UNP Q88VW8
C	129	HIS	-	expression tag	UNP Q88VW8
C	130	HIS	-	expression tag	UNP Q88VW8
D	123	LEU	-	expression tag	UNP Q88VW8
D	124	GLU	-	expression tag	UNP Q88VW8
D	125	HIS	-	expression tag	UNP Q88VW8
D	126	HIS	-	expression tag	UNP Q88VW8
D	127	HIS	-	expression tag	UNP Q88VW8
D	128	HIS	-	expression tag	UNP Q88VW8
D	129	HIS	-	expression tag	UNP Q88VW8
D	130	HIS	-	expression tag	UNP Q88VW8
E	123	LEU	-	expression tag	UNP Q88VW8
E	124	GLU	-	expression tag	UNP Q88VW8
E	125	HIS	-	expression tag	UNP Q88VW8
E	126	HIS	-	expression tag	UNP Q88VW8
E	127	HIS	-	expression tag	UNP Q88VW8
E	128	HIS	-	expression tag	UNP Q88VW8
E	129	HIS	-	expression tag	UNP Q88VW8
E	130	HIS	-	expression tag	UNP Q88VW8
F	123	LEU	-	expression tag	UNP Q88VW8
F	124	GLU	-	expression tag	UNP Q88VW8
F	125	HIS	-	expression tag	UNP Q88VW8
F	126	HIS	-	expression tag	UNP Q88VW8
F	127	HIS	-	expression tag	UNP Q88VW8
F	128	HIS	-	expression tag	UNP Q88VW8
F	129	HIS	-	expression tag	UNP Q88VW8
F	130	HIS	-	expression tag	UNP Q88VW8

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	9	Total O 9 9	0	0
2	B	17	Total O 17 17	0	0
2	C	12	Total O 12 12	0	0
2	D	12	Total O 12 12	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	18	Total	O	0	0
			18	18		
2	F	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: protein lp_1913




• Molecule 1: protein lp_1913




• Molecule 1: protein lp_1913




• Molecule 1: protein lp_1913






● Molecule 1: protein lp_1913



● Molecule 1: protein lp_1913



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	119.36 Å 119.36 Å 67.79 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.92 – 2.70 29.84 – 2.69	Depositor EDS
% Data completeness (in resolution range)	65.3 (19.92-2.70) 87.7 (29.84-2.69)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.09 (at 2.68 Å)	Xtriage
Refinement program	CNS 1.2 & XtalView	Depositor
R, R_{free}	0.231 , 0.276 0.255 , 0.297	Depositor DCC
R_{free} test set	1295 reflections (6.38%)	DCC
Wilson B-factor (Å ²)	44.7	Xtriage
Anisotropy	0.188	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.2	EDS
Estimated twinning fraction	0.053 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 51430 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5614	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 82.55 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.7119e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/932	0.56	0/1261
1	B	0.35	0/937	0.56	0/1268
1	C	0.38	0/932	0.62	0/1261
1	D	0.39	0/945	0.57	0/1280
1	E	0.38	0/932	0.61	0/1261
1	F	0.35	0/932	0.58	0/1261
All	All	0.36	0/5610	0.58	0/7592

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	919	0	947	33	0
1	B	924	0	952	27	0
1	C	919	0	947	21	0
1	D	932	0	955	27	0
1	E	919	0	947	21	0
1	F	919	0	947	24	0
2	A	9	0	0	1	0
2	B	17	0	0	1	0
2	C	12	0	0	0	0
2	D	12	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	18	0	0	1	0
2	F	14	0	0	0	0
All	All	5614	0	5695	144	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:26:MSE:HE1	1:D:51:LYS:HB3	1.47	0.94
1:A:40:ASN:HD22	1:A:57:PRO:HB3	1.45	0.80
1:B:1:MSE:HB2	1:D:70:PRO:HG2	1.62	0.79
1:A:46:LYS:O	1:A:47:LYS:HG2	1.82	0.79
1:F:56:MSE:HG2	1:F:64:ARG:HH11	1.44	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/130 (91%)	108 (92%)	8 (7%)	2 (2%)	11	29
1	B	119/130 (92%)	109 (92%)	9 (8%)	1 (1%)	24	51
1	C	118/130 (91%)	109 (92%)	7 (6%)	2 (2%)	11	29
1	D	120/130 (92%)	109 (91%)	11 (9%)	0	100	100
1	E	118/130 (91%)	112 (95%)	5 (4%)	1 (1%)	24	51
1	F	118/130 (91%)	106 (90%)	10 (8%)	2 (2%)	11	29
All	All	711/780 (91%)	653 (92%)	50 (7%)	8 (1%)	17	42

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	47	LYS
1	F	43	ALA
1	E	47	LYS
1	B	47	LYS
1	F	49	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/101 (95%)	95 (99%)	1 (1%)	82	94
1	B	96/101 (95%)	89 (93%)	7 (7%)	17	39
1	C	96/101 (95%)	93 (97%)	3 (3%)	47	78
1	D	97/101 (96%)	92 (95%)	5 (5%)	29	58
1	E	96/101 (95%)	89 (93%)	7 (7%)	17	39
1	F	96/101 (95%)	90 (94%)	6 (6%)	22	48
All	All	577/606 (95%)	548 (95%)	29 (5%)	30	60

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	44	GLN
1	E	1	MSE
1	F	63	THR
1	D	65	ILE
1	E	3	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	28	ASN
1	C	40	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	27	GLN
1	B	49	GLN
1	C	114	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/130 (89%)	0.35	8 (6%) 20 18	25, 57, 80, 93	0
1	B	117/130 (90%)	0.15	4 (3%) 49 49	28, 54, 84, 105	0
1	C	116/130 (89%)	-0.04	2 (1%) 73 74	23, 44, 72, 84	0
1	D	119/130 (91%)	0.21	8 (6%) 21 19	31, 53, 86, 107	0
1	E	116/130 (89%)	-0.01	3 (2%) 59 59	25, 42, 65, 74	0
1	F	116/130 (89%)	0.49	8 (6%) 20 18	32, 57, 78, 85	0
All	All	700/780 (89%)	0.19	33 (4%) 35 34	23, 51, 81, 107	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	120	LEU	5.0
1	F	120	LEU	4.1
1	B	27	GLN	4.1
1	D	44	GLN	4.0
1	D	20	HIS	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.