



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:42 AM GMT

PDB ID : 3FPX  
Title : Native fungus laccase from Trametes hirsuta  
Authors : Polyakov, K.M.; Fedorova, T.V.; Stepanova, E.V.; Cherkashin, E.A.; Kurzeev, S.A.; Strokopytov, B.V.; Lamzin, V.S.; Koroleva, O.V.  
Deposited on : 2009-01-06  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	499	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	1504	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MAN	A	1513	-	-	-	X
4	NAG	A	1507	-	-	-	X
4	MAN	A	1521	-	-	-	X
5	NAG	A	1509	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4254 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Laccase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	5	0
			3806	2415	653	729	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	VAL	PRO	SEE REMARK 999	UNP B2L9C1

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Cu	0	0
			4	4		

- Molecule 3 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	6	Total	C	N	O	0	0
			72	40	2	30		

- Molecule 4 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 5 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	275	Total	O	0	0
			275	275		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Laccase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.29 Å 76.34 Å 129.66 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.90 – 1.80 19.97 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.5 (65.90-1.80) 94.7 (19.97-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 1.80 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.180 , 0.233 0.186 , 0.235	Depositor DCC
$R_{free}$ test set	2330 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtriage
Anisotropy	0.534	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 57.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	1 of 46097 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4254	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, BMA, NAG, CU, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.94	3/3950 (0.1%)	0.87	7/5424 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	351	VAL	CB-CG2	7.42	1.68	1.52
1	A	117	CYS	CB-SG	-5.80	1.72	1.81
1	A	99	VAL	CB-CG2	5.67	1.64	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	199	ARG	NE-CZ-NH1	6.88	123.74	120.30
1	A	424	ASP	CB-CG-OD2	-5.75	113.12	118.30
1	A	424	ASP	CB-CG-OD1	5.68	123.41	118.30
1	A	450	PHE	CB-CG-CD1	5.49	124.64	120.80
1	A	176	ARG	NE-CZ-NH1	5.47	123.04	120.30
1	A	486	ASP	CB-CG-OD1	5.13	122.91	118.30
1	A	450	PHE	CB-CG-CD2	-5.05	117.26	120.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	228	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3806	0	3593	37	0
2	A	4	0	0	0	0
3	A	72	0	61	0	0
4	A	50	0	43	0	0
5	A	14	0	13	0	0
6	A	28	0	25	1	0
7	A	5	0	0	0	0
8	A	275	0	0	1	0
All	All	4254	0	3735	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:ASP:OD1	1:A:182:VAL:HG22	1.75	0.87
1:A:192:LYS:CD	1:A:254:VAL:HG23	2.13	0.78
1:A:192:LYS:HD2	1:A:254:VAL:HG23	1.73	0.71
1:A:192:LYS:HD3	1:A:254:VAL:HG23	1.78	0.65
1:A:176:ARG:HD3	1:A:181:SER:HA	1.80	0.64
1:A:176:ARG:HG3	1:A:185:LEU:HD21	1.81	0.61
1:A:31:THR:HA	1:A:32:PRO:C	2.25	0.56
1:A:191:THR:HB	1:A:194:LYS:HD2	1.87	0.55
1:A:377:ASN:HD22	1:A:444:ASN:HD21	1.55	0.53
1:A:161:ARG:NH1	1:A:335:SER:O	2.42	0.53
1:A:116:TYR:CE1	1:A:205[A]:CYS:SG	3.02	0.53
1:A:10:ILE:HG22	1:A:52:LEU:HD21	1.92	0.52
1:A:188:ILE:N	1:A:188:ILE:HD12	2.25	0.52
1:A:64:HIS:CD2	8:A:1001:HOH:O	2.64	0.51
1:A:121:ARG:HD2	1:A:203:LEU:HB3	1.93	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:377:ASN:HA	1:A:444:ASN:HD21	1.76	0.50
1:A:147:THR:HG22	1:A:201[A]:VAL:CG1	2.41	0.50
1:A:405:ALA:HB2	1:A:422:PHE:CZ	2.48	0.49
1:A:301:ASN:OD1	1:A:303:VAL:HG22	2.13	0.49
1:A:192:LYS:HA	1:A:250:ALA:O	2.13	0.48
1:A:471:THR:N	1:A:472:PRO:CD	2.78	0.47
1:A:206:ASN:HB3	1:A:207:PRO:CD	2.45	0.47
1:A:419:ASN:N	1:A:420:PRO:CD	2.79	0.45
1:A:66:HIS:C	1:A:66:HIS:CD2	2.90	0.44
1:A:222:GLU:HB3	1:A:245:SER:HB2	2.00	0.44
1:A:301:ASN:CG	1:A:303:VAL:HG22	2.39	0.43
1:A:316:SER:N	1:A:321:GLY:O	2.51	0.43
1:A:133:HIS:ND1	1:A:136[B]:ARG:NH1	2.67	0.42
1:A:231:LEU:HD23	1:A:300:LEU:HD13	2.02	0.42
1:A:192:LYS:HE3	1:A:252:GLN:O	2.20	0.42
1:A:133:HIS:CD2	1:A:221:ILE:HB	2.55	0.42
1:A:136[B]:ARG:NH2	1:A:230:PRO:HG3	2.35	0.41
1:A:471:THR:HA	1:A:474:VAL:HG23	2.02	0.41
6:A:1511:NAG:H2	6:A:1511:NAG:H82	1.79	0.41
1:A:337:PHE:HB3	1:A:463:PHE:CD1	2.56	0.41
1:A:136[B]:ARG:HH21	1:A:230:PRO:HG3	1.86	0.40
1:A:403:THR:HG22	1:A:422:PHE:HB3	2.04	0.40
1:A:463:PHE:HD1	1:A:463:PHE:HA	1.82	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	502/499 (101%)	484 (96%)	15 (3%)	3 (1%)	30 14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	317	PRO
1	A	206	ASN
1	A	58	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	418/414 (101%)	409 (98%)	9 (2%)	60 45

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	TYR
1	A	124	PHE
1	A	205[A]	CYS
1	A	205[B]	CYS
1	A	243	ARG
1	A	318	SER
1	A	387	THR
1	A	450	PHE
1	A	463	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	267	ASN
1	A	360	GLN
1	A	444	ASN
1	A	499	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

12 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAG	A	1504	1,3	14,14,15	0.87	1 (7%)	15,19,21	0.95	1 (6%)
3	NAG	A	1505	3	14,14,15	0.60	0	15,19,21	2.22	4 (26%)
4	NAG	A	1506	1,4	14,14,15	0.77	0	15,19,21	1.57	1 (6%)
4	NAG	A	1507	4	14,14,15	0.79	1 (7%)	15,19,21	2.15	7 (46%)
4	BMA	A	1508	4	11,11,12	0.45	0	14,15,17	0.96	0
6	NAG	A	1510	1,6	14,14,15	0.66	0	15,19,21	1.39	3 (20%)
6	NAG	A	1511	6	14,14,15	0.52	0	15,19,21	2.47	5 (33%)
3	BMA	A	1512	3	11,11,12	0.73	0	14,15,17	1.97	5 (35%)
3	MAN	A	1513	3	11,11,12	0.61	0	14,15,17	1.72	4 (28%)
3	MAN	A	1514	3	11,11,12	0.69	0	14,15,17	1.32	2 (14%)
3	MAN	A	1515	3	11,11,12	0.83	0	14,15,17	3.19	5 (35%)
4	MAN	A	1521	4	11,11,12	0.62	0	14,15,17	1.32	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1504	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1505	3	-	0/6/23/26	0/1/1/1
4	NAG	A	1506	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	1507	4	-	0/6/23/26	0/1/1/1
4	BMA	A	1508	4	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	1510	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	1511	6	-	0/6/23/26	0/1/1/1
3	BMA	A	1512	3	-	0/2/19/22	0/1/1/1
3	MAN	A	1513	3	-	0/2/19/22	0/1/1/1
3	MAN	A	1514	3	-	0/2/19/22	0/1/1/1
3	MAN	A	1515	3	-	0/2/19/22	0/1/1/1
4	MAN	A	1521	4	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1507	NAG	O5-C1	-2.29	1.39	1.43
3	A	1504	NAG	C1-C2	2.37	1.55	1.52

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1505	NAG	C1-O5-C5	-5.71	105.00	112.25
3	A	1512	BMA	C1-C2-C3	-4.15	104.63	109.54
6	A	1511	NAG	C2-N2-C7	-4.06	117.82	123.04
4	A	1507	NAG	O6-C6-C5	-3.54	99.64	111.33
6	A	1511	NAG	C4-C3-C2	-3.45	105.86	111.23
6	A	1510	NAG	C3-C2-N2	-3.45	102.30	110.56
4	A	1507	NAG	C1-O5-C5	-3.33	108.02	112.25
4	A	1507	NAG	O7-C7-C8	-3.29	116.02	122.06
3	A	1505	NAG	O7-C7-C8	-3.25	116.09	122.06
3	A	1512	BMA	O4-C4-C3	-2.92	103.75	110.34
3	A	1513	MAN	O4-C4-C3	-2.76	104.12	110.34
4	A	1507	NAG	C4-C3-C2	-2.51	107.33	111.23
4	A	1507	NAG	O4-C4-C3	-2.33	105.09	110.34
3	A	1504	NAG	C2-N2-C7	-2.17	120.25	123.04
6	A	1511	NAG	C6-C5-C4	-2.09	107.87	113.02
3	A	1514	MAN	O4-C4-C3	-2.04	105.75	110.34
3	A	1512	BMA	O2-C2-C1	2.05	113.31	109.21
4	A	1521	MAN	O3-C3-C2	2.12	113.82	110.00
4	A	1507	NAG	O4-C4-C5	2.13	114.88	109.24
3	A	1513	MAN	C6-C5-C4	2.19	118.41	113.02
3	A	1512	BMA	O5-C5-C6	2.22	112.15	107.35
6	A	1510	NAG	O4-C4-C5	2.23	115.15	109.24
3	A	1514	MAN	C1-C2-C3	2.28	112.24	109.54
6	A	1511	NAG	C3-C2-N2	2.32	116.12	110.56
6	A	1510	NAG	C1-O5-C5	2.36	115.25	112.25

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1513	MAN	O2-C2-C1	2.39	113.99	109.21
4	A	1521	MAN	C1-O5-C5	2.61	115.56	112.25
4	A	1521	MAN	O5-C5-C6	2.63	113.04	107.35
3	A	1512	BMA	O5-C1-C2	2.64	115.14	110.86
3	A	1515	MAN	C2-C3-C4	2.83	115.85	111.04
3	A	1505	NAG	C4-C3-C2	2.85	115.66	111.23
3	A	1505	NAG	O5-C5-C6	3.08	114.02	107.35
4	A	1507	NAG	C8-C7-N2	3.38	122.57	116.11
3	A	1515	MAN	C3-C4-C5	3.49	116.28	110.20
3	A	1515	MAN	O5-C1-C2	4.07	117.46	110.86
3	A	1513	MAN	C1-O5-C5	4.23	117.62	112.25
4	A	1506	NAG	C1-O5-C5	4.90	118.47	112.25
6	A	1511	NAG	C1-O5-C5	6.51	120.51	112.25
3	A	1515	MAN	C1-C2-C3	6.67	117.43	109.54
3	A	1515	MAN	C1-O5-C5	7.26	121.47	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1511	NAG	1	0

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	A	1509	1	14,14,15	0.74	1 (7%)	15,19,21	1.65	2 (13%)
7	PO4	A	500	-	4,4,4	0.78	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1509	1	-	0/6/23/26	0/1/1/1
7	PO4	A	500	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1509	NAG	C1-C2	2.13	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1509	NAG	C3-C4-C5	-4.32	102.67	110.20
5	A	1509	NAG	O5-C5-C6	3.27	114.43	107.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	499/499 (100%)	0.05	20 (4%) 42 36	20, 29, 48, 62	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	320	GLY	9.1
1	A	318	SER	7.9
1	A	319	SER	5.9
1	A	290	THR	3.7
1	A	287	VAL	3.6
1	A	317	PRO	3.5
1	A	497	ASN	3.4
1	A	321	GLY	3.3
1	A	182	VAL	2.9
1	A	55	HIS	2.8
1	A	286	ALA	2.6
1	A	135	SER	2.6
1	A	295	THR	2.6
1	A	285	PRO	2.5
1	A	316	SER	2.4
1	A	439	ILE	2.3
1	A	377	ASN	2.3
1	A	130	ASN	2.2
1	A	178	PRO	2.2
1	A	312	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MAN	A	1521	11/12	0.86	0.21	4.95	54,56,58,59	0
4	NAG	A	1507	14/15	0.93	0.19	2.52	36,41,48,52	0
3	NAG	A	1504	14/15	0.95	0.11	2.40	26,35,38,41	0
3	MAN	A	1513	11/12	0.91	0.18	2.18	32,38,45,48	0
3	MAN	A	1514	11/12	0.91	0.11	1.06	33,40,44,45	0
3	NAG	A	1505	14/15	0.94	0.14	1.05	28,35,42,45	0
6	NAG	A	1510	14/15	0.79	0.27	1.05	49,54,56,57	0
4	NAG	A	1506	14/15	0.90	0.12	0.07	38,40,45,51	0
6	NAG	A	1511	14/15	0.64	0.43	-	59,63,65,65	0
3	MAN	A	1515	11/12	0.82	0.27	-	45,53,59,63	0
4	BMA	A	1508	11/12	0.84	0.25	-	41,48,50,52	0
3	BMA	A	1512	11/12	0.90	0.18	-	37,42,47,50	0

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	1509	14/15	0.92	0.21	3.48	35,40,42,42	0
2	CU	A	1501	1/1	0.99	0.06	-1.96	31,31,31,31	1
2	CU	A	1503	1/1	1.00	0.04	-3.00	27,27,27,27	1
2	CU	A	1500	1/1	0.99	0.05	-6.18	24,24,24,24	1
7	PO4	A	500	5/5	0.96	0.13	-	32,34,35,35	5
2	CU	A	1502	1/1	1.00	0.03	-	26,26,26,26	1

### 6.5 Other polymers ⓘ

There are no such residues in this entry.