



wwPDB X-ray Structure Validation Summary Report ⓘ

May 5, 2016 – 05:12 PM EDT

PDB ID : 5FSW
Title : RNA dependent RNA polymerase QDE-1 from Thielavia terrestris
Authors : Qian, X.; Hamid, F.M.; El Sahili, A.; Darwis, D.A.; Wong, Y.H.; Bhushan, S.;
Makeyev, E.V.; Lescar, J.
Deposited on : 2016-01-08
Resolution : 3.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027457
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1034	<div> <div>4%</div> <div> <div>47%</div> <div>34%</div> <div>8%</div> <div>11%</div> </div> </div>
1	B	1034	<div> <div>10%</div> <div> <div>58%</div> <div>27%</div> <div>1%</div> <div>11%</div> </div> </div>
1	C	1034	<div> <div>18%</div> <div> <div>62%</div> <div>22%</div> <div>5%</div> <div>11%</div> </div> </div>
1	D	1034	<div> <div>22%</div> <div> <div>60%</div> <div>25%</div> <div>1%</div> <div>11%</div> </div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 28130 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA DEPENDENT RNA POLYMERASE QDE-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	924	Total	C	N	O	S	0	0	0
			7358	4679	1307	1335	37			
1	B	922	Total	C	N	O	S	0	0	0
			6924	4462	1186	1244	32			
1	C	922	Total	C	N	O	S	0	0	0
			6924	4462	1186	1244	32			
1	D	922	Total	C	N	O	S	0	0	0
			6924	4462	1186	1244	32			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP G2R911
A	2	GLY	-	EXPRESSION TAG	UNP G2R911
A	3	HIS	-	EXPRESSION TAG	UNP G2R911
A	4	HIS	-	EXPRESSION TAG	UNP G2R911
A	5	HIS	-	EXPRESSION TAG	UNP G2R911
A	6	HIS	-	EXPRESSION TAG	UNP G2R911
A	7	HIS	-	EXPRESSION TAG	UNP G2R911
A	8	HIS	-	EXPRESSION TAG	UNP G2R911
A	9	SER	-	EXPRESSION TAG	UNP G2R911
A	10	SER	-	EXPRESSION TAG	UNP G2R911
A	11	GLY	-	EXPRESSION TAG	UNP G2R911
A	12	VAL	-	EXPRESSION TAG	UNP G2R911
A	13	ASP	-	EXPRESSION TAG	UNP G2R911
A	14	LEU	-	EXPRESSION TAG	UNP G2R911
A	15	GLY	-	EXPRESSION TAG	UNP G2R911
A	16	THR	-	EXPRESSION TAG	UNP G2R911
A	17	GLU	-	EXPRESSION TAG	UNP G2R911
A	18	ASN	-	EXPRESSION TAG	UNP G2R911
A	19	LEU	-	EXPRESSION TAG	UNP G2R911
A	20	TYR	-	EXPRESSION TAG	UNP G2R911
A	21	PHE	-	EXPRESSION TAG	UNP G2R911

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLN	-	EXPRESSION TAG	UNP G2R911
A	23	SER	-	EXPRESSION TAG	UNP G2R911
A	24	MET	-	EXPRESSION TAG	UNP G2R911
B	1	MET	-	EXPRESSION TAG	UNP G2R911
B	2	GLY	-	EXPRESSION TAG	UNP G2R911
B	3	HIS	-	EXPRESSION TAG	UNP G2R911
B	4	HIS	-	EXPRESSION TAG	UNP G2R911
B	5	HIS	-	EXPRESSION TAG	UNP G2R911
B	6	HIS	-	EXPRESSION TAG	UNP G2R911
B	7	HIS	-	EXPRESSION TAG	UNP G2R911
B	8	HIS	-	EXPRESSION TAG	UNP G2R911
B	9	SER	-	EXPRESSION TAG	UNP G2R911
B	10	SER	-	EXPRESSION TAG	UNP G2R911
B	11	GLY	-	EXPRESSION TAG	UNP G2R911
B	12	VAL	-	EXPRESSION TAG	UNP G2R911
B	13	ASP	-	EXPRESSION TAG	UNP G2R911
B	14	LEU	-	EXPRESSION TAG	UNP G2R911
B	15	GLY	-	EXPRESSION TAG	UNP G2R911
B	16	THR	-	EXPRESSION TAG	UNP G2R911
B	17	GLU	-	EXPRESSION TAG	UNP G2R911
B	18	ASN	-	EXPRESSION TAG	UNP G2R911
B	19	LEU	-	EXPRESSION TAG	UNP G2R911
B	20	TYR	-	EXPRESSION TAG	UNP G2R911
B	21	PHE	-	EXPRESSION TAG	UNP G2R911
B	22	GLN	-	EXPRESSION TAG	UNP G2R911
B	23	SER	-	EXPRESSION TAG	UNP G2R911
B	24	MET	-	EXPRESSION TAG	UNP G2R911
C	1	MET	-	EXPRESSION TAG	UNP G2R911
C	2	GLY	-	EXPRESSION TAG	UNP G2R911
C	3	HIS	-	EXPRESSION TAG	UNP G2R911
C	4	HIS	-	EXPRESSION TAG	UNP G2R911
C	5	HIS	-	EXPRESSION TAG	UNP G2R911
C	6	HIS	-	EXPRESSION TAG	UNP G2R911
C	7	HIS	-	EXPRESSION TAG	UNP G2R911
C	8	HIS	-	EXPRESSION TAG	UNP G2R911
C	9	SER	-	EXPRESSION TAG	UNP G2R911
C	10	SER	-	EXPRESSION TAG	UNP G2R911
C	11	GLY	-	EXPRESSION TAG	UNP G2R911
C	12	VAL	-	EXPRESSION TAG	UNP G2R911
C	13	ASP	-	EXPRESSION TAG	UNP G2R911
C	14	LEU	-	EXPRESSION TAG	UNP G2R911
C	15	GLY	-	EXPRESSION TAG	UNP G2R911

Continued on next page...


Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	16	THR	-	EXPRESSION TAG	UNP G2R911
C	17	GLU	-	EXPRESSION TAG	UNP G2R911
C	18	ASN	-	EXPRESSION TAG	UNP G2R911
C	19	LEU	-	EXPRESSION TAG	UNP G2R911
C	20	TYR	-	EXPRESSION TAG	UNP G2R911
C	21	PHE	-	EXPRESSION TAG	UNP G2R911
C	22	GLN	-	EXPRESSION TAG	UNP G2R911
C	23	SER	-	EXPRESSION TAG	UNP G2R911
C	24	MET	-	EXPRESSION TAG	UNP G2R911
D	1	MET	-	EXPRESSION TAG	UNP G2R911
D	2	GLY	-	EXPRESSION TAG	UNP G2R911
D	3	HIS	-	EXPRESSION TAG	UNP G2R911
D	4	HIS	-	EXPRESSION TAG	UNP G2R911
D	5	HIS	-	EXPRESSION TAG	UNP G2R911
D	6	HIS	-	EXPRESSION TAG	UNP G2R911
D	7	HIS	-	EXPRESSION TAG	UNP G2R911
D	8	HIS	-	EXPRESSION TAG	UNP G2R911
D	9	SER	-	EXPRESSION TAG	UNP G2R911
D	10	SER	-	EXPRESSION TAG	UNP G2R911
D	11	GLY	-	EXPRESSION TAG	UNP G2R911
D	12	VAL	-	EXPRESSION TAG	UNP G2R911
D	13	ASP	-	EXPRESSION TAG	UNP G2R911
D	14	LEU	-	EXPRESSION TAG	UNP G2R911
D	15	GLY	-	EXPRESSION TAG	UNP G2R911
D	16	THR	-	EXPRESSION TAG	UNP G2R911
D	17	GLU	-	EXPRESSION TAG	UNP G2R911
D	18	ASN	-	EXPRESSION TAG	UNP G2R911
D	19	LEU	-	EXPRESSION TAG	UNP G2R911
D	20	TYR	-	EXPRESSION TAG	UNP G2R911
D	21	PHE	-	EXPRESSION TAG	UNP G2R911
D	22	GLN	-	EXPRESSION TAG	UNP G2R911
D	23	SER	-	EXPRESSION TAG	UNP G2R911
D	24	MET	-	EXPRESSION TAG	UNP G2R911





- Molecule 1: RNA DEPENDENT RNA POLYMERASE QDE-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.23Å 165.95Å 173.89Å 90.00° 90.10° 90.00°	Depositor
Resolution (Å)	48.91 – 3.19 48.91 – 3.19	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.91-3.19) 99.7 (48.91-3.19)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	22.98 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.213 , 0.251 0.209 , 0.246	Depositor DCC
R_{free} test set	3829 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.396	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 78.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.005 for -h,l,k 0.009 for -h,-l,-k 0.052 for h,-k,-l	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	28130	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.92	8/7528 (0.1%)	1.07	25/10177 (0.2%)
1	B	0.82	6/7063 (0.1%)	0.97	20/9494 (0.2%)
1	C	0.56	0/7063	0.85	6/9494 (0.1%)
1	D	0.56	0/7062	0.82	5/9491 (0.1%)
All	All	0.74	14/28716 (0.0%)	0.93	56/38656 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8
1	B	0	1
1	D	0	1
All	All	0	10

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	308	GLU	CD-OE1	6.83	1.33	1.25
1	B	308	GLU	CG-CD	6.73	1.62	1.51
1	A	284	TYR	CD1-CE1	-6.49	1.29	1.39
1	A	401	GLU	CD-OE1	6.43	1.32	1.25
1	A	349	SER	CB-OG	6.19	1.50	1.42

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	632	ASP	CB-CG-OD2	-13.18	106.44	118.30
1	A	425	ARG	NE-CZ-NH1	11.53	126.06	120.30
1	B	632	ASP	CB-CG-OD1	11.51	128.66	118.30
1	B	333	ARG	NE-CZ-NH1	8.87	124.73	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	650	ASP	CB-CG-OD1	8.67	126.10	118.30

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	132	SER	Peptide
1	A	174	ASN	Peptide
1	A	322	PRO	Peptide
1	A	68	LEU	Peptide
1	A	76	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7358	0	7279	444	0
1	B	6924	0	6423	216	0
1	C	6924	0	6423	181	0
1	D	6924	0	6422	199	0
All	All	28130	0	26547	981	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 981 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:321:SER:O	1:A:323:ALA:HA	1.24	1.30
1:A:943:LYS:HB2	1:A:944:THR:OG1	1.16	1.25
1:A:321:SER:O	1:A:323:ALA:CA	1.86	1.23
1:A:133:LYS:CA	1:A:135:GLY:H	1.56	1.17
1:A:355:ILE:CD1	1:A:405:VAL:HG12	1.75	1.16

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	904/1034 (87%)	779 (86%)	90 (10%)	35 (4%)	4	28
1	B	888/1034 (86%)	779 (88%)	84 (10%)	25 (3%)	6	37
1	C	888/1034 (86%)	786 (88%)	75 (8%)	27 (3%)	5	35
1	D	886/1034 (86%)	787 (89%)	71 (8%)	28 (3%)	5	33
All	All	3566/4136 (86%)	3131 (88%)	320 (9%)	115 (3%)	5	33

5 of 115 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	PRO
1	A	88	GLN
1	A	133	LYS
1	A	137	LEU
1	A	184	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	775/856 (90%)	695 (90%)	80 (10%)	9	36
1	B	655/856 (76%)	585 (89%)	70 (11%)	8	34
1	C	655/856 (76%)	577 (88%)	78 (12%)	6	28
1	D	655/856 (76%)	578 (88%)	77 (12%)	6	29
All	All	2740/3424 (80%)	2435 (89%)	305 (11%)	8	32

5 of 305 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	808	MET
1	C	316	THR
1	D	728	LEU
1	B	877	THR
1	C	84	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	310	ASN
1	B	555	HIS
1	D	463	GLN
1	B	463	GLN
1	B	963	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	924/1034 (89%)	-0.00	42 (4%) 37 23	11, 21, 94, 177	0
1	B	922/1034 (89%)	0.34	100 (10%) 8 4	9, 25, 146, 228	0
1	C	922/1034 (89%)	1.16	185 (20%) 1 1	25, 71, 152, 347	0
1	D	922/1034 (89%)	1.55	229 (24%) 1 1	24, 71, 221, 430	0
All	All	3690/4136 (89%)	0.76	556 (15%) 3 2	9, 53, 158, 430	0

The worst 5 of 556 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	823	HIS	21.6
1	D	936	THR	18.8
1	D	955	ALA	18.5
1	D	940	ALA	16.8
1	D	935	SER	16.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.