



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:09 AM GMT

PDB ID : 2FT3  
Title : Crystal structure of the biglycan dimer core protein  
Authors : Scott, P.G.; Dodd, C.M.; Bergmann, E.M.  
Deposited on : 2006-01-23  
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1476 (3.50-3.30)
Clashscore	102246	1611 (3.50-3.30)
Ramachandran outliers	100387	1571 (3.50-3.30)
Sidechain outliers	100360	1571 (3.50-3.30)
RSRZ outliers	91569	1485 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	
1	B	332	
1	C	332	
1	D	332	
1	E	332	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	332	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	D	2193	-	-	X	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 14674 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Biglycan.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2416	1545	428	432	11			
1	B	303	Total	C	N	O	S	0	0	0
			2416	1545	428	432	11			
1	C	303	Total	C	N	O	S	0	0	0
			2416	1545	428	432	11			
1	D	304	Total	C	N	O	S	0	0	0
			2420	1547	429	433	11			
1	E	303	Total	C	N	O	S	0	0	0
			2416	1545	428	432	11			
1	F	305	Total	C	N	O	S	0	0	0
			2428	1551	431	435	11			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	E	1	Total	C	N	O	0	0
			14	8	1	5		
2	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula:  $C_6H_5O_7$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		
3	C	1	Total	C	O	0	0
			13	6	7		
3	D	1	Total	C	O	0	0
			13	6	7		
3	E	1	Total	C	O	0	0
			13	6	7		
3	F	1	Total	C	O	0	0
			13	6	7		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Biglycan




#### • Molecule 1: Biglycan




#### • Molecule 1: Biglycan





• Molecule 1: Biglycan





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	206.51Å 119.22Å 140.62Å 90.00° 116.61° 90.00°	Depositor
Resolution (Å)	23.00 – 3.40 22.86 – 3.40	Depositor EDS
% Data completeness (in resolution range)	96.5 (23.00-3.40) 96.5 (22.86-3.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 3.37Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.258 , 0.291 0.255 , 0.294	Depositor DCC
$R_{free}$ test set	2040 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	47.1	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 29.3	EDS
Estimated twinning fraction	0.098 for 1/2*h-3/2*k,-1/2*h-1/2*k,-1/2*h +1/2*k-l 0.090 for 1/2*h+3/2*k,1/2*h-1/2*k,-1/2*h- 1/2*k-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.24$	Xtriage
Outliers	1 of 40497 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	14674	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.07	3/2468 (0.1%)	1.23	17/3342 (0.5%)
1	B	1.05	1/2468 (0.0%)	1.23	23/3342 (0.7%)
1	C	1.01	1/2468 (0.0%)	1.28	24/3342 (0.7%)
1	D	1.05	2/2472 (0.1%)	1.29	24/3347 (0.7%)
1	E	1.06	1/2468 (0.0%)	1.36	26/3342 (0.8%)
1	F	1.05	2/2480 (0.1%)	1.25	30/3358 (0.9%)
All	All	1.05	10/14824 (0.1%)	1.27	144/20073 (0.7%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	62	GLN	CD-NE2	-7.44	1.14	1.32
1	E	151	CYS	CB-SG	-6.55	1.71	1.82
1	F	62	GLN	CD-OE1	-6.31	1.10	1.24
1	A	108	TYR	CE1-CZ	-5.78	1.31	1.38
1	B	291	VAL	CA-CB	5.62	1.66	1.54

The worst 5 of 144 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	221	ARG	NE-CZ-NH2	-20.38	110.11	120.30
1	E	244	ARG	NE-CZ-NH2	18.19	129.40	120.30
1	D	317	ARG	NE-CZ-NH2	17.19	128.90	120.30
1	E	244	ARG	NE-CZ-NH1	-15.47	112.56	120.30
1	E	136	ARG	NE-CZ-NH2	-15.26	112.67	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2466	222	0
1	B	2416	0	2466	163	0
1	C	2416	0	2466	188	0
1	D	2420	0	2469	167	0
1	E	2416	0	2466	173	0
1	F	2428	0	2475	191	0
2	A	14	0	13	5	0
2	B	14	0	13	0	0
2	C	14	0	13	0	0
2	D	14	0	13	1	0
2	E	14	0	13	0	0
2	F	14	0	13	0	0
3	A	13	0	5	0	0
3	B	13	0	5	3	0
3	C	13	0	5	3	0
3	D	13	0	5	6	0
3	E	13	0	5	2	0
3	F	13	0	5	2	0
All	All	14674	0	14916	1104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 1104 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:GLU:OE1	1:B:178:ARG:HD3	1.39	1.21
1:D:225:GLY:HA3	3:D:2193:FLC:OB2	1.00	1.15
1:E:27:CYS:HB2	1:E:28:PRO:HD2	1.33	1.10
1:C:121:LEU:HD12	1:C:121:LEU:H	1.15	1.09
1:A:93:ILE:HD11	1:A:114:LEU:HD21	1.32	1.09

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/332 (91%)	216 (72%)	70 (23%)	15 (5%)	3	24
1	B	301/332 (91%)	225 (75%)	68 (23%)	8 (3%)	6	41
1	C	301/332 (91%)	219 (73%)	68 (23%)	14 (5%)	3	26
1	D	302/332 (91%)	230 (76%)	59 (20%)	13 (4%)	3	29
1	E	301/332 (91%)	229 (76%)	53 (18%)	19 (6%)	2	18
1	F	303/332 (91%)	223 (74%)	67 (22%)	13 (4%)	3	29
All	All	1809/1992 (91%)	1342 (74%)	385 (21%)	82 (4%)	3	27

5 of 82 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	26	MET
1	A	291	VAL
1	B	150	ASN
1	C	150	ASN
1	C	291	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/297 (92%)	244 (89%)	29 (11%)	8	36
1	B	273/297 (92%)	253 (93%)	20 (7%)	17	56
1	C	273/297 (92%)	249 (91%)	24 (9%)	12	46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	273/297 (92%)	243 (89%)	30 (11%)	8	34
1	E	273/297 (92%)	245 (90%)	28 (10%)	9	37
1	F	274/297 (92%)	246 (90%)	28 (10%)	9	38
All	All	1639/1782 (92%)	1480 (90%)	159 (10%)	10	40

5 of 159 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	306	VAL
1	D	191	LEU
1	F	191	LEU
1	D	61	LEU
1	D	142	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	62	GLN
1	D	175	ASN
1	F	175	ASN
1	D	105	GLN
1	D	206	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FLC	A	2190	-	3,12,12	0.48	0	3,17,17	3.63	1 (33%)
2	NAG	A	801	1	14,14,15	1.17	1 (7%)	15,19,21	1.63	3 (20%)
3	FLC	B	2191	-	3,12,12	0.90	0	3,17,17	0.80	0
2	NAG	B	803	1	14,14,15	1.02	1 (7%)	15,19,21	1.71	3 (20%)
3	FLC	C	2192	-	3,12,12	1.22	0	3,17,17	4.94	2 (66%)
2	NAG	C	805	1	14,14,15	1.27	2 (14%)	15,19,21	2.11	7 (46%)
3	FLC	D	2193	1	3,12,12	1.08	0	3,17,17	2.95	2 (66%)
2	NAG	D	807	1	14,14,15	0.92	0	15,19,21	1.71	4 (26%)
3	FLC	E	2194	-	3,12,12	1.27	0	3,17,17	3.92	2 (66%)
2	NAG	E	809	1	14,14,15	1.27	2 (14%)	15,19,21	2.16	3 (20%)
3	FLC	F	2195	-	3,12,12	2.12	2 (66%)	3,17,17	5.97	2 (66%)
2	NAG	F	811	1	14,14,15	1.05	1 (7%)	15,19,21	1.91	6 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FLC	A	2190	-	-	0/6/16/16	0/0/0/0
2	NAG	A	801	1	-	0/6/23/26	0/1/1/1
3	FLC	B	2191	-	-	0/6/16/16	0/0/0/0
2	NAG	B	803	1	-	0/6/23/26	0/1/1/1
3	FLC	C	2192	-	-	0/6/16/16	0/0/0/0
2	NAG	C	805	1	-	0/6/23/26	0/1/1/1
3	FLC	D	2193	1	-	0/6/16/16	0/0/0/0
2	NAG	D	807	1	-	0/6/23/26	0/1/1/1
3	FLC	E	2194	-	-	0/6/16/16	0/0/0/0
2	NAG	E	809	1	-	2/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FLC	F	2195	-	-	0/6/16/16	0/0/0/0
2	NAG	F	811	1	-	0/6/23/26	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	805	NAG	C1-C2	2.07	1.55	1.52
2	E	809	NAG	C4-C3	2.07	1.57	1.52
3	F	2195	FLC	OHB-CB	2.09	1.46	1.43
2	B	803	NAG	C1-C2	2.18	1.55	1.52
2	F	811	NAG	C1-C2	2.59	1.56	1.52

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2192	FLC	CB-CA-CAC	-7.21	103.42	114.96
3	E	2194	FLC	CB-CG-CGC	-6.17	105.09	114.96
3	A	2190	FLC	CB-CG-CGC	-6.05	105.28	114.96
3	C	2192	FLC	CB-CG-CGC	-4.58	107.63	114.96
3	D	2193	FLC	CB-CG-CGC	-3.75	108.96	114.96

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	809	NAG	O7-C7-N2-C2
2	E	809	NAG	C8-C7-N2-C2

There are no ring outliers.

7 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	NAG	5	0
3	B	2191	FLC	3	0
3	C	2192	FLC	3	0
3	D	2193	FLC	6	0
2	D	807	NAG	1	0
3	E	2194	FLC	2	0
3	F	2195	FLC	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	303/332 (91%)	0.22	0 <b>100</b> <b>100</b>	8, 38, 52, 57	0
1	B	303/332 (91%)	0.33	4 (1%) 79 74	8, 38, 53, 70	0
1	C	303/332 (91%)	0.26	3 (0%) 84 79	8, 38, 53, 67	0
1	D	304/332 (91%)	0.21	0 <b>100</b> <b>100</b>	8, 38, 52, 57	0
1	E	303/332 (91%)	0.24	2 (0%) 89 85	8, 37, 53, 64	0
1	F	305/332 (91%)	0.25	1 (0%) 94 92	6, 38, 52, 58	0
All	All	1821/1992 (91%)	0.25	10 (0%) <b>91</b> <b>89</b>	6, 38, 53, 70	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	290	GLY	3.7
1	B	288	GLY	3.2
1	B	25	ALA	3.1
1	E	25	ALA	2.8
1	E	294	ALA	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	FLC	B	2191	13/13	0.90	0.30	1.28	49,53,56,57	0
3	FLC	E	2194	13/13	0.92	0.21	-0.35	36,44,47,48	0
3	FLC	A	2190	13/13	0.94	0.20	-0.70	47,50,53,53	0
3	FLC	F	2195	13/13	0.87	0.21	-0.76	38,42,42,43	0
3	FLC	D	2193	13/13	0.95	0.17	-1.15	38,40,44,45	0
3	FLC	C	2192	13/13	0.94	0.18	-1.25	38,43,46,46	0
2	NAG	E	809	14/15	0.72	0.40	-	59,61,62,62	0
2	NAG	A	801	14/15	0.89	0.24	-	51,53,55,55	0
2	NAG	D	807	14/15	0.89	0.27	-	58,59,61,62	0
2	NAG	B	803	14/15	0.80	0.27	-	46,50,51,51	0
2	NAG	C	805	14/15	0.79	0.30	-	53,56,57,58	0
2	NAG	F	811	14/15	0.89	0.16	-	49,50,51,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.