



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 08:44 AM GMT

PDB ID : 3FUQ
Title : Glycosylated SV2 and Gangliosides as Dual Receptors for Botulinum Neurotoxin Serotype F
Authors : Fu, Z.; Chen, C.; Barbieri, J.T.; Kim, J.-J.P.; Baldwin, M.R.
Deposited on : 2009-01-14
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	417	3%	67%	26%	..

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3560 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BoNT/F (Neurotoxin type F).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	399	3285	2086	567	624	8	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	275	275	275	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BoNT/F (Neurotoxin type F)



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.33 Å 74.18 Å 74.77 Å 90.00° 106.36° 90.00°	Depositor
Resolution (Å)	29.17 – 2.10 29.17 – 2.10	Depositor EDS
% Data completeness (in resolution range)	91.4 (29.17-2.10) 91.2 (29.17-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle^1$	12.06 (at 2.10 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R , R_{free}	0.206 , 0.243 0.201 , 0.237	Depositor DCC
R_{free} test set	2350 reflections (9.88%)	DCC
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.9	EDS
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 23800 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3560	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/3352	0.67	1/4538 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1129	ARG	N-CA-C	-5.12	97.18	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3285	0	3200	101	0
2	A	275	0	0	12	0
All	All	3560	0	3200	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (\AA)	Clash overlap (\AA)
1:A:973:ASN:HD22	1:A:974:TYR:H	1.14	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:918:ASN:HD21	1:A:1046:ASN:HD22	1.17	0.93
1:A:1191:ILE:HD11	1:A:1197:ILE:HG22	1.58	0.86
1:A:945:TYR:H	1:A:997:GLN:HE22	1.26	0.79
1:A:973:ASN:HD22	1:A:974:TYR:N	1.83	0.76
1:A:941:ARG:HD2	1:A:1006:ASN:O	1.86	0.76
1:A:1062:ILE:HD12	1:A:1063:ARG:H	1.50	0.75
1:A:918:ASN:HD21	1:A:1046:ASN:ND2	1.85	0.73
1:A:1037:ASN:HD22	1:A:1037:ASN:H	1.35	0.73
1:A:1126:ASN:HD22	1:A:1126:ASN:H	1.37	0.73
1:A:949:VAL:HG23	2:A:502:HOH:O	1.88	0.73
1:A:1020:ASN:HD22	1:A:1034:SER:HA	1.55	0.71
1:A:893:ILE:HD11	1:A:897:VAL:HB	1.73	0.71
1:A:973:ASN:ND2	1:A:974:TYR:H	1.89	0.70
1:A:1171:LEU:HD22	1:A:1213:ILE:HD12	1.71	0.70
1:A:1191:ILE:HG22	1:A:1193:LYS:HG3	1.74	0.69
1:A:953:ASN:HD21	1:A:1055:ASN:H	1.40	0.68
1:A:984:THR:HG21	2:A:600:HOH:O	1.91	0.68
1:A:1020:ASN:ND2	1:A:1034:SER:HA	2.10	0.67
1:A:1022:ARG:HD3	1:A:1032:GLU:OE2	1.96	0.65
1:A:871:ILE:HD13	1:A:927:TYR:OH	1.97	0.65
1:A:1062:ILE:HD12	1:A:1063:ARG:N	2.12	0.63
1:A:1191:ILE:O	1:A:1192:ALA:HB3	1.98	0.62
1:A:939:TRP:HB2	1:A:1063:ARG:HG2	1.82	0.62
1:A:1008:TRP:HE1	1:A:1096:ASN:HD21	1.50	0.60
1:A:1008:TRP:HE1	1:A:1096:ASN:ND2	1.99	0.60
1:A:1141:THR:HG22	2:A:602:HOH:O	2.03	0.59
1:A:1229:ASN:HD22	1:A:1229:ASN:C	2.06	0.58
1:A:1191:ILE:HD12	1:A:1196:LYS:HA	1.86	0.57
1:A:930:ARG:HG2	2:A:645:HOH:O	2.03	0.57
1:A:1229:ASN:HD21	1:A:1233:GLY:H	1.51	0.57
1:A:953:ASN:ND2	1:A:1055:ASN:H	2.03	0.56
1:A:979:TRP:CE2	1:A:1013:ILE:HG21	2.41	0.55
1:A:1133:GLN:HB3	1:A:1139:SER:HA	1.87	0.55
1:A:1126:ASN:H	1:A:1126:ASN:ND2	2.04	0.54
1:A:962:ARG:HG2	1:A:963:ASN:ND2	2.21	0.54
1:A:1048:LEU:HD21	1:A:1050:LYS:HE3	1.89	0.53
1:A:1132:TYR:CE2	1:A:1277:GLU:HA	2.45	0.52
1:A:984:THR:HG22	1:A:1040:ASP:HB3	1.91	0.52
1:A:1126:ASN:HD22	1:A:1126:ASN:N	2.00	0.52
1:A:1179:ARG:HH21	1:A:1179:ARG:HG2	1.75	0.51
1:A:1090:LEU:HG	1:A:1218:SER:HB3	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:878:ASN:HB2	2:A:726:HOH:O	2.11	0.51
1:A:1197:ILE:HA	1:A:1246:VAL:HG12	1.93	0.51
1:A:1177:VAL:O	1:A:1177:VAL:HG13	2.11	0.51
1:A:1144:TYR:CE1	1:A:1271:LYS:HA	2.44	0.51
1:A:1022:ARG:HG2	1:A:1032:GLU:CB	2.41	0.51
1:A:871:ILE:HD12	1:A:925:ILE:CG2	2.41	0.51
1:A:1101:ASN:ND2	1:A:1152:ARG:HH11	2.09	0.50
1:A:1122:PHE:CE1	1:A:1199:LYS:HE2	2.46	0.50
1:A:1153:LYS:HZ3	1:A:1161:ASN:N	2.10	0.49
1:A:927:TYR:CD1	1:A:932:GLN:HG2	2.48	0.49
1:A:1084:GLU:HB2	1:A:1085:PRO:HD3	1.93	0.49
1:A:1037:ASN:HD22	1:A:1037:ASN:N	2.04	0.48
1:A:1084:GLU:CB	1:A:1085:PRO:HD3	2.44	0.48
1:A:1191:ILE:O	1:A:1192:ALA:CB	2.61	0.48
1:A:981:LEU:HD13	1:A:1038:LEU:HD23	1.96	0.48
1:A:1191:ILE:CD1	1:A:1197:ILE:H	2.26	0.47
1:A:1229:ASN:ND2	1:A:1231:ASN:H	2.12	0.47
1:A:1074:LYS:HE2	2:A:640:HOH:O	2.13	0.47
1:A:1179:ARG:O	1:A:1180:ASP:HB2	2.15	0.47
1:A:973:ASN:ND2	1:A:974:TYR:N	2.55	0.46
1:A:1193:LYS:HB3	1:A:1194:PRO:HD2	1.97	0.46
1:A:1022:ARG:HG2	1:A:1032:GLU:HB3	1.96	0.46
1:A:979:TRP:CD2	1:A:1013:ILE:HG21	2.50	0.46
1:A:1109:LEU:N	1:A:1109:LEU:HD22	2.31	0.46
1:A:1119:ASN:OD1	1:A:1124:ASN:ND2	2.48	0.46
1:A:1168:LYS:O	1:A:1169:ASN:HB2	2.15	0.46
1:A:1272:GLU:HB3	2:A:527:HOH:O	2.16	0.45
1:A:1066:LYS:NZ	1:A:1084:GLU:OE2	2.40	0.45
1:A:965:ASN:HB3	2:A:770:HOH:O	2.17	0.45
1:A:953:ASN:H	1:A:973:ASN:HD21	1.65	0.44
1:A:979:TRP:CZ2	1:A:981:LEU:HG	2.53	0.44
1:A:906:GLN:HE22	1:A:1096:ASN:ND2	2.16	0.43
1:A:976:LYS:HD2	1:A:992:VAL:HG11	2.00	0.43
1:A:904:ARG:HG2	2:A:588:HOH:O	2.17	0.43
1:A:1141:THR:HG21	2:A:610:HOH:O	2.18	0.43
1:A:1229:ASN:HD22	1:A:1231:ASN:H	1.67	0.43
1:A:1000:SER:HA	1:A:1141:THR:HG23	2.01	0.43
1:A:898:TYR:O	1:A:907:PHE:HA	2.19	0.43
1:A:1025:ILE:CG1	1:A:1030:ILE:HG13	2.49	0.43
1:A:968:TRP:HA	1:A:980:THR:O	2.19	0.42
1:A:869:ASN:N	1:A:869:ASN:HD22	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:939:TRP:O	1:A:1062:ILE:HD12	2.20	0.42
1:A:1134:LYS:HA	1:A:1135:PRO:HD2	1.88	0.42
1:A:1162:THR:HG22	2:A:690:HOH:O	2.20	0.42
1:A:1126:ASN:HD22	1:A:1127:GLN:N	2.18	0.42
1:A:1016:ASN:ND2	1:A:1018:LEU:H	2.18	0.41
1:A:1125:ILE:HG12	1:A:1125:ILE:O	2.21	0.41
1:A:1161:ASN:N	2:A:764:HOH:O	2.54	0.41
1:A:1016:ASN:C	1:A:1016:ASN:HD22	2.24	0.41
1:A:1149:VAL:CG1	1:A:1150:ILE:N	2.84	0.41
1:A:1191:ILE:HD12	1:A:1197:ILE:H	1.85	0.41
1:A:926:ILE:O	1:A:926:ILE:HG22	2.21	0.41
1:A:1022:ARG:HG2	1:A:1032:GLU:HB2	2.03	0.40
1:A:874:MET:SD	1:A:891:ILE:HD13	2.61	0.40
1:A:894:ASN:HB3	1:A:916:GLU:HG2	2.03	0.40
1:A:928:ASN:O	1:A:1040:ASP:OD1	2.39	0.40
1:A:1122:PHE:HE1	1:A:1199:LYS:HE2	1.86	0.40
1:A:1125:ILE:HG23	1:A:1125:ILE:O	2.21	0.40
1:A:1229:ASN:ND2	1:A:1233:GLY:H	2.18	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	393/417 (94%)	360 (92%)	28 (7%)	5 (1%)	15 9

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	929	GLY
1	A	1120	SER
1	A	869	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1211	GLY
1	A	1192	ALA

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	369/386 (96%)	353 (96%)	16 (4%)	35 34

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	869	ASN
1	A	872	LEU
1	A	896	ASP
1	A	973	ASN
1	A	981	LEU
1	A	991	LEU
1	A	1016	ASN
1	A	1037	ASN
1	A	1062	ILE
1	A	1110	LEU
1	A	1124	ASN
1	A	1126	ASN
1	A	1142	ARG
1	A	1179	ARG
1	A	1229	ASN
1	A	1278	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	869	ASN
1	A	879	ASN
1	A	928	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	950	ASN
1	A	953	ASN
1	A	963	ASN
1	A	964	ASN
1	A	973	ASN
1	A	988	ASN
1	A	997	GLN
1	A	1006	ASN
1	A	1016	ASN
1	A	1020	ASN
1	A	1028	ASN
1	A	1037	ASN
1	A	1046	ASN
1	A	1055	ASN
1	A	1096	ASN
1	A	1101	ASN
1	A	1124	ASN
1	A	1126	ASN
1	A	1154	ASN
1	A	1229	ASN
1	A	1230	ASN
1	A	1243	ASN
1	A	1244	ASN
1	A	1253	ASN
1	A	1276	GLN
1	A	1278	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/417 (95%)	0.18	12 (3%) 54 62	10, 18, 35, 48	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1210	LEU	6.7
1	A	1191	ILE	6.3
1	A	868	ASP	5.0
1	A	1230	ASN	4.6
1	A	1231	ASN	4.4
1	A	1278	ASN	4.2
1	A	1161	ASN	3.4
1	A	1084	GLU	2.9
1	A	1192	ALA	2.5
1	A	1193	LYS	2.4
1	A	1079	THR	2.2
1	A	1180	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.