



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:09 AM GMT

PDB ID : 2FV1  
Title : UGL\_D88N/dGlcA-GlcNAc  
Authors : Itoh, T.; Hashimoto, W.; Mikami, B.; Murata, K.  
Deposited on : 2006-01-28  
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2417 (1.76-1.72)
Clashscore	102246	2570 (1.76-1.72)
Ramachandran outliers	100387	2544 (1.76-1.72)
Sidechain outliers	100360	2544 (1.76-1.72)
RSRZ outliers	91569	2420 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	<div> <div>8%</div> <div>83%</div> <div>15%</div> <div>.</div> </div>
1	B	377	<div> <div>15%</div> <div>81%</div> <div>18%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NDG	A	401	-	-	-	X
2	GAD	A	402	-	-	-	X
2	NDG	B	401	-	-	-	X
2	GAD	B	402	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unsaturated glucuronyl hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	0	10	0
			3094	1941	564	583	6			
1	B	377	Total	C	N	O	S	0	7	0
			3081	1934	562	579	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	88	ASN	ASP	ENGINEERED	UNP Q9RC92
B	88	ASN	ASP	ENGINEERED	UNP Q9RC92

- Molecule 2 is a polymer of unknown type called SUGAR (2,6-ANHYDRO-3-DEOXY-D-ERYTHRO-HEX-2-ENONIC ACID).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			26	14	1	11		
2	B	2	Total	C	N	O	0	0
			26	14	1	11		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	439	Total	O	0	0
			439	439		
3	B	350	Total	O	0	0
			350	350		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Unsaturated glucuronyl hydrolase



#### • Molecule 1: Unsaturated glucuronyl hydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.73Å 94.89Å 95.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.95 – 1.73 33.64 – 1.73	Depositor EDS
% Data completeness (in resolution range)	95.6 (14.95-1.73) 95.8 (33.64-1.73)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 1.73Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.179 , 0.212 0.181 , 0.213	Depositor DCC
$R_{free}$ test set	7994 reflections (10.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.9	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 54.9	EDS
Estimated twinning fraction	0.013 for -h,l,k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 79806 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7016	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAD, NDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/3212	0.57	0/4346
1	B	0.30	0/3188	0.57	1/4316 (0.0%)
All	All	0.30	0/6400	0.57	1/8662 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	336	GLY	N-CA-C	-5.27	99.93	113.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	355	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3094	0	2925	53	0
1	B	3081	0	2912	64	0
2	A	26	0	20	2	0
2	B	26	0	20	5	0
3	A	439	0	0	6	0
3	B	350	0	0	7	0
All	All	7016	0	5877	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (118) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339[A]:HIS:NE2	2:B:401:NDG:H6C2	1.86	0.90
1:A:377:ARG:OXT	1:A:377:ARG:HD3	1.80	0.81
1:A:131:ILE:H	1:A:176:HIS:HD2	1.38	0.72
1:B:22:ASP:HB3	1:B:69[A]:ARG:NH2	2.04	0.71
1:B:193:HIS:HB3	1:B:209:THR:HB	1.73	0.70
1:A:119:VAL:HA	1:A:122[A]:ARG:NH1	2.05	0.70
1:A:86:HIS:HD2	1:A:88:ASN:H	1.40	0.69
1:B:28:SER:OG	1:B:348:ASP:HB3	1.93	0.68
1:B:111:LYS:HD2	3:B:1649:HOH:O	1.94	0.68
1:B:19:LYS:HE2	3:B:1745:HOH:O	1.94	0.66
1:B:22:ASP:HB3	1:B:69[A]:ARG:HH21	1.61	0.66
1:A:86:HIS:CD2	1:A:88:ASN:HB2	2.32	0.65
1:A:131:ILE:H	1:A:176:HIS:CD2	2.16	0.63
1:B:116:ALA:O	1:B:119:VAL:HG22	1.99	0.63
1:A:87:HIS:CE1	1:A:150:CYS:HB2	2.34	0.62
1:A:133:ALA:CB	1:A:147:ILE:HG12	2.30	0.61
1:B:77:ARG:HG3	1:B:82:GLU:HB3	1.82	0.61
1:B:86:HIS:CD2	1:B:88:ASN:HB2	2.36	0.60
1:B:87:HIS:CD2	1:B:150:CYS:SG	2.95	0.60
1:A:176:HIS:HE1	3:A:1633:HOH:O	1.84	0.60
1:A:168:GLU:OE1	1:A:171:ARG:HD3	2.02	0.60
1:B:65:GLU:HB3	1:B:69[B]:ARG:NH2	2.17	0.59
1:B:298:GLU:CD	1:B:298:GLU:H	2.07	0.58
1:B:179:LYS:HE2	1:B:197:PHE:CE2	2.39	0.58
1:B:130:ILE:HG22	1:B:179:LYS:HD3	1.86	0.57
1:A:174:GLU:HG3	1:A:237:LEU:HD13	1.87	0.57

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:377:ARG:HH21	1:B:377:ARG:HG2	1.70	0.57
1:A:119:VAL:HA	1:A:122[A]:ARG:HH11	1.71	0.56
1:A:111:LYS:HE2	1:A:115:ASP:OD1	2.06	0.55
1:A:119:VAL:HG13	1:A:122[A]:ARG:HH12	1.71	0.54
1:A:75[A]:ARG:HE	1:A:112:LEU:HD13	1.72	0.54
1:A:28:SER:OG	1:A:348:ASP:HB3	2.08	0.54
1:B:339[A]:HIS:HB2	1:B:344:ILE:HB	1.90	0.54
1:B:125:ARG:HH11	1:B:132:GLN:HE21	1.55	0.54
1:A:193:HIS:HB3	1:A:209:THR:HB	1.90	0.53
1:A:39:ASN:ND2	1:A:77:ARG:HH11	2.06	0.53
1:B:176:HIS:HE1	3:B:1639:HOH:O	1.90	0.53
1:B:268:GLU:HB2	3:B:1754:HOH:O	2.08	0.53
1:B:272[A]:GLU:HG3	1:B:274:SER:H	1.72	0.53
1:A:75[A]:ARG:HH21	1:A:112:LEU:HB2	1.74	0.53
1:B:136:PRO:HD2	1:B:139:ASP:HB2	1.91	0.53
1:A:119:VAL:HG22	1:A:122[A]:ARG:NH1	2.25	0.52
1:A:141:GLU:OE2	1:A:142:ASN:ND2	2.43	0.52
1:A:119:VAL:HG22	1:A:122[A]:ARG:HH11	1.74	0.52
1:B:26:HIS:HD2	1:B:43:THR:OG1	1.92	0.52
1:B:39:ASN:ND2	1:B:77:ARG:HH11	2.06	0.52
1:A:147:ILE:HD12	1:A:194:THR:HG22	1.92	0.51
1:B:210:HIS:HB2	3:B:1587:HOH:O	2.10	0.51
1:B:124:TRP:CE3	1:B:172:VAL:HG13	2.46	0.51
1:B:125:ARG:HH11	1:B:132:GLN:NE2	2.08	0.51
1:B:339[B]:HIS:HD2	3:B:1071:HOH:O	1.94	0.51
1:A:75[A]:ARG:HG2	1:A:75[A]:ARG:HH11	1.75	0.50
1:A:120:LEU:HD12	1:A:123:ARG:HG3	1.93	0.50
1:B:134:TRP:CD1	2:B:401:NDG:H8C2	2.47	0.50
1:A:121:MET:SD	1:A:172:VAL:HG21	2.52	0.50
1:A:154:LEU:N	1:A:155:PRO:CD	2.75	0.50
1:A:147:ILE:HD12	1:A:194:THR:CG2	2.42	0.50
1:B:339[A]:HIS:CE1	2:B:401:NDG:H6C2	2.45	0.49
1:B:136:PRO:HD2	1:B:139:ASP:OD2	2.11	0.49
1:A:149:ASP:OD2	1:A:225:TRP:NE1	2.46	0.49
1:B:87:HIS:H	1:B:87:HIS:HD1	1.59	0.48
1:A:86:HIS:HD2	1:A:88:ASN:N	2.09	0.48
1:B:302:GLU:HG2	1:B:306:PHE:CE2	2.49	0.48
1:B:179:LYS:HE2	1:B:197:PHE:CZ	2.49	0.48
1:B:377:ARG:NH2	1:B:377:ARG:HG2	2.28	0.47
1:B:65:GLU:O	1:B:69[B]:ARG:HG3	2.13	0.47
1:A:149:ASP:N	1:A:149:ASP:OD1	2.47	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:128:ALA:O	1:B:130:ILE:HG23	2.14	0.47
1:B:158:LEU:O	1:B:162:GLU:HG3	2.14	0.47
1:B:149:ASP:OD2	1:B:149:ASP:N	2.48	0.47
1:A:26:HIS:HD2	1:A:43:THR:OG1	1.98	0.46
1:B:254:LEU:HD12	1:B:305[A]:ARG:HH22	1.80	0.46
1:B:305[A]:ARG:NH2	1:B:306:PHE:CE1	2.83	0.46
1:A:188:ASP:O	1:A:189:ASP:HB2	2.16	0.46
1:B:212:GLY:O	1:B:341:ARG:HD2	2.16	0.46
1:B:254:LEU:HD12	1:B:305[A]:ARG:NH2	2.30	0.45
1:A:75[A]:ARG:NH2	3:A:1208:HOH:O	2.49	0.45
1:B:132:GLN:HG3	1:B:142:ASN:O	2.17	0.45
1:B:234:SER:OG	1:B:243:LEU:HB2	2.17	0.45
1:A:147:ILE:HG22	1:A:149:ASP:OD1	2.16	0.45
1:A:264:TYR:HA	1:A:277:ARG:HA	1.99	0.45
1:B:130:ILE:CG2	1:B:179:LYS:HD3	2.47	0.44
1:A:61[B]:GLU:OE2	1:A:65:GLU:HG2	2.17	0.44
1:A:83:ASN:HB3	3:A:1517:HOH:O	2.17	0.44
1:A:86:HIS:CD2	1:A:88:ASN:H	2.28	0.44
1:A:374:GLU:HB2	1:A:377:ARG:HB2	1.99	0.44
1:B:23[A]:ARG:HH21	1:B:23[A]:ARG:HG3	1.83	0.44
1:A:119:VAL:HG13	1:A:122[A]:ARG:NH1	2.32	0.43
1:A:150:CYS:HB3	3:A:1633:HOH:O	2.19	0.43
1:B:52:TRP:CE2	1:B:67:ALA:HB1	2.54	0.43
1:A:234[B]:SER:OG	1:A:243:LEU:HB2	2.18	0.43
2:A:401:NDG:H2	2:A:401:NDG:H8C1	1.70	0.42
1:A:107[B]:GLU:OE2	1:A:110:ARG:HD3	2.20	0.42
1:B:88:ASN:ND2	2:B:401:NDG:O7	2.53	0.42
1:A:87:HIS:ND1	1:A:150:CYS:HB2	2.33	0.42
1:B:23[A]:ARG:HE	1:B:36:LEU:HD12	1.85	0.42
1:A:247:LYS:O	1:A:251[A]:ARG:HG3	2.20	0.42
1:B:166:ASP:HA	1:B:167:PRO:HD2	1.91	0.42
1:B:66:GLY:HA2	1:B:69[B]:ARG:HD2	2.01	0.42
1:B:39:ASN:HA	1:B:44:ASP:OD1	2.20	0.42
1:A:124:TRP:HA	1:A:131:ILE:HD13	2.02	0.41
1:A:162:GLU:HB2	3:A:1327:HOH:O	2.20	0.41
2:A:402:GAD:H2	3:A:1744:HOH:O	2.19	0.41
1:B:125:ARG:NH1	1:B:132:GLN:NE2	2.67	0.41
1:A:187:GLY:HA3	1:B:185:VAL:HB	2.01	0.41
1:A:179:LYS:HE2	1:A:197:PHE:CE2	2.55	0.41
1:B:176:HIS:CE1	3:B:1639:HOH:O	2.71	0.41
1:B:298:GLU:HG3	1:B:303[A]:ARG:NH1	2.36	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:ASN:ND2	1:B:328:GLU:HG2	2.36	0.41
1:A:145:ARG:NH2	1:A:194:THR:HG21	2.35	0.40
1:B:305[A]:ARG:NH2	1:B:306:PHE:HE1	2.19	0.40
1:B:32:ASN:HD21	1:B:328:GLU:HG2	1.86	0.40
1:B:154:LEU:N	1:B:155:PRO:CD	2.84	0.40
1:B:338:TYR:OH	2:B:401:NDG:H5	2.21	0.40
1:B:124:TRP:CD2	1:B:172:VAL:HG13	2.56	0.40
1:B:300:ASP:HA	1:B:301:PRO:HD2	1.97	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/377 (102%)	372 (97%)	13 (3%)	0	100	100
1	B	382/377 (101%)	360 (94%)	19 (5%)	3 (1%)	24	7
All	All	767/754 (102%)	732 (95%)	32 (4%)	3 (0%)	39	20

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	136	PRO
1	B	88	ASN
1	B	141	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/304 (103%)	301 (96%)	13 (4%)	37	12
1	B	311/304 (102%)	305 (98%)	6 (2%)	65	42
All	All	625/608 (103%)	606 (97%)	19 (3%)	54	22

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	61[A]	GLU
1	A	61[B]	GLU
1	A	87	HIS
1	A	118	ASP
1	A	123	ARG
1	A	141	GLU
1	A	149	ASP
1	A	211[A]	GLN
1	A	211[B]	GLN
1	A	228	TYR
1	A	339	HIS
1	A	377	ARG
1	B	39	ASN
1	B	65	GLU
1	B	149	ASP
1	B	228	TYR
1	B	272[A]	GLU
1	B	272[B]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	HIS
1	A	39	ASN
1	A	83	ASN
1	A	86	HIS
1	A	176	HIS
1	A	193	HIS
1	A	223	GLN
1	A	304	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	26	HIS
1	B	39	ASN
1	B	86	HIS
1	B	132	GLN
1	B	176	HIS
1	B	223	GLN
1	B	304	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NDG	A	401	2	15,15,15	0.61	0	17,21,21	0.83	0
2	GAD	A	402	1,2	7,11,11	2.11	2 (28%)	8,15,15	1.06	1 (12%)
2	NDG	B	401	2	15,15,15	0.61	0	17,21,21	0.79	0
2	GAD	B	402	1,2	7,11,11	2.14	2 (28%)	8,15,15	1.01	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDG	A	401	2	-	0/6/26/26	0/1/1/1
2	GAD	A	402	1,2	-	0/0/17/17	0/1/1/1
2	NDG	B	401	2	-	0/6/26/26	0/1/1/1
2	GAD	B	402	1,2	-	0/0/17/17	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	GAD	C3-C4	2.17	1.53	1.50
2	B	402	GAD	C3-C4	2.19	1.53	1.50
2	A	402	GAD	C4-C5	5.05	1.39	1.32
2	B	402	GAD	C4-C5	5.14	1.39	1.32

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	GAD	O5-C1-C2	-2.82	107.77	111.06
2	B	402	GAD	O5-C1-C2	-2.71	107.90	111.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NDG	1	0
2	A	402	GAD	1	0
2	B	401	NDG	5	0

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	377/377 (100%)	0.40	29 (7%)	16 20	7, 14, 31, 46	0
1	B	377/377 (100%)	0.99	56 (14%)	3 4	7, 17, 45, 52	0
All	All	754/754 (100%)	0.69	85 (11%)	7 8	7, 15, 41, 52	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	134	TRP	14.7
1	B	136	PRO	12.2
1	A	134	TRP	11.7
1	B	137	LYS	11.5
1	B	124	TRP	11.2
1	B	140	PRO	10.9
1	B	131	ILE	9.9
1	B	138	GLY	9.9
1	B	135	GLY	9.1
1	A	376	GLY	8.4
1	B	128	ALA	8.4
1	B	376	GLY	8.4
1	B	141	GLU	8.1
1	B	126	ALA	8.0
1	B	129	GLY	8.0
1	A	136	PRO	7.8
1	B	139	ASP	7.1
1	B	127	ASP	6.6
1	B	143	GLY	6.5
1	B	142	ASN	6.4
1	B	133	ALA	6.1
1	B	85	ASP	5.7
1	A	141	GLU	5.7
1	A	140	PRO	5.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	199	PRO	5.5
1	B	87	HIS	5.5
1	B	210	HIS	5.4
1	B	200	GLU	5.1
1	B	125	ARG	4.9
1	B	130	ILE	4.7
1	B	122	ARG	4.6
1	B	123	ARG	4.5
1	B	301	PRO	4.3
1	B	297	ASP	4.3
1	B	377	ARG	4.3
1	B	298	GLU	4.3
1	B	132	GLN	4.2
1	B	80	ARG	4.1
1	A	377	ARG	4.1
1	A	138	GLY	3.9
1	B	201	ASN	3.9
1	B	81	PHE	3.9
1	B	202	GLY	3.8
1	A	135	GLY	3.8
1	A	139	ASP	3.7
1	A	225	TRP	3.7
1	A	152	LEU	3.6
1	B	152	LEU	3.5
1	A	124	TRP	3.3
1	B	86	HIS	3.1
1	B	299	SER	3.1
1	B	1	MET	3.1
1	A	137	LYS	3.0
1	A	149	ASP	3.0
1	B	144	GLY	2.9
1	B	167	PRO	2.9
1	B	203	ASN	2.9
1	A	46	PHE	2.8
1	A	1	MET	2.8
1	A	297	ASP	2.7
1	B	171	ARG	2.7
1	B	225	TRP	2.6
1	B	46	PHE	2.5
1	A	91	PHE	2.5
1	A	126	ALA	2.4
1	A	131	ILE	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	142	ASN	2.3
1	B	121	MET	2.3
1	B	83	ASN	2.3
1	A	284	ILE	2.3
1	A	356	TYR	2.3
1	B	145	ARG	2.2
1	A	151	LEU	2.2
1	A	127	ASP	2.2
1	A	122[A]	ARG	2.1
1	A	150	CYS	2.1
1	B	224	ALA	2.1
1	A	200	GLU	2.1
1	B	69[A]	ARG	2.1
1	B	228	TYR	2.0
1	A	45	GLY	2.0
1	B	351	ILE	2.0
1	B	356	TYR	2.0
1	B	120	LEU	2.0
1	B	79	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	GAD	A	402	11/11	0.68	0.54	6.93	17,43,54,56	0
2	GAD	B	402	11/11	0.57	0.53	5.00	18,44,54,57	0
2	NDG	A	401	15/15	0.27	0.66	4.88	46,50,54,55	0
2	NDG	B	401	15/15	0.52	0.61	2.69	49,50,55,55	0

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.