



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:09 AM GMT

PDB ID : 2FVS
Title : A Structural Study of the CA Dinucleotide Step in the Integrase Processing Site of Moloney Murine Leukemia Virus
Authors : Montano, S.P.; Cote, M.L.; Roth, M.J.; Georgiadis, M.M.
Deposited on : 2006-01-31
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	16	<div> <div style="width: 31%; background-color: red;"></div> <div style="width: 69%; background-color: yellow;"></div> </div> <div> <div style="width: 31%; background-color: red;"></div> <div style="width: 69%; background-color: yellow;"></div> </div>
2	A	255	<div> <div style="width: 9%; background-color: red;"></div> <div style="width: 73%; background-color: green;"></div> <div style="width: 22%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> </div> <div> <div style="width: 9%; background-color: red;"></div> <div style="width: 73%; background-color: green;"></div> <div style="width: 22%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*AP*CP*AP*AP*TP*GP*AP*TP*CP*AP*TP*TP*GP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	16	Total	C	N	O	P	0	0	0
			325	157	59	94	15			

- Molecule 2 is a protein called reverse transcriptase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	255	Total	C	N	O	S	0	0	0
			2041	1311	356	367	7			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	85	Total	O	0	0
			85	85		
3	B	1	Total	O	0	0
			1	1		

● Molecule 1: 5'-D(*CP*AP*CP*AP*AP*TP*GP*AP*TP*CP*AP*TP*TP*GP*TP*G)-3'



- Chain A:  9% 73% 22% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	54.24Å 145.68Å 46.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.35 46.94 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.35) 96.4 (46.94-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.29Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.243 , 0.265 0.242 , 0.263	Depositor DCC
R_{free} test set	765 reflections (4.87%)	DCC
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 34.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 17052 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2452	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.32	0/364	0.70	0/560
2	A	0.37	0/2097	0.61	0/2858
All	All	0.36	0/2461	0.63	0/3418

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	325	0	183	18	0
2	A	2041	0	2056	49	0
3	A	85	0	0	2	0
3	B	1	0	0	1	0
All	All	2452	0	2239	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:173:ARG:HA	2:A:173:ARG:HE	1.25	1.02

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:218:ILE:HB	2:A:229:ALA:HB3	1.59	0.84
1:B:9:DT:H2"	1:B:10:DC:H5'	1.69	0.74
2:A:173:ARG:HA	2:A:173:ARG:NE	2.02	0.73
2:A:274:LYS:HG2	2:A:275:GLU:HG3	1.71	0.72
2:A:206:ASP:HB3	2:A:250:LEU:HD13	1.70	0.71
2:A:103:LYS:HE3	2:A:110:ARG:HH22	1.55	0.70
1:B:2:DA:H2"	1:B:3:DC:H5'	1.74	0.68
2:A:26:LEU:HD23	2:A:26:LEU:O	1.96	0.65
2:A:61:ILE:HD11	2:A:117:GLU:HG3	1.80	0.64
1:B:15:DT:H2"	1:B:16:DG:C8	2.33	0.63
2:A:173:ARG:CA	2:A:173:ARG:HE	2.02	0.63
1:B:14:DG:H1'	1:B:15:DT:H5"	1.81	0.62
1:B:14:DG:H2"	1:B:15:DT:C5'	2.31	0.60
2:A:174:ASP:OD1	2:A:175:PRO:HD2	2.01	0.60
1:B:4:DA:H2"	1:B:5:DA:H5'	1.85	0.59
1:B:14:DG:H2"	1:B:15:DT:H5'	1.84	0.58
2:A:103:LYS:HE3	2:A:110:ARG:NH2	2.17	0.58
2:A:233:GLU:O	2:A:237:GLN:HG3	2.04	0.57
1:B:5:DA:H2"	1:B:6:DT:C5'	2.34	0.57
2:A:71:ARG:NH1	2:A:175:PRO:HG3	2.20	0.56
2:A:277:GLN:O	2:A:278:ARG:HB2	2.05	0.56
2:A:75:LYS:HB3	2:A:76:PRO:HD3	1.87	0.55
1:B:15:DT:H1'	3:B:63:HOH:O	2.05	0.55
2:A:25:TRP:C	2:A:27:SER:H	2.10	0.55
2:A:131:ASN:HD21	2:A:134:ASN:ND2	2.05	0.55
1:B:8:DA:H1'	1:B:9:DT:H5"	1.88	0.54
2:A:102:LYS:HD2	2:A:106:THR:HA	1.90	0.54
2:A:62:LYS:HE2	2:A:62:LYS:HA	1.90	0.53
1:B:9:DT:H2"	1:B:10:DC:C5'	2.39	0.53
2:A:176:GLU:CD	2:A:177:MET:H	2.12	0.53
2:A:103:LYS:CB	2:A:104:PRO:HD3	2.39	0.53
1:B:5:DA:H2"	1:B:6:DT:H5'	1.91	0.52
2:A:265:GLN:OE1	2:A:274:LYS:HD2	2.10	0.52
2:A:174:ASP:O	2:A:175:PRO:C	2.49	0.51
2:A:220:LEU:HD22	2:A:227:LEU:HD23	1.92	0.50
2:A:237:GLN:O	2:A:241:ARG:HG3	2.12	0.49
2:A:179:ILE:HG22	2:A:180:SER:N	2.29	0.48
2:A:265:GLN:HA	2:A:273:LEU:O	2.14	0.47
2:A:174:ASP:HB3	2:A:178:GLY:CA	2.45	0.47
2:A:62:LYS:HE2	2:A:63:GLN:H	1.80	0.47
2:A:179:ILE:O	2:A:180:SER:C	2.53	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:68:GLN:HG2	2:A:72:LEU:HD22	1.98	0.46
2:A:119:ASN:HB3	2:A:193:LYS:HD3	1.98	0.46
2:A:203:LEU:HD23	3:A:283:HOH:O	2.16	0.45
1:B:10:DC:H5'	1:B:10:DC:H6	1.82	0.44
2:A:25:TRP:C	2:A:27:SER:N	2.71	0.44
2:A:214:HIS:N	2:A:215:PRO:HD3	2.32	0.44
2:A:158:LEU:HD11	2:A:199:PHE:HA	2.00	0.43
1:B:9:DT:H1'	1:B:10:DC:H5''	1.99	0.43
2:A:99:LEU:HA	2:A:100:PRO:HD3	1.79	0.43
2:A:132:PRO:O	2:A:136:LEU:HD23	2.19	0.42
2:A:103:LYS:NZ	2:A:110:ARG:HH12	2.17	0.42
2:A:257:LYS:HD2	2:A:257:LYS:O	2.20	0.42
2:A:28:ASP:C	2:A:30:PRO:HD3	2.41	0.42
2:A:41:LEU:O	2:A:251:GLY:HA3	2.20	0.41
1:B:14:DG:C1'	1:B:15:DT:H5''	2.50	0.41
2:A:120:LYS:HG3	3:A:333:HOH:O	2.20	0.41
1:B:14:DG:C2'	1:B:15:DT:H5''	2.49	0.41
1:B:9:DT:H6	1:B:9:DT:H5'	1.85	0.41
2:A:103:LYS:HB2	2:A:104:PRO:HD3	2.02	0.41
2:A:277:GLN:NE2	2:A:277:GLN:HA	2.36	0.41
2:A:195:SER:HB2	2:A:196:PRO:HD3	2.03	0.41
2:A:197:THR:O	2:A:201:GLU:HG3	2.21	0.40
2:A:72:LEU:HD12	2:A:72:LEU:HA	1.90	0.40
1:B:10:DC:C6	1:B:10:DC:H5'	2.57	0.40
2:A:43:VAL:HG23	2:A:44:ARG:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	253/255 (99%)	234 (92%)	9 (4%)	10 (4%)	4 1

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	104	PRO
2	A	108	ASP
2	A	175	PRO
2	A	106	THR
2	A	180	SER
2	A	223	VAL
2	A	103	LYS
2	A	26	LEU
2	A	181	GLY
2	A	179	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	224/224 (100%)	218 (97%)	6 (3%)	52 67

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	62	LYS
2	A	72	LEU
2	A	173	ARG
2	A	175	PRO
2	A	176	GLU
2	A	177	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	31	GLN
2	A	79	GLN
2	A	84	GLN
2	A	91	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	134	ASN
2	A	144	GLN
2	A	214	HIS
2	A	237	GLN
2	A	245	GLN
2	A	249	ASN
2	A	277	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	B	16/16 (100%)	-0.02	0	100	100	38, 60, 70, 71	0
2	A	255/255 (100%)	0.54	23 (9%)	12	18	20, 41, 85, 101	0
All	All	271/271 (100%)	0.50	23 (8%)	13	21	20, 42, 84, 101	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	179	ILE	8.3
2	A	175	PRO	7.9
2	A	177	MET	7.3
2	A	105	GLY	5.2
2	A	106	THR	4.4
2	A	180	SER	4.1
2	A	107	ASN	4.1
2	A	25	TRP	3.2
2	A	178	GLY	3.1
2	A	275	GLU	3.1
2	A	237	GLN	3.0
2	A	103	LYS	2.9
2	A	108	ASP	2.8
2	A	26	LEU	2.7
2	A	145	TRP	2.5
2	A	136	LEU	2.3
2	A	29	PHE	2.3
2	A	109	TYR	2.3
2	A	64	TYR	2.3
2	A	264	LYS	2.2
2	A	176	GLU	2.1
2	A	30	PRO	2.1
2	A	104	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.