



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:15 AM GMT

PDB ID : 2GAZ  
Title : Mycobacterial lipoglycan presentation by CD1d  
Authors : Zajonc, D.M.  
Deposited on : 2006-03-09  
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2700 (2.64-2.60)
Clashscore	102246	3065 (2.64-2.60)
Ramachandran outliers	100387	3015 (2.64-2.60)
Sidechain outliers	100360	3015 (2.64-2.60)
RSRZ outliers	91569	2706 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	<div> <div>5%</div> <div>78%</div> <div>15%</div> <div>6%</div> </div>
2	B	99	<div> <div>%</div> <div>82%</div> <div>15%</div> <div>••</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1d1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	S	0	0	0
			2161	1377	373	398	13			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	HIS	ASP	SEE REMARK 999	UNP P11609
A	280	HIS	-	EXPRESSION TAG	UNP P11609
A	281	HIS	-	EXPRESSION TAG	UNP P11609
A	282	HIS	-	EXPRESSION TAG	UNP P11609
A	283	HIS	-	EXPRESSION TAG	UNP P11609
A	284	HIS	-	EXPRESSION TAG	UNP P11609
A	285	HIS	-	EXPRESSION TAG	UNP P11609

- Molecule 2 is a protein called beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	98	Total	C	N	O	S	4	0	0
			810	517	137	149	7			

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 5 is (2R)-3-[(HYDROXY{[(2R,3R,5S,6R)-3,4,5-TRIHYDROXY-2,6-BIS(ALPHA-D-MANNOPYRANOSYLOXY)CYCLOHEXYL]OXY}PHOSPHORYL)OXY]PROPANE-1,2-DIYL DIHEXADECANOATE (three-letter code: XPX) (formula: C<sub>53</sub>H<sub>99</sub>O<sub>23</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	P	0	0
			77	53	23	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	32	Total	O	0	0
			32	32		
6	B	13	Total	O	0	0
			13	13		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-cell surface glycoprotein CD1d1



- Molecule 2: beta-2-microglobulin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.84Å 110.76Å 107.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.34 – 2.61 48.32 – 2.61	Depositor EDS
% Data completeness (in resolution range)	96.7 (48.34-2.61) 96.7 (48.32-2.61)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.59 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.212 , 0.280 0.211 , 0.277	Depositor DCC
$R_{free}$ test set	760 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.2	Xtriage
Anisotropy	0.206	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.8	EDS
Estimated twinning fraction	0.020 for -h,l,k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 15274 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3182	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, XPX, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.70	0/2224	0.76	1/3023 (0.0%)
2	B	0.74	0/836	0.74	0/1133
All	All	0.71	0/3060	0.76	1/4156 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	VAL	CB-CA-C	-5.81	100.36	111.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2161	0	2062	27	0
2	B	810	0	783	11	0
3	A	28	0	26	1	0
4	A	61	0	52	0	0
5	A	77	0	98	4	0
6	A	32	0	0	1	0
6	B	13	0	0	1	0
All	All	3182	0	3021	36	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:33:PRO:HG3	2:B:62:PHE:CZ	2.20	0.76
2:B:33:PRO:HG3	2:B:62:PHE:CE1	2.19	0.76
2:B:21:ASN:HB3	2:B:70:PHE:CE1	2.22	0.75
1:A:59:SER:H	1:A:62:GLN:NE2	1.89	0.69
1:A:118:VAL:HG11	5:A:525:XPX:H231	1.76	0.68
1:A:264:ARG:HG3	1:A:276:ILE:HG12	1.75	0.66
1:A:59:SER:H	1:A:62:GLN:HE21	1.43	0.66
5:A:525:XPX:H121	5:A:525:XPX:H9	1.78	0.64
1:A:124:TYR:CE2	1:A:136:VAL:HG21	2.34	0.63
1:A:15:MET:HG2	2:B:62:PHE:HE2	1.66	0.59
1:A:124:TYR:CZ	1:A:136:VAL:HG21	2.38	0.59
1:A:30:VAL:HG12	1:A:37:THR:OG1	2.07	0.54
1:A:73:TYR:CD1	5:A:525:XPX:H332	2.43	0.54
1:A:106:MET:HG3	1:A:172:VAL:HG11	1.91	0.53
1:A:14:GLN:HB3	1:A:100:LEU:HB3	1.90	0.53
1:A:182:ASP:O	1:A:185:LYS:HG2	2.08	0.53
1:A:209:HIS:HE1	2:B:99:MET:OXT	1.93	0.52
3:A:510:NAG:H61	6:A:534:HOH:O	2.10	0.51
1:A:160:VAL:HG22	5:A:525:XPX:H342	1.93	0.50
1:A:85:VAL:HG11	1:A:94:TYR:CZ	2.46	0.50
1:A:195:SER:HB3	1:A:206:LEU:HD12	1.94	0.49
1:A:69:MET:SD	1:A:163:LEU:HD11	2.53	0.48
2:B:57:SER:HB2	6:B:107:HOH:O	2.13	0.48
1:A:219:TRP:HB3	1:A:266:LYS:HB2	1.95	0.48
1:A:219:TRP:CE3	1:A:266:LYS:HG3	2.50	0.47
1:A:264:ARG:HD3	1:A:274:ASP:OD2	2.16	0.46
1:A:123:LYS:O	1:A:125:VAL:HG13	2.16	0.45
1:A:96:ILE:HG23	1:A:120:PHE:HE1	1.82	0.45
1:A:209:HIS:CE1	2:B:99:MET:OXT	2.70	0.44
2:B:29:GLN:HA	2:B:61:SER:HB2	2.00	0.43
1:A:18:PHE:N	1:A:18:PHE:CD1	2.87	0.43
1:A:51:LYS:HA	1:A:51:LYS:HD3	1.91	0.42
1:A:125:VAL:HG23	1:A:126:VAL:HG23	2.02	0.41
2:B:10:TYR:CD1	2:B:10:TYR:N	2.89	0.41
1:A:39:ARG:NE	2:B:53:ASP:OD2	2.42	0.41
2:B:71:THR:HA	2:B:72:PRO:HD2	1.92	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/285 (93%)	258 (97%)	7 (3%)	0	100	100
2	B	96/99 (97%)	89 (93%)	6 (6%)	1 (1%)	19	37
All	All	361/384 (94%)	347 (96%)	13 (4%)	1 (0%)	46	70

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	47	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/249 (94%)	221 (94%)	13 (6%)	26	49
2	B	92/93 (99%)	86 (94%)	6 (6%)	21	41
All	All	326/342 (95%)	307 (94%)	19 (6%)	25	47

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ARG
1	A	61	GLN
1	A	64	GLU
1	A	69	MET
1	A	79	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	88	MET
1	A	127	ARG
1	A	164	LEU
1	A	206	LEU
1	A	209	HIS
1	A	218	VAL
1	A	226	ASP
1	A	265	VAL
2	B	45	LYS
2	B	58	LYS
2	B	59	ASP
2	B	64	ILE
2	B	70	PHE
2	B	71	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	203	HIS
1	A	230	GLN
2	B	31	HIS
2	B	67	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	520	1,4	14,14,15	0.71	0	15,19,21	1.84	2 (13%)
4	NAG	A	521	4	14,14,15	0.52	0	15,19,21	1.46	2 (13%)
4	BMA	A	522	4	11,11,12	0.60	0	14,15,17	2.42	4 (28%)
4	MAN	A	523	4	11,11,12	0.73	0	14,15,17	1.22	1 (7%)
4	MAN	A	524	4	11,11,12	0.61	0	14,15,17	1.30	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	520	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	521	4	-	0/6/23/26	0/1/1/1
4	BMA	A	522	4	-	0/2/19/22	0/1/1/1
4	MAN	A	523	4	-	0/2/19/22	0/1/1/1
4	MAN	A	524	4	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	522	BMA	C6-C5-C4	-2.30	107.33	113.02
4	A	520	NAG	O3-C3-C2	-2.01	105.12	109.11
4	A	521	NAG	O7-C7-N2	2.00	125.94	121.86
4	A	523	MAN	O2-C2-C1	2.70	114.62	109.21
4	A	522	BMA	O5-C1-C2	3.03	115.78	110.86
4	A	522	BMA	C1-C2-C3	3.24	113.37	109.54
4	A	524	MAN	C1-O5-C5	3.97	117.28	112.25
4	A	521	NAG	C1-O5-C5	4.67	118.17	112.25
4	A	520	NAG	C1-O5-C5	5.66	119.44	112.25
4	A	522	BMA	C1-O5-C5	7.23	121.42	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAG	A	501	1	14,14,15	0.56	0	15,19,21	2.06	2 (13%)
3	NAG	A	510	1	14,14,15	0.67	0	15,19,21	1.48	3 (20%)
5	XPX	A	525	-	79,79,79	0.98	2 (2%)	99,103,103	1.54	13 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	501	1	-	0/6/23/26	0/1/1/1
3	NAG	A	510	1	-	0/6/23/26	0/1/1/1
5	XPX	A	525	-	-	0/62/126/126	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	525	XPX	O1-C1	2.04	1.47	1.41
5	A	525	XPX	O2-C48	3.48	1.52	1.43

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	525	XPX	C12-C13-C30	-2.78	105.57	112.07
3	A	510	NAG	C3-C4-C5	-2.32	106.15	110.20
5	A	525	XPX	C49-C48-C47	-2.31	106.35	111.44
5	A	525	XPX	C11-C6-C10	-2.18	107.63	113.02

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	NAG	C6-C5-C4	-2.12	107.79	113.02
5	A	525	XPX	C30-O19-C31	-2.10	110.97	116.85
3	A	510	NAG	O4-C4-C5	2.12	114.86	109.24
5	A	525	XPX	O1-C52-C47	2.14	113.09	107.49
5	A	525	XPX	C3-C4-C	2.23	114.09	110.20
5	A	525	XPX	C7-O2-C48	2.95	125.72	118.01
5	A	525	XPX	O7-C6-C10	3.05	115.40	109.68
5	A	525	XPX	O19-C31-C32	3.05	121.21	111.90
3	A	510	NAG	C1-O5-C5	3.38	116.54	112.25
5	A	525	XPX	O2-C7-C8	3.41	116.41	108.10
5	A	525	XPX	O17-C14-C15	4.13	120.51	111.53
5	A	525	XPX	O-C-C4	4.97	119.01	109.68
3	A	501	NAG	C1-O5-C5	7.19	121.37	112.25
5	A	525	XPX	O2-C48-C49	7.81	127.32	107.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	510	NAG	1	0
5	A	525	XPX	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	269/285 (94%)	0.31	15 (5%) 28 21	36, 44, 57, 69	0
2	B	98/99 (98%)	0.04	1 (1%) 84 81	38, 45, 52, 64	1 (1%)
All	All	367/384 (95%)	0.24	16 (4%) 38 31	36, 44, 56, 69	1 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	108	PRO	5.3
1	A	109	GLY	4.0
1	A	201	HIS	3.7
1	A	88	MET	3.4
1	A	21	ARG	3.3
1	A	93	ASP	3.0
1	A	271	GLY	2.6
1	A	107	TYR	2.5
1	A	19	ALA	2.5
1	A	110	ASN	2.5
1	A	22	SER	2.4
2	B	99	MET	2.4
1	A	255	ALA	2.3
1	A	199	SER	2.2
1	A	262	ALA	2.1
1	A	86	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MAN	A	524	11/12	0.79	0.28	1.55	65,67,69,70	0
4	NAG	A	520	14/15	0.94	0.18	0.06	36,46,50,52	0
4	MAN	A	523	11/12	0.73	0.32	-	69,71,71,71	0
4	NAG	A	521	14/15	0.83	0.26	-	57,61,63,68	0
4	BMA	A	522	11/12	0.70	0.26	-	66,69,70,71	0

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	XPX	A	525	77/77	0.84	0.23	1.16	43,61,81,83	0
3	NAG	A	501	14/15	0.70	0.39	-	53,55,57,57	0
3	NAG	A	510	14/15	0.90	0.22	-	46,51,54,54	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.