



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:35 PM GMT

PDB ID : 1GC1
Title : HIV-1 GP120 CORE COMPLEXED WITH CD4 AND A NEUTRALIZING HUMAN ANTIBODY
Authors : Kwong, P.D.; Wyatt, R.; Robinson, J.; Sweet, R.W.; Sodroski, J.; Hendrickson, W.A.
Deposited on : 1998-06-15
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	G	321	
2	C	185	
3	L	213	
4	H	229	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENVELOPE PROTEIN GP120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	297	Total	C	N	O	S	0	0	0
			2300	1441	401	438	20			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	94	ASN	ASP	CONFLICT	UNP P04578
G	346	ALA	ASP	CONFLICT	UNP P04578
G	470	PRO	LEU	CONFLICT	UNP P04578

- Molecule 2 is a protein called CD4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	181	Total	C	N	O	S	0	0	0
			1412	885	247	276	4			

- Molecule 3 is a protein called ANTIBODY 17B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	0	0
			1645	1028	281	332	4			

- Molecule 4 is a protein called ANTIBODY 17B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	229	Total	C	N	O	S	0	0	0
			1723	1088	291	339	5			

- Molecule 5 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	G	2	Total	C	N	O	0	0
			24	14	1	9		

- Molecule 7 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	G	2	Total	C	N	O	0	0
			24	14	1	9		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	G	2	Total	C	N	O	0	0
			24	14	1	9		
7	G	2	Total	C	N	O	0	0
			24	14	1	9		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	116	Total	O	0	0
			116	116		
8	G	208	Total	O	1	0
			208	208		
8	H	143	Total	O	0	0
			143	143		
8	L	136	Total	O	0	0
			136	136		



• Molecule 4: ANTIBODY 17B



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	71.64Å 88.13Å 196.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 2.50	Depositor
% Data completeness (in resolution range)	74.9 (5.00-2.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.210 , 0.302	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7877	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, NDG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.39	0/2346	0.75	0/3177
2	C	0.38	0/1432	0.71	0/1930
3	L	0.40	0/1682	0.74	0/2287
4	H	0.41	0/1763	0.71	0/2402
All	All	0.40	0/7223	0.73	0/9796

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2300	0	2242	145	0
2	C	1412	0	1444	79	0
3	L	1645	0	1591	77	0
4	H	1723	0	1699	64	0
5	G	98	0	91	9	0
6	G	24	0	22	0	0
7	G	72	0	66	10	0
8	C	116	0	0	10	0
8	G	208	0	0	30	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	H	143	0	0	4	0
8	L	136	0	0	10	0
All	All	7877	0	7155	363	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 363 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:138:PRO:HD3	4:H:224:LYS:HE2	1.45	0.96
2:C:111:GLY:HA2	2:C:148:GLN:HG3	1.49	0.94
1:G:124:PRO:HD2	1:G:198:THR:HA	1.52	0.90
3:L:151:LYS:HE2	3:L:154:ASN:HA	1.59	0.85
1:G:363:GLN:HE21	1:G:470:PRO:HG3	1.43	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	293/321 (91%)	242 (83%)	40 (14%)	11 (4%)	4	5
2	C	179/185 (97%)	150 (84%)	24 (13%)	5 (3%)	6	9
3	L	211/213 (99%)	185 (88%)	19 (9%)	7 (3%)	5	6
4	H	227/229 (99%)	201 (88%)	16 (7%)	10 (4%)	3	3
All	All	910/948 (96%)	778 (86%)	99 (11%)	33 (4%)	4	5

5 of 33 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	99	ASP
1	G	118	PRO
1	G	258	GLN
1	G	350	ARG
1	G	352	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	262/283 (93%)	240 (92%)	22 (8%)	14	25
2	C	164/167 (98%)	144 (88%)	20 (12%)	6	11
3	L	184/184 (100%)	161 (88%)	23 (12%)	6	10
4	H	193/193 (100%)	174 (90%)	19 (10%)	10	19
All	All	803/827 (97%)	719 (90%)	84 (10%)	8	16

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	139	GLN
3	L	40	PRO
4	H	139	LEU
2	C	167	LYS
3	L	3	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	462	ASN
2	C	137	ASN
4	H	163	GLN
1	G	478	ASN
2	C	66	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	NDG	G	734	1,6	14,14,15	0.60	0	15,19,21	1.21	1 (6%)
6	FUC	G	735	6	10,10,11	0.41	0	14,14,16	0.60	0
7	NAG	G	795	1,7	14,14,15	0.50	0	15,19,21	1.11	1 (6%)
7	FUC	G	796	7	10,10,11	0.59	0	14,14,16	0.72	0
7	NAG	G	892	1,7	14,14,15	0.50	0	15,19,21	0.58	0
7	FUC	G	893	7	10,10,11	0.54	0	14,14,16	1.00	1 (7%)
7	NAG	G	948	1,7	14,14,15	0.57	0	15,19,21	0.73	0
7	FUC	G	949	7	10,10,11	0.55	0	14,14,16	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NDG	G	734	1,6	-	0/6/23/26	0/1/1/1
6	FUC	G	735	6	-	0/0/17/20	0/1/1/1
7	NAG	G	795	1,7	-	0/6/23/26	0/1/1/1
7	FUC	G	796	7	-	0/0/17/20	0/1/1/1
7	NAG	G	892	1,7	-	0/6/23/26	0/1/1/1
7	FUC	G	893	7	-	0/0/17/20	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	G	948	1,7	-	0/6/23/26	0/1/1/1
7	FUC	G	949	7	-	0/0/17/20	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	795	NAG	C4-C3-C2	-2.57	107.24	111.23
7	G	893	FUC	C1-C2-C3	2.44	112.43	109.54
6	G	734	NDG	C1-O-C5	2.82	115.82	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	G	795	NAG	2	0
7	G	796	FUC	1	0
7	G	892	NAG	4	0
7	G	948	NAG	4	0
7	G	949	FUC	2	0

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	G	697	1	14,14,15	0.55	0	15,19,21	0.69	0
5	NAG	G	762	1	14,14,15	0.43	0	15,19,21	0.72	0
5	NAG	G	776	1	14,14,15	0.41	0	15,19,21	0.41	0
5	NAG	G	789	1	14,14,15	0.53	0	15,19,21	0.59	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	G	832	1	14,14,15	0.70	1 (7%)	15,19,21	0.72	0
5	NAG	G	839	1	14,14,15	0.47	0	15,19,21	0.90	0
5	NAG	G	886	1	14,14,15	0.52	0	15,19,21	0.89	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	G	697	1	-	0/6/23/26	0/1/1/1
5	NAG	G	762	1	-	0/6/23/26	0/1/1/1
5	NAG	G	776	1	-	0/6/23/26	0/1/1/1
5	NAG	G	789	1	-	0/6/23/26	0/1/1/1
5	NAG	G	832	1	-	0/6/23/26	0/1/1/1
5	NAG	G	839	1	-	0/6/23/26	0/1/1/1
5	NAG	G	886	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	832	NAG	C1-C2	2.01	1.55	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	886	NAG	C4-C3-C2	-2.33	107.61	111.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	697	NAG	1	0
5	G	789	NAG	3	0
5	G	832	NAG	2	0
5	G	839	NAG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.