



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:16 AM GMT

PDB ID : 2GFO  
Title : Structure of the Catalytic Domain of Human Ubiquitin Carboxyl-terminal Hydrolase 8  
Authors : Walker, J.R.; Avvakumov, G.V; Xue, S.; Newman, E.M.; Finerty Jr., P.J.; Butler-Cole, C.; Weigelt, J.; Sundstrom, M.; Arrowsmith, C.; Edwards, A.; Bochkarev, A.; Dhe-Paganon, S.; Structural Genomics Consortium (SGC)  
Deposited on : 2006-03-22  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	<div> <div>5%</div> <div>73%</div> <div>11%</div> <div>•</div> <div>14%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2938 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	Se	0	0	0
			2746	1748	479	504	11	4			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	715	MSE	-	CLONING ARTIFACT	UNP P40818
A	716	GLY	-	CLONING ARTIFACT	UNP P40818
A	717	SER	-	CLONING ARTIFACT	UNP P40818
A	718	SER	-	CLONING ARTIFACT	UNP P40818
A	719	HIS	-	EXPRESSION TAG	UNP P40818
A	720	HIS	-	EXPRESSION TAG	UNP P40818
A	721	HIS	-	EXPRESSION TAG	UNP P40818
A	722	HIS	-	EXPRESSION TAG	UNP P40818
A	723	HIS	-	EXPRESSION TAG	UNP P40818
A	724	HIS	-	EXPRESSION TAG	UNP P40818
A	725	SER	-	CLONING ARTIFACT	UNP P40818
A	726	SER	-	CLONING ARTIFACT	UNP P40818
A	727	GLY	-	CLONING ARTIFACT	UNP P40818
A	728	LEU	-	CLONING ARTIFACT	UNP P40818
A	729	VAL	-	CLONING ARTIFACT	UNP P40818
A	730	PRO	-	CLONING ARTIFACT	UNP P40818
A	731	ARG	-	CLONING ARTIFACT	UNP P40818
A	732	GLY	-	CLONING ARTIFACT	UNP P40818
A	733	SER	-	CLONING ARTIFACT	UNP P40818
A	788	MSE	MET	MODIFIED RESIDUE	UNP P40818
A	834	MSE	MET	MODIFIED RESIDUE	UNP P40818
A	877	MSE	MET	MODIFIED RESIDUE	UNP P40818
A	950	MSE	MET	MODIFIED RESIDUE	UNP P40818

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	191	Total 191	O 191	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 8



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.17Å 67.17Å 194.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	18.20 – 2.00 18.20 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.3 (18.20-2.00) 99.3 (18.20-2.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.168 , 0.210 0.171 , 0.208	Depositor DCC
$R_{free}$ test set	1684 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.5	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 58.9	EDS
Estimated twinning fraction	0.087 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 33136 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2938	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.73	1/2806 (0.0%)	0.74	1/3777 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	796	CYS	CB-SG	8.12	1.96	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	763	ARG	NE-CZ-NH1	5.35	122.97	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2746	0	2690	52	0
2	A	1	0	0	0	0
3	A	191	0	0	6	0
All	All	2938	0	2690	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 10.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:877:MSE:HE2	1:A:950:MSE:CE	1.58	1.34
1:A:877:MSE:HE2	1:A:950:MSE:HE2	1.27	1.14
1:A:877:MSE:CE	1:A:950:MSE:HE2	1.82	1.09
1:A:877:MSE:HE2	1:A:950:MSE:HE1	1.35	1.08
1:A:877:MSE:CE	1:A:950:MSE:CE	2.33	1.06
1:A:796:CYS:SG	1:A:834:MSE:SE	2.74	0.95
1:A:971:PHE:CZ	1:A:998:ILE:HD11	2.03	0.94
1:A:808:ASN:HD22	1:A:811:GLN:NE2	1.69	0.89
1:A:808:ASN:HD21	1:A:835:LYS:NZ	1.81	0.79
1:A:808:ASN:HD21	1:A:835:LYS:HZ3	1.31	0.76
1:A:839:THR:HG23	1:A:841:GLN:H	1.49	0.76
1:A:932:SER:OG	1:A:945:THR:HG22	1.85	0.75
1:A:763:ARG:HB3	1:A:763:ARG:HH11	1.51	0.74
1:A:965:GLN:HG3	1:A:1040:TYR:OH	1.88	0.74
1:A:808:ASN:HD22	1:A:811:GLN:HE21	1.35	0.74
1:A:945:THR:HB	3:A:84:HOH:O	1.88	0.74
1:A:1030:PHE:O	1:A:1097:LYS:HE2	1.90	0.72
1:A:796:CYS:HB3	3:A:113:HOH:O	1.90	0.70
1:A:971:PHE:CE2	1:A:998:ILE:HD11	2.28	0.67
1:A:877:MSE:SE	1:A:950:MSE:HE2	2.46	0.66
1:A:808:ASN:ND2	1:A:811:GLN:NE2	2.42	0.65
1:A:971:PHE:HZ	1:A:998:ILE:HD11	1.60	0.64
1:A:763:ARG:CB	1:A:763:ARG:HH11	2.10	0.63
1:A:763:ARG:HB3	1:A:763:ARG:NH1	2.15	0.62
1:A:763:ARG:HH11	1:A:763:ARG:CG	2.14	0.60
1:A:848:LYS:HG3	3:A:157:HOH:O	2.02	0.60
1:A:1017:ASP:OD1	1:A:1018:GLY:N	2.35	0.60
1:A:1087:GLU:HG2	3:A:41:HOH:O	2.03	0.59
1:A:1091:ILE:HD11	1:A:1095:SER:HB3	1.86	0.58
1:A:982:ARG:HB2	1:A:992:ARG:O	2.06	0.56
1:A:808:ASN:ND2	1:A:835:LYS:NZ	2.54	0.53
1:A:802:ALA:HA	1:A:834:MSE:HE2	1.92	0.52
1:A:1051:TYR:HB3	1:A:1106:TYR:HB3	1.93	0.50
1:A:927:GLN:HE21	1:A:950:MSE:SE	2.44	0.50
1:A:796:CYS:HA	1:A:834:MSE:SE	2.62	0.49
1:A:808:ASN:ND2	1:A:811:GLN:HE21	2.06	0.49
1:A:950:MSE:HA	1:A:950:MSE:HE3	1.94	0.48
1:A:1097:LYS:HE3	3:A:183:HOH:O	2.12	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:788:MSE:HA	1:A:872:LEU:HD22	1.96	0.48
1:A:860:GLN:NE2	1:A:871:GLU:OE1	2.39	0.47
1:A:796:CYS:CB	3:A:113:HOH:O	2.57	0.47
1:A:950:MSE:CA	1:A:950:MSE:HE3	2.45	0.47
1:A:808:ASN:HD22	1:A:811:GLN:HE22	1.59	0.46
1:A:966:ASP:OD1	1:A:969:ARG:NH2	2.49	0.45
1:A:763:ARG:HG2	1:A:763:ARG:HH11	1.81	0.44
1:A:763:ARG:NH1	1:A:763:ARG:CB	2.77	0.44
1:A:877:MSE:CE	1:A:950:MSE:HE1	2.23	0.43
1:A:784:ASN:HB2	1:A:864:TYR:O	2.19	0.43
1:A:956:LEU:HD13	1:A:1026:THR:HB	2.01	0.42
1:A:811:GLN:HG2	1:A:811:GLN:H	1.47	0.42
1:A:848:LYS:HE2	1:A:852:ILE:HD11	2.02	0.42
1:A:936:CYS:O	1:A:940:HIS:HA	2.20	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/396 (85%)	324 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/347 (87%)	291 (97%)	10 (3%)	45	43

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	763	ARG
1	A	811	GLN
1	A	839	THR
1	A	843	ARG
1	A	899	ASP
1	A	945	THR
1	A	973	LYS
1	A	991	ARG
1	A	998	ILE
1	A	1008	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	806	ASN
1	A	808	ASN
1	A	811	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/396 (84%)	0.15	20 (5%)	25 27	24, 37, 60, 80	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	980	ASN	10.9
1	A	819	LEU	6.0
1	A	979	ASP	5.2
1	A	1016	TYR	5.0
1	A	1019	ARG	5.0
1	A	1018	GLY	4.6
1	A	817	SER	4.2
1	A	898	ASN	4.1
1	A	900	HIS	4.0
1	A	899	ASP	4.0
1	A	978	THR	3.4
1	A	888	ASP	2.9
1	A	1017	ASP	2.9
1	A	763	ARG	2.8
1	A	981	ASN	2.8
1	A	837	LEU	2.7
1	A	762	ILE	2.4
1	A	1056	VAL	2.3
1	A	812	ASP	2.2
1	A	901	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	1200	1/1	0.99	0.03	-8.01	43,43,43,43	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.