wwPDB X-ray Structure Validation Summary Report (i)

Feb 1, 2016 – 08:54 AM GMT

PDB ID	:	3GIN
Title	:	Crystal structure of E454K-CBD1
Authors	:	Chaptal, V.; Mercado-Besserer, G.; Abramson, J.
Deposited on	:	2009-03-05
Resolution	:	2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	% 66% 6% 28%
1	B	160	% 61% 11% 28%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sodium/calcium exchanger 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	116	895	563	145	184	3	0	0	0
1	B	116	896	563	145	185	3	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	350	MET	-	EXPRESSION TAG	UNP P23685
A	351	ALA	-	EXPRESSION TAG	UNP P23685
A	352	HIS	-	EXPRESSION TAG	UNP P23685
A	353	HIS	-	EXPRESSION TAG	UNP P23685
A	354	HIS	-	EXPRESSION TAG	UNP P23685
A	355	HIS	-	EXPRESSION TAG	UNP P23685
A	356	HIS	-	EXPRESSION TAG	UNP P23685
A	357	HIS	-	EXPRESSION TAG	UNP P23685
A	358	SER	-	EXPRESSION TAG	UNP P23685
A	359	ALA	-	EXPRESSION TAG	UNP P23685
A	360	ALA	-	EXPRESSION TAG	UNP P23685
A	361	LEU	-	EXPRESSION TAG	UNP P23685
A	362	GLU	-	EXPRESSION TAG	UNP P23685
A	363	VAL	-	EXPRESSION TAG	UNP P23685
A	364	LEU	-	EXPRESSION TAG	UNP P23685
A	365	PHE	-	EXPRESSION TAG	UNP P23685
A	366	GLN	-	EXPRESSION TAG	UNP P23685
A	367	GLY	-	EXPRESSION TAG	UNP P23685
A	368	PRO	-	EXPRESSION TAG	UNP P23685
A	369	GLY	-	EXPRESSION TAG	UNP P23685
A	454	LYS	GLU	ENGINEERED	UNP P23685
B	350	MET	-	EXPRESSION TAG	UNP P23685
B	351	ALA	-	EXPRESSION TAG	UNP P23685
B	352	HIS	-	EXPRESSION TAG	UNP P23685
B	353	HIS	-	EXPRESSION TAG	UNP P23685

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	354	HIS	-	EXPRESSION TAG	UNP P23685
B	355	HIS	-	EXPRESSION TAG	UNP P23685
B	356	HIS	-	EXPRESSION TAG	UNP P23685
B	357	HIS	-	EXPRESSION TAG	UNP P23685
B	358	SER	-	EXPRESSION TAG	UNP P23685
B	359	ALA	-	EXPRESSION TAG	UNP P23685
B	360	ALA	-	EXPRESSION TAG	UNP P23685
B	361	LEU	-	EXPRESSION TAG	UNP P23685
B	362	GLU	-	EXPRESSION TAG	UNP P23685
B	363	VAL	-	EXPRESSION TAG	UNP P23685
B	364	LEU	-	EXPRESSION TAG	UNP P23685
B	365	PHE	-	EXPRESSION TAG	UNP P23685
B	366	GLN	-	EXPRESSION TAG	UNP P23685
B	367	GLY	-	EXPRESSION TAG	UNP P23685
B	368	PRO	-	EXPRESSION TAG	UNP P23685
B	369	GLY	-	EXPRESSION TAG	UNP P23685
B	454	LYS	GLU	ENGINEERED	UNP P23685

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
2	B	3	3	3	0	0
			Total	Ca		
2	A	4	4	4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	55	55	55	0	0
			Total	O		
3	B	43	43	43	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sodium/calcium exchanger 1

Chain A:



- Molecule 1: Sodium/calcium exchanger 1

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α, β, γ	91.86Å 91.86Å 65.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.96 – 2.40 64.95 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.3 (64.96-2.40) 98.4 (64.95-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.21 (at 2.40Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.259 0.201 , 0.264	Depositor DCC
R_{free} test set	535 reflections (4.99%)	DCC
Wilson B-factor (Å^2)	31.2	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 42.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Outliers	0 of 11268 reflections	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	1896	wwPDB-VP
Average B, all atoms (Å^2)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.71% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.77	0/909	0.76	0/1231
1	B	0.77	0/910	0.76	0/1232
All	All	0.77	0/1819	0.76	0/2463

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	895	0	857	10	0
1	B	896	0	857	13	0
2	A	4	0	0	0	0
2	B	3	0	0	0	0
3	A	55	0	0	0	0
3	B	43	0	0	0	0
All	All	1896	0	1714	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:383:CYS:SG	1:B:444:ILE:HD13	2.19	0.83
1:A:485:CYS:HG	1:B:485:CYS:HG	0.78	0.75
1:B:431:PHE:CE2	1:B:438:LYS:HG2	2.27	0.70
1:A:383:CYS:HB2	1:A:387:CYS:SG	2.39	0.63
1:A:485:CYS:HG	1:B:485:CYS:CB	2.17	0.58

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/160 (70%)	111 (99%)	1 (1%)	0	100	100
1	B	112/160 (70%)	110 (98%)	2 (2%)	0	100	100
All	All	224/320 (70%)	221 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/134 (75%)	99 (99%)	1 (1%)	82	93
1	B	101/134 (75%)	99 (98%)	2 (2%)	63	81

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	201/268 (75%)	198 (98%)	3 (2%)	72 87

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	493	THR
1	B	402	THR
1	B	493	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	382	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	116/160 (72%)	0.08	2 (1%)	73	72	20, 27, 38, 42	0
1	B	116/160 (72%)	0.12	1 (0%)	85	85	18, 29, 38, 43	0
All	All	232/320 (72%)	0.10	3 (1%)	79	79	18, 28, 38, 43	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	483	LEU	3.3
1	A	448	ASP	3.0
1	B	501	HIS	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
2	CA	A	1	1/1	0.94	0.18	-0.45	51,51,51,51	0
2	CA	B	4	1/1	0.92	0.13	-1.21	51,51,51,51	0
2	CA	B	3	1/1	0.88	0.12	-1.35	49,49,49,49	0
2	CA	A	2	1/1	0.94	0.09	-1.56	61,61,61,61	0
2	CA	B	6	1/1	0.77	0.07	-3.62	62,62,62,62	1
2	CA	A	7	1/1	0.88	0.09	-3.67	64,64,64,64	0
2	CA	A	5	1/1	0.97	0.05	-	37,37,37,37	1

6.5 Other polymers (i)

There are no such residues in this entry.