



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:18 AM GMT

PDB ID : 2GJM  
Title : Crystal structure of Buffalo lactoperoxidase at 2.75Å resolution  
Authors : Sheikh, I.A.; Ethayathulla, A.S.; Singh, A.K.; Singh, N.; Sharma, S.; Singh, T.P.  
Deposited on : 2006-03-31  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	583	<div> <div>5%</div> <div>65%</div> <div>30%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	IOD	A	2006	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	IOD	A	2007	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 5088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactoperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	583	Total	C	N	O	S	0	1	0
			4701	2994	835	847	25			

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 4 is a polymer of unknown type called SUGAR (4-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 5 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 6 is CARBONATE ION (three-letter code: CO3) (formula: CO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	1	3		


- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Ca 1 1	0	0

- Molecule 8 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	8	Total I 8 8	0	0

- Molecule 9 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total C Fe N O 43 34 1 4 4	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	183	Total O 183 183	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Lactoperoxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.24Å 80.47Å 77.35Å 90.00° 102.71° 90.00°	Depositor
Resolution (Å)	20.00 – 2.75 25.27 – 2.73	Depositor EDS
% Data completeness (in resolution range)	96.0 (20.00-2.75) 95.4 (25.27-2.73)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.72Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.182 , 0.216 0.184 , 0.216	Depositor DCC
$R_{free}$ test set	798 reflections (5.15%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.8	Xtriage
Anisotropy	0.711	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 63.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 16582 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5088	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, SCN, NAG, CA, BMA, HEM, IOD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/4827	0.72	5/6544 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	153	CYS	C-N-CD	-8.60	101.68	120.60
1	A	355	PRO	CA-N-CD	-7.25	101.35	111.50
1	A	154	PRO	CA-N-CD	-7.25	101.35	111.50
1	A	154	PRO	CA-C-N	-6.28	103.39	117.20
1	A	158	SER	N-CA-C	5.87	126.85	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4701	0	4618	131	0
2	A	56	0	50	0	0
3	A	39	0	34	1	0
4	A	50	0	43	1	0
5	A	3	0	0	0	0
6	A	4	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	1	0	0	0	0
8	A	8	0	0	5	0
9	A	43	0	30	2	0
10	A	183	0	0	10	0
All	All	5088	0	4775	135	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (135) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:211:LEU:HD23	1:A:254:LEU:HD22	1.44	1.00
1:A:109:GLU:HB3	1:A:112:LYS:HE3	1.48	0.94
1:A:153:CYS:HB2	1:A:154:PRO:HD2	1.47	0.93
1:A:153:CYS:CB	1:A:154:PRO:HD2	1.99	0.93
8:A:2005:IOD:I	10:A:2191:HOH:O	2.61	0.88
1:A:50:ARG:HG2	1:A:51:LYS:N	1.89	0.86
1:A:114:GLN:HG3	10:A:2161:HOH:O	1.77	0.83
1:A:17:ARG:HD2	10:A:2170:HOH:O	1.77	0.81
1:A:120:ILE:HB	10:A:2161:HOH:O	1.82	0.80
1:A:153:CYS:CB	1:A:154:PRO:CD	2.62	0.77
1:A:333:ARG:HH11	1:A:421:ASN:ND2	1.85	0.74
1:A:49:GLN:O	1:A:50:ARG:HB3	1.89	0.71
1:A:109:GLU:HG3	1:A:111:SER:H	1.55	0.71
1:A:573:SER:OG	1:A:574:PRO:HD3	1.90	0.71
1:A:452:GLN:NE2	1:A:458:LYS:HG2	2.06	0.70
1:A:154:PRO:HG2	1:A:156(B):TYR:CD1	2.26	0.69
1:A:197:MET:HG2	1:A:257:HIS:CD2	2.28	0.69
1:A:79:ASP:O	1:A:388:LYS:HD3	1.95	0.67
1:A:99:PHE:O	1:A:101:PRO:HD3	1.93	0.67
1:A:521:THR:OG1	1:A:524:GLN:HG3	1.95	0.66
1:A:171:LEU:HD21	1:A:288:ILE:HG22	1.77	0.66
1:A:155:THR:O	1:A:156:PRO:O	2.13	0.66
1:A:2:ASP:HB3	10:A:2162:HOH:O	1.95	0.65
1:A:77:VAL:HG13	1:A:390:LYS:HG3	1.79	0.65
1:A:514:TRP:CE2	1:A:515:GLU:HG3	2.32	0.65
1:A:355:PRO:HB2	1:A:356:GLU:HG3	1.78	0.64
1:A:237:ASP:OD2	1:A:239:ARG:HD3	1.98	0.64
1:A:159:LEU:HD22	1:A:160:ALA:H	1.63	0.63
1:A:318:ASN:C	1:A:318:ASN:HD22	2.02	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:460:LEU:HD21	1:A:482:ALA:HB1	1.83	0.61
1:A:406:LEU:HD22	1:A:417:LEU:HB2	1.82	0.60
1:A:318:ASN:HD22	1:A:319:SER:N	2.00	0.60
1:A:285:GLY:O	1:A:289:GLN:HG3	2.02	0.59
1:A:334:PHE:CB	1:A:481:ASN:HD21	2.17	0.57
1:A:256:GLU:OE2	1:A:259:ARG:NH1	2.38	0.57
3:A:586:NAG:H61	3:A:587:MAN:H2	1.86	0.57
1:A:107:SER:O	1:A:108:ASN:CB	2.54	0.55
1:A:525:ARG:O	1:A:529:GLN:HG3	2.06	0.55
1:A:171:LEU:HD21	1:A:288:ILE:CG2	2.35	0.55
1:A:272:ASN:O	1:A:276:LEU:HD23	2.06	0.55
1:A:152:VAL:O	1:A:153:CYS:C	2.45	0.55
1:A:290:ILE:HD12	1:A:531:PHE:HD1	1.72	0.55
1:A:242:GLU:OE1	1:A:243:GLN:HG2	2.07	0.54
1:A:213:PHE:CD1	1:A:231:PRO:HG2	2.42	0.54
1:A:116:GLU:CD	1:A:411:HIS:HD1	2.10	0.54
1:A:349:ASN:O	1:A:350:TYR:HB2	2.06	0.54
1:A:205:ASP:HB2	1:A:210:TYR:CZ	2.43	0.54
1:A:242:GLU:OE1	1:A:243:GLN:CG	2.56	0.54
1:A:132:LYS:O	1:A:133:ASN:HB2	2.08	0.53
1:A:92:ILE:HD11	1:A:249:ALA:HB1	1.91	0.53
1:A:153:CYS:HB3	1:A:154:PRO:HD2	1.89	0.53
1:A:531:PHE:CE2	1:A:570:LEU:HD22	2.44	0.53
1:A:153:CYS:HB3	1:A:154:PRO:CD	2.38	0.52
1:A:287:PHE:O	1:A:291:ILE:HG12	2.10	0.52
1:A:18:ARG:CB	1:A:18:ARG:HH11	2.22	0.52
1:A:200:ASN:ND2	8:A:2004:IOD:I	3.13	0.51
1:A:2:ASP:O	1:A:4:ASN:N	2.37	0.51
1:A:226:THR:O	1:A:229:ARG:HG3	2.11	0.51
1:A:164:ILE:HG22	1:A:165:ASN:N	2.25	0.51
1:A:163:GLN:HG2	1:A:428:HIS:CE1	2.46	0.51
1:A:40:GLY:HA2	10:A:2132:HOH:O	2.10	0.51
1:A:535:ARG:HD3	1:A:568:ASP:O	2.11	0.50
1:A:184:ALA:O	1:A:188:ARG:HG3	2.11	0.49
1:A:271:TRP:CG	1:A:275:LYS:HG2	2.48	0.49
1:A:378:ASP:N	1:A:379:PRO:CD	2.76	0.49
1:A:528:LEU:HD22	1:A:531:PHE:CZ	2.47	0.48
9:A:605:HEM:HBB2	9:A:605:HEM:HMB1	1.96	0.48
1:A:322:PRO:O	1:A:506:ILE:HG22	2.13	0.48
1:A:109:GLU:CB	1:A:112:LYS:HE3	2.32	0.47
1:A:360:PRO:HB2	1:A:363:THR:HG23	1.95	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:531:PHE:HE2	1:A:570:LEU:HD22	1.79	0.47
1:A:63:GLU:OE2	1:A:67:LYS:NZ	2.44	0.47
1:A:181:PRO:HD2	8:A:2002:IOD:I	2.84	0.47
1:A:404:ASN:O	1:A:414:GLY:HA2	2.15	0.47
1:A:290:ILE:HD12	1:A:531:PHE:CD1	2.50	0.46
1:A:74:GLU:HG2	1:A:341:SER:OG	2.15	0.46
1:A:260:LEU:O	1:A:264:LEU:HG	2.16	0.46
1:A:48:THR:CG2	1:A:50:ARG:HD3	2.45	0.46
1:A:528:LEU:HD22	1:A:531:PHE:CE1	2.51	0.46
1:A:535:ARG:HD3	1:A:569:LYS:HA	1.97	0.46
1:A:566:THR:O	1:A:566:THR:CG2	2.64	0.46
1:A:155:THR:C	1:A:156:PRO:O	2.54	0.46
1:A:107:SER:O	1:A:108:ASN:HB3	2.15	0.46
1:A:370:ARG:O	1:A:374:ASP:HB3	2.16	0.46
1:A:405:LYS:HA	1:A:413(A):HIS:O	2.15	0.45
1:A:491:VAL:HB	1:A:495:LEU:HB3	1.98	0.45
1:A:215:ASN:N	10:A:2140:HOH:O	2.43	0.45
1:A:308:LYS:HE3	1:A:309:TRP:CZ2	2.50	0.45
1:A:210:TYR:OH	1:A:376:GLY:HA2	2.17	0.45
1:A:310:ILE:HG12	1:A:504:GLN:HB2	1.99	0.45
1:A:543:ILE:HG23	8:A:2006:IOD:I	2.87	0.45
1:A:334:PHE:HA	1:A:481:ASN:HD21	1.82	0.45
1:A:345:ARG:NH2	1:A:374:ASP:OD2	2.49	0.44
1:A:112:LYS:HB2	1:A:112:LYS:NZ	2.32	0.44
1:A:524:GLN:O	1:A:528:LEU:HG	2.17	0.44
1:A:422:LEU:HD21	1:A:478:ILE:HB	1.98	0.44
1:A:32:TRP:CE2	1:A:325:SER:HB3	2.52	0.44
1:A:215:ASN:HB3	10:A:2140:HOH:O	2.17	0.44
1:A:223:PHE:CZ	1:A:412:LYS:HB3	2.53	0.44
1:A:279:GLU:O	1:A:283:ILE:HG13	2.18	0.43
1:A:125:CYS:SG	1:A:127:PRO:HG3	2.58	0.43
1:A:105:LEU:N	1:A:105:LEU:HD12	2.33	0.43
1:A:5:SER:HA	1:A:6:PRO:HD3	1.87	0.43
1:A:37:TYR:CZ	1:A:161:ARG:HB3	2.53	0.43
1:A:334:PHE:HB2	1:A:481:ASN:HD21	1.82	0.43
1:A:297:LEU:HB2	1:A:298:PRO:HD3	2.00	0.43
1:A:462:LYS:HE2	1:A:462:LYS:HB2	1.82	0.42
1:A:92:ILE:HD11	1:A:249:ALA:CB	2.49	0.42
1:A:130:PHE:HE2	1:A:144:MET:CE	2.33	0.42
1:A:62:ARG:O	1:A:65:SER:HB3	2.19	0.42
1:A:579:GLU:H	1:A:579:GLU:HG2	1.46	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:LEU:HD12	1:A:572:LEU:HD11	2.01	0.42
1:A:292:THR:HA	1:A:296:TYR:HB3	2.00	0.42
4:A:589:NAG:H62	4:A:590:BMA:C1	2.50	0.42
1:A:333:ARG:NH1	1:A:421:ASN:ND2	2.61	0.42
1:A:120:ILE:CB	10:A:2161:HOH:O	2.57	0.42
1:A:392:MET:HB3	1:A:485:MET:CE	2.49	0.42
1:A:226:THR:O	1:A:229:ARG:NH2	2.52	0.41
1:A:330:PHE:HE2	1:A:424:ARG:HB3	1.85	0.41
1:A:543:ILE:HA	8:A:2006:IOD:I	2.91	0.41
1:A:351:GLN:O	1:A:352:PRO:C	2.58	0.41
1:A:52:THR:HB	1:A:56:PHE:N	2.36	0.41
1:A:211:LEU:HD11	1:A:250:HIS:HB3	2.03	0.41
1:A:137:LEU:HA	1:A:141:GLY:O	2.19	0.41
1:A:579:GLU:HB2	1:A:580:ASN:H	1.45	0.41
1:A:476:ILE:HG23	1:A:477:TRP:N	2.36	0.41
1:A:151:PHE:CD1	1:A:151:PHE:N	2.88	0.41
1:A:452:GLN:O	1:A:456:LYS:N	2.54	0.41
1:A:115:CYS:HB2	10:A:2015:HOH:O	2.21	0.41
1:A:333:ARG:NH1	9:A:605:HEM:HBA2	2.36	0.40
1:A:284:LEU:O	1:A:287:PHE:HB3	2.21	0.40
1:A:511:ARG:O	1:A:516:ASN:ND2	2.48	0.40
1:A:361:LEU:HD13	1:A:365:PHE:CZ	2.56	0.40
1:A:252:LEU:HD12	1:A:252:LEU:HA	1.94	0.40
1:A:557:TYR:CD1	1:A:558:PRO:HA	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	582/583 (100%)	528 (91%)	40 (7%)	14 (2%)	<b>7</b> <b>22</b>

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	107	SER
1	A	108	ASN
1	A	152	VAL
1	A	153	CYS
1	A	155	THR
1	A	156	PRO
1	A	158	SER
1	A	579	GLU
1	A	50	ARG
1	A	154	PRO
1	A	156(B)	TYR
1	A	157	GLN
1	A	156(A)	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	510/509 (100%)	467 (92%)	43 (8%)	14	34

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ARG
1	A	43	LEU
1	A	50	ARG
1	A	77	VAL
1	A	84	LEU
1	A	104	GLU
1	A	110	HIS
1	A	142	LYS
1	A	152	VAL
1	A	155	THR
1	A	156(A)	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	159	LEU
1	A	171	LEU
1	A	186	ARG
1	A	187	LEU
1	A	200	ASN
1	A	226	THR
1	A	239	ARG
1	A	245	LEU
1	A	252	LEU
1	A	259	ARG
1	A	260	LEU
1	A	301	LEU
1	A	307	GLN
1	A	318	ASN
1	A	332	PHE
1	A	345	ARG
1	A	361	LEU
1	A	373	LYS
1	A	403	ARG
1	A	435	SER
1	A	446	LYS
1	A	448	LEU
1	A	464	LEU
1	A	478	ILE
1	A	488	ARG
1	A	495	LEU
1	A	504	GLN
1	A	507	ARG
1	A	526	ASP
1	A	549	LEU
1	A	566	THR
1	A	579	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	ASN
1	A	200	ASN
1	A	307	GLN
1	A	314	GLN
1	A	318	ASN
1	A	349	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	421	ASN
1	A	452	GLN
1	A	481	ASN
1	A	504	GLN
1	A	529	GLN
1	A	553	GLN
1	A	555	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

11 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	581	1,2	14,14,15	0.49	0	15,19,21	1.70	1 (6%)
2	NAG	A	582	2	14,14,15	0.56	0	15,19,21	1.37	1 (6%)
2	NAG	A	583	1,2	14,14,15	0.48	0	15,19,21	0.96	0
2	NAG	A	584	2	14,14,15	0.63	0	15,19,21	1.07	2 (13%)
3	NAG	A	585	1,3	14,14,15	0.46	0	15,19,21	0.83	1 (6%)
3	NAG	A	586	3	14,14,15	0.64	0	15,19,21	0.86	1 (6%)
3	MAN	A	587	3	11,11,12	0.54	0	14,15,17	0.42	0
4	NAG	A	588	1,4	14,14,15	0.72	0	15,19,21	1.38	1 (6%)
4	NAG	A	589	4	14,14,15	0.70	0	15,19,21	0.77	0
4	BMA	A	590	4	11,11,12	0.95	1 (9%)	14,15,17	2.10	3 (21%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MAN	A	591	4	11,11,12	1.03	0	14,15,17	1.13	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	581	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	582	2	-	0/6/23/26	0/1/1/1
2	NAG	A	583	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	584	2	-	0/6/23/26	0/1/1/1
3	NAG	A	585	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	586	3	-	0/6/23/26	0/1/1/1
3	MAN	A	587	3	-	0/2/19/22	0/1/1/1
4	NAG	A	588	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	589	4	-	0/6/23/26	0/1/1/1
4	BMA	A	590	4	-	0/2/19/22	0/1/1/1
4	MAN	A	591	4	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	590	BMA	O5-C5	2.07	1.48	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	581	NAG	C2-N2-C7	-5.68	115.74	123.04
2	A	582	NAG	C2-N2-C7	-4.83	116.83	123.04
4	A	591	MAN	C1-C2-C3	-2.69	106.36	109.54
4	A	591	MAN	C1-O5-C5	-2.43	109.16	112.25
3	A	586	NAG	C2-N2-C7	-2.35	120.01	123.04
3	A	585	NAG	C2-N2-C7	-2.05	120.41	123.04
2	A	584	NAG	C2-N2-C7	-2.04	120.42	123.04
4	A	590	BMA	C1-C2-C3	2.46	112.45	109.54
2	A	584	NAG	C4-C3-C2	2.46	115.05	111.23
4	A	590	BMA	O5-C1-C2	3.23	116.10	110.86
4	A	588	NAG	C4-C3-C2	4.03	117.49	111.23
4	A	590	BMA	C1-O5-C5	6.29	120.22	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	586	NAG	1	0
3	A	587	MAN	1	0
4	A	589	NAG	1	0
4	A	590	BMA	1	0

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	CO3	A	1001	-	0,3,3	0.00	-	0,3,3	0.00	-
5	SCN	A	1502	-	2,2,2	1.89	1 (50%)	1,1,1	0.89	0
9	HEM	A	605	1	30,50,50	2.71	9 (30%)	24,82,82	2.99	10 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CO3	A	1001	-	-	0/0/0/0	0/0/0/0
5	SCN	A	1502	-	-	0/0/0/0	0/0/0/0
9	HEM	A	605	1	-	0/10/54/54	0/0/8/8

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	605	HEM	C3B-C4B	-6.61	1.46	1.51
9	A	605	HEM	C3D-C4D	-5.35	1.44	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	605	HEM	C2C-C1C	-3.89	1.45	1.52
9	A	605	HEM	C1C-NC	2.48	1.39	1.36
9	A	605	HEM	C3C-CAC	2.48	1.56	1.51
5	A	1502	SCN	C-S	2.55	1.80	1.63
9	A	605	HEM	CAA-C2A	2.61	1.56	1.52
9	A	605	HEM	C3B-CAB	4.28	1.59	1.51
9	A	605	HEM	FE-ND	4.57	2.21	1.97
9	A	605	HEM	FE-NC	6.84	2.22	1.95

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	605	HEM	CBA-CAA-C2A	-3.24	106.72	112.53
9	A	605	HEM	CAA-C2A-C1A	-2.67	124.11	127.01
9	A	605	HEM	C2D-C3D-C4D	2.26	105.34	101.50
9	A	605	HEM	C3B-C4B-CHC	2.52	126.71	123.16
9	A	605	HEM	CMB-C2B-C3B	4.11	126.79	116.53
9	A	605	HEM	CAD-C3D-C2D	4.15	125.15	113.22
9	A	605	HEM	CAD-C3D-C4D	4.83	129.51	112.47
9	A	605	HEM	CBD-CAD-C3D	5.67	130.05	113.55
9	A	605	HEM	CAD-CBD-CGD	6.11	137.94	113.02
9	A	605	HEM	CMC-C2C-C3C	6.14	131.85	116.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	605	HEM	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	583/583 (100%)	-0.12	31 (5%) 30 23	8, 26, 52, 85	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	578	ARG	8.0
1	A	108	ASN	7.9
1	A	109	GLU	6.1
1	A	158	SER	6.0
1	A	157	GLN	5.8
1	A	579	GLU	5.8
1	A	111	SER	5.6
1	A	107	SER	5.4
1	A	156(B)	TYR	5.2
1	A	156	PRO	5.0
1	A	106	GLY	4.8
1	A	105	LEU	4.8
1	A	580	ASN	4.1
1	A	218	PRO	4.1
1	A	110	HIS	4.0
1	A	112	LYS	3.9
1	A	113	THR	3.2
1	A	355	PRO	3.2
1	A	156(A)	PRO	2.9
1	A	2	ASP	2.8
1	A	3	GLU	2.4
1	A	114	GLN	2.4
1	A	559	HIS	2.4
1	A	155	THR	2.3
1	A	270	HIS	2.3
1	A	307	GLN	2.2
1	A	4	ASN	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	120	ILE	2.2
1	A	215	ASN	2.1
1	A	217	LYS	2.1
1	A	152	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	583	14/15	0.91	0.20	0.67	45,49,51,53	0
3	NAG	A	585	14/15	0.94	0.17	0.19	39,40,44,45	0
2	NAG	A	581	14/15	0.87	0.17	-0.12	49,51,55,60	0
3	MAN	A	587	11/12	0.61	0.34	-	46,47,48,48	11
4	NAG	A	588	14/15	0.84	0.27	-	45,49,51,51	0
4	MAN	A	591	11/12	0.59	0.55	-	48,49,49,49	11
3	NAG	A	586	14/15	0.82	0.35	-	47,50,51,51	0
2	NAG	A	582	14/15	0.86	0.43	-	63,66,67,67	0
4	BMA	A	590	11/12	0.66	0.49	-	49,49,49,49	11
4	NAG	A	589	14/15	0.67	0.44	-	48,49,50,50	14
2	NAG	A	584	14/15	0.80	0.48	-	56,57,58,58	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
8	IOD	A	2007	1/1	0.97	0.13	2.12	10,10,10,10	1
9	HEM	A	605	43/43	0.95	0.14	0.01	13,19,23,25	0
7	CA	A	1503	1/1	0.99	0.10	-0.95	14,14,14,14	0
8	IOD	A	2004	1/1	0.97	0.12	-1.03	33,33,33,33	1
5	SCN	A	1502	3/3	0.94	0.12	-1.79	14,14,14,15	0
8	IOD	A	2001	1/1	0.97	0.06	-2.18	47,47,47,47	1
8	IOD	A	2006	1/1	0.93	0.12	-2.30	37,37,37,37	1
8	IOD	A	2005	1/1	0.92	0.10	-3.55	37,37,37,37	1
8	IOD	A	2002	1/1	0.99	0.05	-4.94	42,42,42,42	1
8	IOD	A	2008	1/1	0.92	0.12	-	24,24,24,24	1
6	CO3	A	1001	4/4	0.95	0.20	-	31,32,33,33	0
8	IOD	A	2003	1/1	0.97	0.07	-	47,47,47,47	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.