



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2016 – 09:10 PM EDT

PDB ID : 5GLH
Title : Human endothelin receptor type-B in complex with ET-1
Authors : Shihoya, W.; Nishizawa, T.; Okuta, A.; Tani, K.; Fujiyoshi, Y.; Dohmae, N.;
Nureki, O.; Doi, T.
Deposited on : 2016-07-11
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	
2	B	21	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endothelin Receptor Subtype-B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	452	Total	C	N	O	S	0	0	0
			3468	2263	577	607	21			


- Molecule 2 is a protein called Peptide from Endothelin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	21	Total	C	N	O	S	0	0	0
			171	109	25	32	5			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	O	0	0
			4	4		

- Molecule 1: Endothelin Receptor Subtype-B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	72.97Å 172.97Å 109.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.21 – 2.80 46.21 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.21-2.80) 87.0 (46.21-2.60)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.61Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.234 , 0.277 0.229 , 0.275	Depositor DCC
R_{free} test set	1608 reflections (10.12%)	DCC
Wilson B-factor (Å ²)	56.9	Xtriage
Anisotropy	0.700	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 63.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3643	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3541	0.39	0/4828
2	B	0.23	0/175	0.43	0/234
All	All	0.24	0/3716	0.39	0/5062

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3468	0	3475	41	0
2	B	171	0	157	8	0
3	A	4	0	0	0	0
All	All	3643	0	3632	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:PRO:HG3	2:B:20:ILE:HD11	1.73	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:289:THR:HG21	1:A:329:VAL:HG21	1.78	0.66
1:A:286:LEU:HD21	1:A:333:ALA:HB2	1.79	0.63
1:A:1010:ASP:OD1	1:A:1148:ARG:NH2	2.34	0.61
1:A:1005:GLU:OE2	1:A:1008:ARG:NH1	2.34	0.61
1:A:254:ILE:HD12	2:B:16:HIS:HB2	1.83	0.60
1:A:1029:ILE:HD13	1:A:1067:PHE:HB2	1.83	0.60
1:A:1054:THR:OG1	1:A:1057:VAL:O	2.19	0.59
1:A:346:LYS:NZ	2:B:2:SER:O	2.36	0.59
1:A:257:LEU:HB3	2:B:2:SER:HB2	1.84	0.57
1:A:90:CYS:SG	1:A:91:GLN:N	2.81	0.54
1:A:182:LYS:NZ	2:B:21:TRP:OXT	2.38	0.52
1:A:1155:THR:HG23	1:A:1157:THR:H	1.76	0.51
1:A:390:SER:O	1:A:392:ARG:N	2.41	0.50
1:A:273:LYS:NZ	2:B:21:TRP:OXT	2.45	0.49
1:A:342:ALA:HB2	1:A:367:LEU:HG	1.93	0.48
1:A:247:TYR:HD1	2:B:9:LYS:HB3	1.78	0.48
1:A:263:THR:HG22	1:A:265:PHE:H	1.79	0.48
1:A:1105:GLN:OE1	1:A:1138:TRP:NE1	2.45	0.47
1:A:387:TYR:CE1	1:A:394:LYS:HG3	2.49	0.47
1:A:175:LYS:NZ	1:A:240:PHE:O	2.47	0.47
1:A:154:ALA:O	1:A:158:ASN:ND2	2.45	0.46
1:A:1033:LEU:HD12	1:A:1034:THR:HG22	1.97	0.45
1:A:1011:GLU:OE1	1:A:1145:ARG:NH2	2.48	0.44
1:A:93:PRO:HG3	1:A:358:CYS:SG	2.57	0.44
1:A:356:ASN:HD22	1:A:356:ASN:N	2.15	0.44
1:A:1065:LYS:O	1:A:1069:GLN:HG2	2.17	0.44
1:A:390:SER:HB3	1:A:393:PHE:HB2	2.00	0.43
1:A:1137:ARG:HD2	1:A:1138:TRP:N	2.33	0.43
1:A:126:ILE:HG21	1:A:141:ALA:HB2	1.99	0.43
1:A:321:VAL:O	1:A:325:VAL:HB	2.18	0.43
1:A:136:PRO:HG3	1:A:199:ARG:HH12	1.85	0.42
1:A:147:ASP:O	1:A:151:ILE:HG12	2.20	0.41
1:A:1135:LYS:HB3	1:A:1135:LYS:HE2	1.85	0.41
1:A:302:ARG:HA	1:A:1003:ILE:HG22	2.01	0.41
1:A:247:TYR:CD1	2:B:9:LYS:HB3	2.55	0.41
1:A:1083:LYS:HD2	1:A:1112:ALA:HB1	2.02	0.41
1:A:340:HIS:O	1:A:344:ILE:HG12	2.21	0.40
1:A:1047:ASP:OD2	1:A:1053:ASN:ND2	2.55	0.40
1:A:175:LYS:HE3	1:A:237:ALA:O	2.20	0.40
1:A:1033:LEU:HG	1:A:1033:LEU:H	1.71	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/498 (90%)	428 (96%)	18 (4%)	0	100	100
2	B	19/21 (90%)	19 (100%)	0	0	100	100
All	All	465/519 (90%)	447 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/425 (85%)	341 (95%)	19 (5%)	28	61
2	B	21/21 (100%)	20 (95%)	1 (5%)	31	66
All	All	381/446 (85%)	361 (95%)	20 (5%)	29	62

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	241	ASP
1	A	302	ARG
1	A	1011	GLU
1	A	1033	LEU
1	A	1040	ASN
1	A	1048	LYS
1	A	1055	ASN
1	A	1059	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1062	GLU
1	A	1080	ARG
1	A	1092	ASP
1	A	1094	VAL
1	A	1104	PHE
1	A	1137	ARG
1	A	325	VAL
1	A	356	ASN
1	A	387	TYR
1	A	395	ASN
1	A	397	PHE
2	B	1	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1055	ASN
1	A	356	ASN
1	A	378	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/498 (90%)	0.40	33 (7%) 18 10	47, 80, 144, 162	0
2	B	21/21 (100%)	0.01	0 100 100	63, 73, 79, 91	0
All	All	473/519 (91%)	0.38	33 (6%) 19 11	47, 79, 142, 162	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1021	THR	10.0
1	A	1038	SER	9.5
1	A	390	SER	8.1
1	A	1032	LEU	4.6
1	A	395	ASN	4.5
1	A	1034	THR	3.8
1	A	391	LYS	3.7
1	A	389	VAL	3.6
1	A	387	TYR	3.5
1	A	1037	PRO	3.5
1	A	397	PHE	3.3
1	A	1059	THR	3.3
1	A	388	LEU	3.2
1	A	93	PRO	3.2
1	A	1036	SER	3.0
1	A	99	THR	2.9
1	A	353	ASN	2.8
1	A	352	GLN	2.8
1	A	1022	GLU	2.7
1	A	89	PRO	2.7
1	A	102	TYR	2.7
1	A	1033	LEU	2.6
1	A	393	PHE	2.6
1	A	385	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	383	PRO	2.5
1	A	1019	LYS	2.4
1	A	123	LEU	2.4
1	A	386	LEU	2.4
1	A	1054	THR	2.4
1	A	94	ILE	2.3
1	A	384	ILE	2.1
1	A	1051	GLY	2.0
1	A	1041	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.