



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:39 PM GMT

PDB ID : 1GN4
Title : H145E MUTANT OF MYCOBACTERIUM TUBERCULOSIS IRON-SUPEROXIDE DISMUTASE.
Authors : Bunting, K.A.; Cooper, J.B.; Badasso, M.O.; Tickle, I.J.; Newton, M.; Wood, S.P.; Zhang, Y.; Young, D.B.
Deposited on : 2001-10-02
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	<div> <div></div> <div>40%</div> <div>43%</div> <div>12%</div> <div>• •</div> </div>
1	B	207	<div> <div></div> <div>32%</div> <div>52%</div> <div>9%</div> <div>• •</div> </div>
1	C	207	<div> <div></div> <div>38%</div> <div>41%</div> <div>16%</div> <div>• •</div> </div>
1	D	207	<div> <div></div> <div>40%</div> <div>43%</div> <div>11%</div> <div>• •</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6560 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUPEROXIDE DISMUTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	0	0
			1567	1011	267	288	1			
1	B	198	Total	C	N	O	S	0	0	0
			1567	1011	267	288	1			
1	C	198	Total	C	N	O	S	0	0	0
			1567	1011	267	288	1			
1	D	198	Total	C	N	O	S	0	0	0
			1567	1011	267	288	1			

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		
2	D	1	Total	Mn	0	0
			1	1		
2	C	1	Total	Mn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	70	Total	O	0	0
			70	70		
3	B	81	Total	O	0	0
			81	81		
3	C	69	Total	O	0	0
			69	69		
3	D	68	Total	O	0	0
			68	68		



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	107.66Å 102.86Å 73.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 12.54 – 2.50	Depositor EDS
% Data completeness (in resolution range)	94.4 (20.00-2.50) 95.0 (12.54-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 2.50Å)	Xtriage
Refinement program	CCP4	Depositor
R, R_{free}	0.172 , 0.225 0.153 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.434	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 86.5	EDS
Estimated twinning fraction	0.024 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 27288 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6560	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.75	44/1615 (2.7%)	2.24	87/2203 (3.9%)
1	B	1.76	53/1615 (3.3%)	2.24	95/2203 (4.3%)
1	C	1.75	50/1615 (3.1%)	2.25	98/2203 (4.4%)
1	D	1.62	39/1615 (2.4%)	2.19	84/2203 (3.8%)
All	All	1.72	186/6460 (2.9%)	2.23	364/8812 (4.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	0
1	C	1	0
1	D	1	0
All	All	3	0

The worst 5 of 186 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	117	ALA	CA-CB	10.24	1.74	1.52
1	C	177	PHE	N-CA	10.22	1.66	1.46
1	A	175	VAL	N-CA	9.98	1.66	1.46
1	C	150	PRO	CA-CB	9.91	1.73	1.53
1	A	114	HIS	N-CA	9.87	1.66	1.46

The worst 5 of 364 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	40	ALA	N-CA-CB	-12.73	92.28	110.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	THR	CA-CB-CG2	-11.59	96.18	112.40
1	C	153	ILE	CA-CB-CG2	10.55	131.99	110.90
1	D	117	ALA	N-CA-CB	-10.32	95.66	110.10
1	B	67	ALA	N-CA-CB	-10.23	95.78	110.10

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	20	ILE	CB
1	C	153	ILE	CB
1	D	161	MET	CA

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1567	0	1488	44	0
1	B	1567	0	1488	44	0
1	C	1567	0	1488	53	0
1	D	1567	0	1488	46	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	70	0	0	1	0
3	B	81	0	0	0	0
3	C	69	0	0	5	0
3	D	68	0	0	3	0
All	All	6560	0	5952	180	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 180 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:88:GLY:O	1:C:89:ASP:HB2	1.39	1.07
1:B:95:LEU:HD21	1:B:190:VAL:HG12	1.41	1.02
1:A:88:GLY:O	1:A:89:ASP:HB2	1.66	0.92
1:C:88:GLY:O	1:C:89:ASP:CB	2.22	0.86
1:B:95:LEU:CD2	1:B:190:VAL:HG12	2.06	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/207 (95%)	177 (90%)	17 (9%)	2 (1%)	19	34
1	B	196/207 (95%)	184 (94%)	10 (5%)	2 (1%)	19	34
1	C	196/207 (95%)	175 (89%)	20 (10%)	1 (0%)	34	55
1	D	196/207 (95%)	182 (93%)	12 (6%)	2 (1%)	19	34
All	All	784/828 (95%)	718 (92%)	59 (8%)	7 (1%)	21	37

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	89	ASP
1	A	89	ASP
1	B	148	ASN
1	D	148	ASN
1	B	166	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/163 (96%)	140 (90%)	16 (10%)	9	17
1	B	156/163 (96%)	141 (90%)	15 (10%)	10	19
1	C	156/163 (96%)	142 (91%)	14 (9%)	12	22
1	D	156/163 (96%)	141 (90%)	15 (10%)	10	19
All	All	624/652 (96%)	564 (90%)	60 (10%)	10	19

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	105	SER
1	C	30	SER
1	D	146	GLN
1	B	199	SER
1	C	64	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	114	HIS
1	C	62	ASN
1	D	62	ASN
1	B	82	ASN
1	D	19	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/207 (95%)	-1.04	0 100 100	4, 20, 49, 89	0
1	B	198/207 (95%)	-0.88	0 100 100	7, 27, 61, 86	0
1	C	198/207 (95%)	-1.00	0 100 100	5, 20, 53, 84	0
1	D	198/207 (95%)	-0.95	0 100 100	7, 23, 57, 110	0
All	All	792/828 (95%)	-0.97	0 100 100	4, 22, 56, 110	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MN	A	1200	1/1	0.99	0.07	-0.75	13,13,13,13	0
2	MN	C	1200	1/1	0.99	0.06	-1.48	18,18,18,18	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MN	D	1200	1/1	0.99	0.06	-2.80	18,18,18,18	0
2	MN	B	1200	1/1	0.98	0.04	-6.17	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.