



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:56 PM GMT

PDB ID : 4GQ2
Title : S. pombe Nup120-Nup37 complex
Authors : Liu, X.; Mitchell, J.; Wozniak, R.; Blobel, G.; Fan, J.
Deposited on : 2012-08-22
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	M	950	<div> <div>5%</div> <div>80%</div> <div>13%</div> <div>• 5%</div> </div>
2	P	393	<div> <div>10%</div> <div>74%</div> <div>17%</div> <div>• 6%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoporin nup120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	M	904	Total	C	N	O	S	0	0	0
			7274	4701	1149	1397	27			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	0	ARG	-	EXPRESSION TAG	UNP O43044

- Molecule 2 is a protein called Nup37.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	371	Total	C	N	O	S	0	0	0
			2840	1794	484	546	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	-1	GLY	-	EXPRESSION TAG	UNP O36030
P	0	SER	-	EXPRESSION TAG	UNP O36030

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	M	97	Total	O	0	0
			97	97		
3	P	25	Total	O	0	0
			25	25		

L385	
M386	
G387	
PHE	
THR	
ARG	
LEU	

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.47Å 123.02Å 172.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.40 34.72 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (35.00-2.40) 99.8 (34.72-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.210 , 0.258 0.210 , 0.258	Depositor DCC
R_{free} test set	2996 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	49.7	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 59341 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10236	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	M	0.47	7/7437 (0.1%)	0.60	3/10091 (0.0%)
2	P	0.52	3/2906 (0.1%)	0.64	1/3971 (0.0%)
All	All	0.48	10/10343 (0.1%)	0.61	4/14062 (0.0%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	M	321	PHE	C-N	6.36	1.48	1.34
1	M	728	TRP	CD2-CE2	5.61	1.48	1.41
2	P	260	TRP	CD2-CE2	5.45	1.47	1.41
1	M	616	TRP	CD2-CE2	5.28	1.47	1.41
1	M	75	TRP	CD2-CE2	5.25	1.47	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	321	PHE	O-C-N	16.48	149.07	122.70
1	M	321	PHE	CA-C-N	-12.35	90.04	117.20
1	M	321	PHE	C-N-CA	-8.74	99.86	121.70
2	P	345	TYR	N-CA-C	8.07	132.80	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	7274	0	7170	73	0
2	P	2840	0	2773	49	0
3	M	97	0	0	2	0
3	P	25	0	0	0	0
All	All	10236	0	9943	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:564:GLU:H	1:M:565:THR:HA	1.14	1.08
1:M:564:GLU:N	1:M:565:THR:HA	1.71	1.03
1:M:634:THR:O	1:M:638:THR:HG23	1.75	0.87
1:M:31:ASN:H	1:M:163:ARG:HH22	1.32	0.78
1:M:843:TYR:OH	1:M:901:HIS:HE1	1.68	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	M	888/950 (94%)	840 (95%)	45 (5%)	3 (0%)	46 63
2	P	367/393 (93%)	338 (92%)	25 (7%)	4 (1%)	17 25
All	All	1255/1343 (93%)	1178 (94%)	70 (6%)	7 (1%)	30 43

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	338	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	M	294	ASN
2	P	244	ASN
2	P	250	GLY
1	M	608	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	831/876 (95%)	781 (94%)	50 (6%)	24	37
2	P	321/344 (93%)	299 (93%)	22 (7%)	19	30
All	All	1152/1220 (94%)	1080 (94%)	72 (6%)	22	35

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	541	PHE
1	M	805	ASP
2	P	302	ASN
1	M	556	LEU
1	M	699	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	537	ASN
1	M	705	GLN
2	P	286	ASN
1	M	444	GLN
2	P	118	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	M	904/950 (95%)	0.02	46 (5%) 32 32	26, 56, 112, 159	0
2	P	371/393 (94%)	0.36	41 (11%) 7 7	30, 56, 128, 184	0
All	All	1275/1343 (94%)	0.12	87 (6%) 20 20	26, 56, 118, 184	0

The worst 5 of 87 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	321	PHE	8.6
2	P	54	THR	8.4
2	P	224	SER	7.1
2	P	226	THR	6.6
2	P	220	ALA	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.