



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 11:21 AM EST

PDB ID : 5GQX
Title : Crystal structure of branching enzyme W610N mutant from *Cyanothece* sp. ATCC 51142 in complex with maltoheptaose
Authors : Suzuki, R.; Suzuki, E.
Deposited on : 2016-08-08
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442

i


X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	793	 <div> <div></div> <div>2%</div> <div>87%</div> <div>8%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	801	-	-	-	X
2	GLC	A	804	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	807	-	-	-	X
2	GLC	A	810	-	-	-	X
2	GLC	A	813	-	-	-	X
2	GLC	A	816	-	-	-	X
2	GLC	A	826	-	-	-	X
2	GLC	A	837	-	-	-	X
2	GLC	A	838	-	-	-	X
3	GOL	A	840	-	-	-	X
3	GOL	A	841	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7245 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 1,4-alpha-glucan branching enzyme GlgB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	755	Total	C	N	O	S	0	0	0
			6267	4050	1042	1151	24			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP B1WPM8
A	-18	GLY	-	expression tag	UNP B1WPM8
A	-17	SER	-	expression tag	UNP B1WPM8
A	-16	SER	-	expression tag	UNP B1WPM8
A	-15	HIS	-	expression tag	UNP B1WPM8
A	-14	HIS	-	expression tag	UNP B1WPM8
A	-13	HIS	-	expression tag	UNP B1WPM8
A	-12	HIS	-	expression tag	UNP B1WPM8
A	-11	HIS	-	expression tag	UNP B1WPM8
A	-10	HIS	-	expression tag	UNP B1WPM8
A	-9	SER	-	expression tag	UNP B1WPM8
A	-8	SER	-	expression tag	UNP B1WPM8
A	-7	GLY	-	expression tag	UNP B1WPM8
A	-6	LEU	-	expression tag	UNP B1WPM8
A	-5	VAL	-	expression tag	UNP B1WPM8
A	-4	PRO	-	expression tag	UNP B1WPM8
A	-3	ARG	-	expression tag	UNP B1WPM8
A	-2	GLY	-	expression tag	UNP B1WPM8
A	-1	SER	-	expression tag	UNP B1WPM8
A	0	HIS	-	expression tag	UNP B1WPM8
A	610	ASN	TRP	engineered mutation	UNP B1WPM8

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	530	Total 530	O 530	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1,4-alpha-glucan branching enzyme GlgB



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.03Å 134.03Å 185.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.38 – 2.30 37.58 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.38-2.30) 99.9 (37.58-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	14.37 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.149 , 0.184 0.161 , 0.191	Depositor DCC
R_{free} test set	3785 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7245	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	15/6485 (0.2%)	0.99	15/8824 (0.2%)

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	203	TYR	CE1-CZ	9.18	1.50	1.38
1	A	525	ASP	CB-CG	-7.13	1.36	1.51
1	A	187	SER	CB-OG	-6.75	1.33	1.42
1	A	460	GLU	CD-OE1	6.20	1.32	1.25
1	A	135	GLU	CD-OE2	6.12	1.32	1.25

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	525	ASP	CB-CG-OD1	-7.37	111.66	118.30
1	A	412	ARG	NE-CZ-NH2	-7.29	116.66	120.30
1	A	242	ARG	NE-CZ-NH2	7.03	123.81	120.30
1	A	59	ARG	NE-CZ-NH2	-6.89	116.85	120.30
1	A	203	TYR	CB-CA-C	-6.70	96.99	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6267	0	5853	20	0
2	A	435	0	371	12	0
3	A	12	0	16	1	0
4	A	1	0	0	0	0
5	A	530	0	0	1	0
All	All	7245	0	6240	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:490:THR:O	1:A:491:GLU:HB2	1.95	0.65
1:A:249:GLN:HB3	1:A:250:PRO:HD2	1.78	0.64
2:A:820:GLC:H62	2:A:821:GLC:C5	2.34	0.58
2:A:815:GLC:O5	2:A:821:GLC:O4	2.23	0.57
1:A:490:THR:O	1:A:491:GLU:CB	2.50	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	753/793 (95%)	732 (97%)	20 (3%)	1 (0%)	56 68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	598	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	668 / 700 (95%)	646 (97%)	22 (3%)	45 61

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	240	LYS
1	A	373	ASP
1	A	725	LYS
1	A	307	ASP
1	A	369	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	35	ASN
1	A	65	ASN
1	A	619	GLN
1	A	634	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 42 ligands modelled in this entry, 1 is monoatomic - leaving 41 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	A	801	2	11,11,12	1.28	2 (18%)	15,15,17	2.70	6 (40%)
2	GLC	A	802	2	11,11,12	0.94	0	15,15,17	2.25	4 (26%)
2	GLC	A	803	2	11,11,12	1.10	1 (9%)	15,15,17	1.80	4 (26%)
2	GLC	A	804	2	11,11,12	1.01	0	15,15,17	1.80	5 (33%)
2	GLC	A	805	2	11,11,12	1.20	1 (9%)	15,15,17	2.61	7 (46%)
2	GLC	A	806	2	11,11,12	1.23	1 (9%)	15,15,17	1.91	4 (26%)
2	GLC	A	807	2	11,11,12	2.42	5 (45%)	15,15,17	2.52	6 (40%)
2	GLC	A	808	2	12,12,12	0.79	0	17,17,17	1.03	0
2	GLC	A	809	2	11,11,12	1.26	2 (18%)	15,15,17	2.41	4 (26%)
2	GLC	A	810	2	11,11,12	1.38	1 (9%)	15,15,17	2.50	4 (26%)
2	GLC	A	811	2	11,11,12	0.81	0	15,15,17	2.11	4 (26%)
2	GLC	A	812	2	12,12,12	1.06	0	17,17,17	1.35	1 (5%)
2	GLC	A	813	2	11,11,12	1.08	1 (9%)	15,15,17	1.74	4 (26%)
2	GLC	A	814	2	11,11,12	0.93	1 (9%)	15,15,17	2.07	7 (46%)
2	GLC	A	815	2	11,11,12	0.42	0	15,15,17	1.79	4 (26%)
2	GLC	A	816	2	11,11,12	1.33	1 (9%)	15,15,17	1.65	4 (26%)
2	GLC	A	817	2	11,11,12	0.88	0	15,15,17	1.33	3 (20%)
2	GLC	A	818	2	11,11,12	0.58	0	15,15,17	1.62	5 (33%)
2	GLC	A	819	2	11,11,12	0.74	0	15,15,17	1.42	1 (6%)
2	GLC	A	820	2	11,11,12	0.69	0	15,15,17	2.02	5 (33%)
2	GLC	A	821	2	11,11,12	0.80	0	15,15,17	1.82	4 (26%)
2	GLC	A	822	2	12,12,12	0.92	0	17,17,17	1.80	4 (23%)
2	GLC	A	823	2	11,11,12	0.88	0	15,15,17	1.77	5 (33%)
2	GLC	A	824	2	11,11,12	1.08	1 (9%)	15,15,17	1.66	2 (13%)
2	GLC	A	825	2	11,11,12	1.09	2 (18%)	15,15,17	2.25	6 (40%)
2	GLC	A	826	2	12,12,12	0.93	0	17,17,17	2.18	7 (41%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	A	827	2	11,11,12	0.81	0	15,15,17	1.50	4 (26%)
2	GLC	A	828	2	11,11,12	0.64	0	15,15,17	1.76	5 (33%)
2	GLC	A	829	2	11,11,12	1.13	1 (9%)	15,15,17	3.46	8 (53%)
2	GLC	A	830	2	11,11,12	1.12	1 (9%)	15,15,17	1.31	2 (13%)
2	GLC	A	831	2	11,11,12	0.65	0	15,15,17	1.60	3 (20%)
2	GLC	A	832	2	11,11,12	0.66	0	15,15,17	1.72	5 (33%)
2	GLC	A	833	2	12,12,12	0.95	0	17,17,17	2.20	5 (29%)
2	GLC	A	834	2	11,11,12	0.71	0	15,15,17	2.06	2 (13%)
2	GLC	A	835	2	11,11,12	1.13	1 (9%)	15,15,17	1.98	7 (46%)
2	GLC	A	836	2	11,11,12	0.91	0	15,15,17	1.73	3 (20%)
2	GLC	A	837	2	12,12,12	1.28	1 (8%)	17,17,17	2.21	6 (35%)
2	GLC	A	838	2	11,11,12	1.98	2 (18%)	15,15,17	2.74	8 (53%)
2	GLC	A	839	2	11,11,12	1.68	1 (9%)	15,15,17	1.94	5 (33%)
3	GOL	A	840	-	5,5,5	1.24	1 (20%)	5,5,5	1.34	1 (20%)
3	GOL	A	841	-	5,5,5	1.00	0	5,5,5	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	801	2	-	0/2/19/22	0/1/1/1
2	GLC	A	802	2	-	0/2/19/22	0/1/1/1
2	GLC	A	803	2	-	0/2/19/22	0/1/1/1
2	GLC	A	804	2	-	0/2/19/22	0/1/1/1
2	GLC	A	805	2	-	0/2/19/22	0/1/1/1
2	GLC	A	806	2	-	0/2/19/22	0/1/1/1
2	GLC	A	807	2	-	0/2/19/22	0/1/1/1
2	GLC	A	808	2	-	0/2/22/22	0/1/1/1
2	GLC	A	809	2	-	0/2/19/22	0/1/1/1
2	GLC	A	810	2	-	0/2/19/22	0/1/1/1
2	GLC	A	811	2	-	0/2/19/22	0/1/1/1
2	GLC	A	812	2	-	0/2/22/22	0/1/1/1
2	GLC	A	813	2	-	0/2/19/22	0/1/1/1
2	GLC	A	814	2	-	0/2/19/22	0/1/1/1
2	GLC	A	815	2	-	0/2/19/22	0/1/1/1
2	GLC	A	816	2	-	0/2/19/22	0/1/1/1
2	GLC	A	817	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	818	2	-	0/2/19/22	0/1/1/1
2	GLC	A	819	2	-	0/2/19/22	0/1/1/1
2	GLC	A	820	2	-	0/2/19/22	0/1/1/1
2	GLC	A	821	2	-	0/2/19/22	0/1/1/1
2	GLC	A	822	2	-	0/2/22/22	0/1/1/1
2	GLC	A	823	2	-	0/2/19/22	0/1/1/1
2	GLC	A	824	2	-	0/2/19/22	0/1/1/1
2	GLC	A	825	2	-	0/2/19/22	0/1/1/1
2	GLC	A	826	2	-	0/2/22/22	0/1/1/1
2	GLC	A	827	2	-	0/2/19/22	0/1/1/1
2	GLC	A	828	2	-	0/2/19/22	0/1/1/1
2	GLC	A	829	2	-	0/2/19/22	0/1/1/1
2	GLC	A	830	2	-	0/2/19/22	0/1/1/1
2	GLC	A	831	2	-	0/2/19/22	0/1/1/1
2	GLC	A	832	2	-	0/2/19/22	0/1/1/1
2	GLC	A	833	2	-	0/2/22/22	0/1/1/1
2	GLC	A	834	2	-	0/2/19/22	0/1/1/1
2	GLC	A	835	2	-	0/2/19/22	0/1/1/1
2	GLC	A	836	2	-	0/2/19/22	0/1/1/1
2	GLC	A	837	2	-	0/2/22/22	0/1/1/1
2	GLC	A	838	2	-	0/2/19/22	0/1/1/1
2	GLC	A	839	2	-	0/2/19/22	0/1/1/1
3	GOL	A	840	-	-	0/4/4/4	0/0/0/0
3	GOL	A	841	-	-	0/4/4/4	0/0/0/0

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	838	GLC	O5-C1	-3.69	1.37	1.43
2	A	813	GLC	O5-C1	-2.66	1.39	1.43
2	A	824	GLC	O5-C1	-2.55	1.39	1.43
2	A	809	GLC	O2-C2	-2.15	1.38	1.43
2	A	829	GLC	O5-C1	-2.12	1.40	1.43

The worst 5 of 174 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	GLC	O5-C1-C2	-5.41	102.24	110.89
2	A	806	GLC	O5-C1-C2	-4.78	103.26	110.89
2	A	829	GLC	O6-C6-C5	-4.77	95.38	111.30
2	A	820	GLC	O4-C4-C3	-4.22	100.84	110.36
2	A	836	GLC	C1-O5-C5	-4.20	105.96	112.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

10 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	GLC	2	0
2	A	805	GLC	1	0
2	A	812	GLC	1	0
2	A	813	GLC	1	0
2	A	814	GLC	1	0
2	A	815	GLC	1	0
2	A	819	GLC	2	0
2	A	820	GLC	4	0
2	A	821	GLC	5	0
3	A	840	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	755/793 (95%)	-0.35	13 (1%) 73 79	13, 22, 44, 83	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	758	VAL	5.8
1	A	456	TYR	4.0
1	A	35	ASN	2.5
1	A	457	GLY	2.4
1	A	450	GLU	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GLC	A	804	11/12	0.92	0.22	6.33	34,42,48,50	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GLC	A	807	11/12	0.70	0.29	6.28	32,57,68,68	0
2	GLC	A	801	11/12	0.73	0.27	5.73	32,62,72,73	0
2	GLC	A	837	12/12	0.86	0.22	5.10	44,57,59,62	0
3	GOL	A	840	6/6	0.88	0.18	4.80	39,46,55,55	0
2	GLC	A	813	11/12	0.95	0.17	4.33	31,39,45,52	0
2	GLC	A	826	12/12	0.85	0.33	4.27	38,56,65,68	0
2	GLC	A	838	11/12	0.90	0.19	4.26	33,42,53,53	0
2	GLC	A	810	11/12	0.90	0.24	3.26	31,41,53,55	0
3	GOL	A	841	6/6	0.89	0.18	2.52	33,43,45,51	0
2	GLC	A	816	11/12	0.93	0.17	2.33	26,33,36,45	0
2	GLC	A	803	11/12	0.92	0.17	1.64	41,46,50,51	0
2	GLC	A	802	11/12	0.89	0.15	1.16	45,54,61,63	0
2	GLC	A	818	11/12	0.96	0.13	1.06	26,30,34,39	0
2	GLC	A	835	11/12	0.96	0.12	0.88	25,27,30,33	0
2	GLC	A	824	11/12	0.95	0.12	-0.03	28,33,40,44	0
2	GLC	A	817	11/12	0.97	0.09	-0.18	22,25,27,31	0
2	GLC	A	812	12/12	0.95	0.10	-0.20	32,35,44,45	0
2	GLC	A	832	11/12	0.96	0.10	-0.27	24,31,40,44	0
2	GLC	A	823	11/12	0.97	0.09	-0.38	28,30,38,40	0
2	GLC	A	831	11/12	0.96	0.07	-1.39	26,29,33,35	0
2	GLC	A	820	11/12	0.93	0.30	-	60,63,72,75	0
2	GLC	A	830	11/12	0.95	0.17	-	36,37,40,41	0
2	GLC	A	834	11/12	0.98	0.10	-	22,26,30,30	0
2	GLC	A	821	11/12	0.94	0.31	-	63,65,72,72	0
2	GLC	A	829	11/12	0.96	0.21	-	43,47,49,53	0
2	GLC	A	827	11/12	0.87	0.32	-	59,69,76,79	0
2	GLC	A	805	11/12	0.94	0.17	-	29,40,42,54	0
2	GLC	A	815	11/12	0.86	0.30	-	40,57,65,68	0
4	MG	A	842	1/1	0.98	0.02	-	23,23,23,23	0
2	GLC	A	814	11/12	0.77	0.35	-	56,71,78,84	0
2	GLC	A	806	11/12	0.90	0.26	-	37,49,59,60	0
2	GLC	A	833	12/12	0.92	0.32	-	36,52,58,60	0
2	GLC	A	836	11/12	0.94	0.33	-	40,46,56,67	0
2	GLC	A	828	11/12	0.93	0.23	-	53,57,64,66	0
2	GLC	A	811	11/12	0.87	0.39	-	59,68,74,75	0
2	GLC	A	825	11/12	0.88	0.26	-	51,61,66,76	0
2	GLC	A	809	11/12	0.94	0.19	-	27,34,39,40	0
2	GLC	A	819	11/12	0.94	0.22	-	40,49,54,60	0
2	GLC	A	822	12/12	0.88	0.27	-	40,55,59,63	0
2	GLC	A	839	11/12	0.93	0.28	-	42,45,55,58	0
2	GLC	A	808	12/12	0.85	0.27	-	35,59,64,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.