



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:58 PM GMT

PDB ID : 4GU4
Title : Crystal structure of the T1L reovirus attachment protein sigma1 in complex with alpha-2,3-sialyllactose
Authors : Reiss, K.; Stehle, T.
Deposited on : 2012-08-29
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1051 (3.60-3.40)
Clashscore	102246	1157 (3.60-3.40)
Ramachandran outliers	100387	1120 (3.60-3.40)
Sidechain outliers	100360	1121 (3.60-3.40)
RSRZ outliers	91569	1058 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	 80% 13% 6%
1	B	219	 87% 6% 6%
1	C	219	 79% 14% 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer capsid protein sigma-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	205	Total	C	N	O	S	0	0	0
			1589	1016	265	301	7			
1	B	205	Total	C	N	O	S	0	0	0
			1607	1024	267	309	7			
1	C	205	Total	C	N	O	S	0	0	0
			1598	1020	267	304	7			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	252	GLY	-	EXPRESSION TAG	UNP P04506
A	253	VAL	-	EXPRESSION TAG	UNP P04506
A	254	LEU	-	EXPRESSION TAG	UNP P04506
A	255	ASN	-	EXPRESSION TAG	UNP P04506
A	256	GLN	-	EXPRESSION TAG	UNP P04506
A	257	GLY	-	EXPRESSION TAG	UNP P04506
A	258	VAL	-	EXPRESSION TAG	UNP P04506
A	259	THR	-	EXPRESSION TAG	UNP P04506
A	260	SER	-	EXPRESSION TAG	UNP P04506
B	252	GLY	-	EXPRESSION TAG	UNP P04506
B	253	VAL	-	EXPRESSION TAG	UNP P04506
B	254	LEU	-	EXPRESSION TAG	UNP P04506
B	255	ASN	-	EXPRESSION TAG	UNP P04506
B	256	GLN	-	EXPRESSION TAG	UNP P04506
B	257	GLY	-	EXPRESSION TAG	UNP P04506
B	258	VAL	-	EXPRESSION TAG	UNP P04506
B	259	THR	-	EXPRESSION TAG	UNP P04506
B	260	SER	-	EXPRESSION TAG	UNP P04506
C	252	GLY	-	EXPRESSION TAG	UNP P04506
C	253	VAL	-	EXPRESSION TAG	UNP P04506
C	254	LEU	-	EXPRESSION TAG	UNP P04506
C	255	ASN	-	EXPRESSION TAG	UNP P04506
C	256	GLN	-	EXPRESSION TAG	UNP P04506

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	257	GLY	-	EXPRESSION TAG	UNP P04506
C	258	VAL	-	EXPRESSION TAG	UNP P04506
C	259	THR	-	EXPRESSION TAG	UNP P04506
C	260	SER	-	EXPRESSION TAG	UNP P04506

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	3	Total	C	N	O	0	0
			43	23	1	19		
2	B	3	Total	C	N	O	0	0
			43	23	1	19		


- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	2	Total	C	N	O	0	0
			32	17	1	14		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Outer capsid protein sigma-1



- Molecule 1: Outer capsid protein sigma-1



- Molecule 1: Outer capsid protein sigma-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	149.43Å 149.43Å 165.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.91 – 3.50 48.91 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.8 (48.91-3.50) 97.7 (48.91-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.186 , 0.197 0.200 , 0.210	Depositor DCC
R_{free} test set	2664 reflections (11.06%)	DCC
Wilson B-factor (Å ²)	76.6	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 67.2	EDS
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 42023 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4912	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, BGC, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/1630	0.66	0/2231
1	B	0.49	0/1648	0.65	0/2254
1	C	0.45	0/1639	0.64	0/2242
All	All	0.48	0/4917	0.65	0/6727

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1589	0	1508	24	0
1	B	1607	0	1543	10	0
1	C	1598	0	1525	20	0
2	A	43	0	37	0	0
2	B	43	0	37	0	0
3	C	32	0	28	0	0
All	All	4912	0	4678	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:409:LEU:HB2	1:A:433:VAL:HG13	1.50	0.91
1:A:420:ASN:O	1:A:421:ASN:CB	2.30	0.80
1:A:409:LEU:HB2	1:A:433:VAL:CG1	2.12	0.79
1:A:420:ASN:O	1:A:421:ASN:HB2	1.82	0.77
1:A:311:TYR:HD2	1:B:324:VAL:HB	1.62	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/219 (93%)	197 (97%)	5 (2%)	1 (0%)	34	78
1	B	203/219 (93%)	200 (98%)	3 (2%)	0	100	100
1	C	203/219 (93%)	202 (100%)	1 (0%)	0	100	100
All	All	609/657 (93%)	599 (98%)	9 (2%)	1 (0%)	52	88

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	421	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/192 (88%)	168 (99%)	1 (1%)	90	97
1	B	177/192 (92%)	176 (99%)	1 (1%)	90	97
1	C	172/192 (90%)	171 (99%)	1 (1%)	90	97
All	All	518/576 (90%)	515 (99%)	3 (1%)	90	97

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	290	THR
1	B	283	THR
1	C	283	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	421	ASN
1	A	445	ASN
1	C	297	ASN
1	C	322	ASN
1	C	445	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SIA	A	501	2	16,20,21	0.43	0	18,28,31	1.41	4 (22%)
2	GAL	A	502	2	11,11,12	0.28	0	14,15,17	0.88	1 (7%)
2	BGC	A	503	2	12,12,12	0.26	0	17,17,17	0.39	0
2	SIA	B	501	2	16,20,21	0.43	0	18,28,31	1.41	4 (22%)
2	GAL	B	502	2	11,11,12	0.30	0	14,15,17	0.69	0
2	BGC	B	503	2	12,12,12	0.17	0	17,17,17	0.51	0
3	SIA	C	501	3	16,20,21	0.43	0	18,28,31	1.46	5 (27%)
3	GAL	C	502	3	12,12,12	0.27	0	17,17,17	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SIA	A	501	2	-	0/14/34/38	0/1/1/1
2	GAL	A	502	2	-	0/2/19/22	0/1/1/1
2	BGC	A	503	2	-	0/2/22/22	0/1/1/1
2	SIA	B	501	2	-	0/14/34/38	0/1/1/1
2	GAL	B	502	2	-	0/2/19/22	0/1/1/1
2	BGC	B	503	2	-	0/2/22/22	0/1/1/1
3	SIA	C	501	3	-	0/14/34/38	0/1/1/1
3	GAL	C	502	3	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	SIA	C6-C5-N5	-2.40	106.88	111.07
2	A	501	SIA	C6-C5-N5	-2.38	106.92	111.07
3	C	501	SIA	O9-C9-C8	-2.12	106.48	111.10
3	C	501	SIA	C6-C5-N5	-2.02	107.56	111.07
2	B	501	SIA	O6-C6-C5	2.22	112.11	108.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	205/219 (93%)	-0.53	0 100 100	55, 77, 109, 124	0
1	B	205/219 (93%)	-0.44	0 100 100	56, 79, 116, 135	0
1	C	205/219 (93%)	-0.48	0 100 100	56, 91, 112, 121	0
All	All	615/657 (93%)	-0.48	0 100 100	55, 81, 112, 135	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SIA	B	501	20/21	0.96	0.19	0.22	65,76,107,107	0
2	SIA	A	501	20/21	0.94	0.18	0.04	82,88,104,106	0
3	SIA	C	501	20/21	0.93	0.16	-0.73	77,89,107,108	0
3	GAL	C	502	12/12	0.91	0.18	-	110,123,126,126	0
2	BGC	A	503	12/12	0.92	0.28	-	118,124,126,127	0
2	BGC	B	503	12/12	0.91	0.25	-	119,124,131,131	0
2	GAL	B	502	11/12	0.96	0.10	-	98,110,116,118	0
2	GAL	A	502	11/12	0.96	0.15	-	97,104,111,114	0

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.