



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:00 AM GMT

PDB ID : 3GVH  
Title : Crystal structure of Lactate/malate dehydrogenase from Brucella melitensis  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2009-03-31  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	<div> <div>4%</div> <div>90%</div> <div>7% ..</div> </div>
1	B	324	<div> <div>3%</div> <div>92%</div> <div>5% .</div> </div>
1	C	324	<div> <div>3%</div> <div>88%</div> <div>8% ..</div> </div>
1	D	324	<div> <div>3%</div> <div>89%</div> <div>7% .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAD	B	400	-	-	-	X
2	NAD	C	400	-	-	-	X
2	NAD	D	400	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2324	1483	386	444	11			
1	B	313	Total	C	N	O	S	0	1	0
			2297	1467	380	439	11			
1	C	313	Total	C	N	O	S	0	1	0
			2301	1470	381	439	11			
1	D	313	Total	C	N	O	S	0	1	0
			2302	1468	380	443	11			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP Q2YLR9
A	-2	PRO	-	EXPRESSION TAG	UNP Q2YLR9
A	-1	GLY	-	EXPRESSION TAG	UNP Q2YLR9
A	0	SER	-	EXPRESSION TAG	UNP Q2YLR9
B	-3	GLY	-	EXPRESSION TAG	UNP Q2YLR9
B	-2	PRO	-	EXPRESSION TAG	UNP Q2YLR9
B	-1	GLY	-	EXPRESSION TAG	UNP Q2YLR9
B	0	SER	-	EXPRESSION TAG	UNP Q2YLR9
C	-3	GLY	-	EXPRESSION TAG	UNP Q2YLR9
C	-2	PRO	-	EXPRESSION TAG	UNP Q2YLR9
C	-1	GLY	-	EXPRESSION TAG	UNP Q2YLR9
C	0	SER	-	EXPRESSION TAG	UNP Q2YLR9
D	-3	GLY	-	EXPRESSION TAG	UNP Q2YLR9
D	-2	PRO	-	EXPRESSION TAG	UNP Q2YLR9
D	-1	GLY	-	EXPRESSION TAG	UNP Q2YLR9
D	0	SER	-	EXPRESSION TAG	UNP Q2YLR9

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	152	Total	O	0	0
			152	152		
3	B	145	Total	O	0	0
			145	145		
3	C	149	Total	O	0	0
			149	149		
3	D	174	Total	O	0	0
			174	174		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Malate dehydrogenase




#### • Molecule 1: Malate dehydrogenase



#### • Molecule 1: Malate dehydrogenase



#### • Molecule 1: Malate dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.75Å 114.35Å 148.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	90.00 – 2.30 19.73 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (90.00-2.30) 100.0 (19.73-2.30)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.15 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.174 , 0.218 0.172 , 0.216	Depositor DCC
$R_{free}$ test set	3526 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	29.1	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 44.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 69915 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10020	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.83	1/2363 (0.0%)	0.76	2/3205 (0.1%)
1	B	0.80	0/2337	0.75	1/3168 (0.0%)
1	C	0.81	0/2341	0.81	5/3172 (0.2%)
1	D	0.82	0/2342	0.76	1/3175 (0.0%)
All	All	0.82	1/9383 (0.0%)	0.77	9/12720 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	GLU	CG-CD	6.15	1.61	1.51

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	13	MET	CG-SD-CE	7.54	112.26	100.20
1	C	54	ASP	CB-CG-OD1	-6.78	112.20	118.30
1	B	54	ASP	CB-CG-OD2	-6.57	112.38	118.30
1	C	54	ASP	CB-CG-OD2	5.92	123.63	118.30
1	C	284	ARG	NE-CZ-NH1	-5.74	117.43	120.30
1	D	54	ASP	CB-CG-OD1	-5.68	113.19	118.30
1	C	89	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	A	54	ASP	CB-CG-OD1	-5.50	113.35	118.30
1	A	252	ARG	NE-CZ-NH1	-5.19	117.71	120.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2324	0	2343	15	0
1	B	2297	0	2319	14	0
1	C	2301	0	2330	15	0
1	D	2302	0	2321	15	1
2	A	44	0	26	0	0
2	B	44	0	26	1	0
2	C	44	0	26	3	0
2	D	44	0	26	0	0
3	A	152	0	0	2	0
3	B	145	0	0	1	1
3	C	149	0	0	2	0
3	D	174	0	0	3	0
All	All	10020	0	9417	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:LEU:HD23	1:B:200:MET:HE3	1.31	1.12
1:B:197:LEU:HA	1:B:200:MET:HE2	1.50	0.93
1:B:197:LEU:HD23	1:B:200:MET:CE	1.98	0.93
3:C:434:HOH:O	1:D:13:MET:HE1	1.69	0.92
1:B:81:VAL:HG21	1:B:95:ILE:HG21	1.63	0.79
1:B:197:LEU:HA	1:B:200:MET:CE	2.17	0.75
1:D:13:MET:HA	1:D:13:MET:HE2	1.73	0.71
1:A:13:MET:HA	1:A:13:MET:HE3	1.77	0.67
1:D:37:GLU:OE1	3:D:676:HOH:O	2.13	0.66
1:A:81:VAL:HG13	1:A:82:PRO:HD2	1.77	0.66
1:A:269:MET:CE	1:A:302:VAL:HG13	2.26	0.65
1:B:81:VAL:CG2	1:B:95:ILE:HG21	2.25	0.65
1:A:159:GLU:HG2	3:A:365:HOH:O	1.97	0.64
1:B:269:MET:HE1	1:B:302:VAL:HG13	1.80	0.63
1:D:38:GLY:HA2	3:D:676:HOH:O	2.01	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:13:MET:HA	1:D:13:MET:CE	2.36	0.54
1:B:269:MET:CE	1:B:302:VAL:HG13	2.37	0.54
1:B:197:LEU:CD2	1:B:200:MET:HE3	2.22	0.54
1:A:269:MET:HE1	1:A:302:VAL:HG13	1.91	0.53
1:A:205:GLN:NE2	3:A:666:HOH:O	2.42	0.53
1:C:119:ILE:O	2:C:400:NAD:H2N	2.09	0.52
1:C:35:ILE:HG23	2:C:400:NAD:C2A	2.40	0.52
1:A:142:VAL:HB	1:A:256:VAL:CG1	2.40	0.51
1:A:256:VAL:HG21	1:A:277:ILE:HD11	1.93	0.51
1:D:234:ALA:HB3	1:D:235:PRO:HD3	1.94	0.50
1:C:66:TYR:O	1:C:69:ILE:HG12	2.12	0.50
1:D:269:MET:HE3	1:D:305:VAL:HG12	1.94	0.50
1:A:269:MET:HE2	1:A:302:VAL:HG13	1.94	0.49
1:D:35:ILE:O	1:D:36:ALA:C	2.52	0.48
1:B:269:MET:HE1	1:B:302:VAL:CG1	2.44	0.47
1:C:173:LEU:CD2	1:C:274:PRO:HD3	2.43	0.47
1:C:35:ILE:HD13	2:C:400:NAD:C5A	2.45	0.47
1:C:5:LYS:NZ	1:C:70:GLU:O	2.35	0.47
1:D:134:SER:HB2	1:D:136:LEU:HD13	1.97	0.46
1:C:89:ARG:HH11	1:C:89:ARG:CG	2.29	0.46
1:A:131:GLN:HB3	1:A:141:VAL:HG11	1.98	0.45
1:B:2:ALA:N	3:B:653:HOH:O	2.48	0.45
1:C:121:ASN:ND2	3:C:420:HOH:O	2.49	0.44
1:A:78:THR:HG22	1:A:119:ILE:HD12	2.00	0.44
1:D:142:VAL:HB	1:D:256:VAL:CG1	2.48	0.44
1:D:253:VAL:HA	1:D:275:THR:O	2.19	0.43
1:D:13:MET:HE2	1:D:13:MET:CA	2.44	0.43
1:A:13:MET:CA	1:A:13:MET:HE3	2.48	0.42
1:C:51:SER:N	1:C:52:PRO:CD	2.82	0.42
1:C:194:LEU:HB2	1:C:195:PRO:HD3	2.02	0.42
1:B:119:ILE:O	2:B:400:NAD:H2N	2.20	0.42
1:A:142:VAL:HB	1:A:256:VAL:HG11	2.02	0.42
1:D:194:LEU:HD23	1:D:197:LEU:HD12	2.02	0.42
1:A:173:LEU:CD2	1:A:274:PRO:HD3	2.50	0.41
1:B:81:VAL:CG2	1:B:95:ILE:CG2	2.96	0.41
1:C:242:MET:CE	1:C:256:VAL:HG22	2.51	0.41
1:A:121:ASN:HD22	1:A:122:PRO:HA	1.84	0.41
1:C:13:MET:CE	1:C:13:MET:HA	2.51	0.41
1:C:203:THR:HG21	1:C:208:LEU:HD13	2.03	0.41
1:D:131:GLN:HG3	3:D:587:HOH:O	2.20	0.41
1:C:315:ILE:O	1:C:315:ILE:HG22	2.21	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:260:LEU:HD12	1:D:266:VAL:HB	2.03	0.40
1:B:234:ALA:N	1:B:235:PRO:CD	2.85	0.40
1:C:266:VAL:HG13	1:C:269:MET:HE3	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:37:GLU:OE2	3:B:675:HOH:O[3_445]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/324 (98%)	304 (96%)	11 (4%)	1 (0%)	46	57
1	B	310/324 (96%)	302 (97%)	8 (3%)	0	100	100
1	C	310/324 (96%)	303 (98%)	5 (2%)	2 (1%)	30	36
1	D	310/324 (96%)	304 (98%)	6 (2%)	0	100	100
All	All	1246/1296 (96%)	1213 (97%)	30 (2%)	3 (0%)	52	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	200	MET
1	C	318	SER
1	A	85	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/249 (96%)	233 (98%)	6 (2%)	55	73
1	B	237/249 (95%)	232 (98%)	5 (2%)	61	78
1	C	238/249 (96%)	233 (98%)	5 (2%)	61	78
1	D	239/249 (96%)	234 (98%)	5 (2%)	61	78
All	All	953/996 (96%)	932 (98%)	21 (2%)	60	77

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	ARG
1	A	131	GLN
1	A	172	VAL
1	A	207	LYS
1	A	280	ASN
1	A	308	LEU
1	B	89	ARG
1	B	131	GLN
1	B	153	PHE
1	B	252	ARG
1	B	267	LYS
1	C	13	MET
1	C	89	ARG
1	C	131	GLN
1	C	208	LEU
1	C	310	GLU
1	D	69	ILE
1	D	89	ARG
1	D	131	GLN
1	D	207	LYS
1	D	208	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	GLN
1	B	213	GLN
1	C	102	GLN
1	C	121	ASN
1	C	162	ASN
1	C	213	GLN
1	D	213	GLN
1	D	297	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	A	400	-	38,48,48	0.85	1 (2%)	47,73,73	2.04	8 (17%)
2	NAD	B	400	-	38,48,48	0.74	1 (2%)	47,73,73	2.10	8 (17%)
2	NAD	C	400	-	38,48,48	0.78	1 (2%)	47,73,73	1.93	8 (17%)
2	NAD	D	400	-	38,48,48	0.76	1 (2%)	47,73,73	1.97	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	400	-	-	0/22/62/62	0/5/5/5
2	NAD	B	400	-	-	0/22/62/62	0/5/5/5
2	NAD	C	400	-	-	0/22/62/62	0/5/5/5
2	NAD	D	400	-	-	0/22/62/62	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	400	NAD	O4D-C1D	2.04	1.43	1.41
2	D	400	NAD	O4B-C1B	2.47	1.44	1.41
2	B	400	NAD	O4D-C1D	2.52	1.44	1.41
2	A	400	NAD	O4D-C1D	3.43	1.45	1.41

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	400	NAD	N3A-C2A-N1A	-9.81	121.38	128.89
2	B	400	NAD	N3A-C2A-N1A	-8.43	122.44	128.89
2	C	400	NAD	N3A-C2A-N1A	-8.02	122.76	128.89
2	A	400	NAD	N3A-C2A-N1A	-7.92	122.83	128.89
2	B	400	NAD	C1B-N9A-C4A	-5.26	119.01	126.94
2	C	400	NAD	C1B-N9A-C4A	-4.50	120.15	126.94
2	C	400	NAD	C4A-C5A-N7A	-3.58	106.19	109.48
2	D	400	NAD	PN-O3-PA	-3.55	122.75	132.73
2	C	400	NAD	O7N-C7N-N7N	-3.43	117.77	122.59
2	A	400	NAD	PN-O3-PA	-3.14	123.92	132.73
2	A	400	NAD	O7N-C7N-N7N	-3.13	118.18	122.59
2	B	400	NAD	PN-O3-PA	-3.11	124.00	132.73
2	C	400	NAD	PN-O3-PA	-2.99	124.33	132.73
2	A	400	NAD	C4A-C5A-N7A	-2.83	106.87	109.48
2	D	400	NAD	C1B-N9A-C4A	-2.73	122.82	126.94
2	B	400	NAD	O7N-C7N-N7N	-2.63	118.89	122.59
2	D	400	NAD	O7N-C7N-N7N	-2.53	119.03	122.59
2	D	400	NAD	C4A-C5A-N7A	-2.49	107.18	109.48
2	A	400	NAD	C1B-N9A-C4A	-2.41	123.30	126.94
2	B	400	NAD	C4A-C5A-N7A	-2.38	107.28	109.48
2	C	400	NAD	O3-PN-O5D	-2.29	96.86	102.94
2	B	400	NAD	O7N-C7N-C3N	-2.14	117.25	119.59
2	D	400	NAD	C3N-C2N-N1N	2.69	123.46	120.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	400	NAD	C3N-C2N-N1N	2.74	123.52	120.36
2	D	400	NAD	O7N-C7N-C3N	3.31	123.20	119.59
2	A	400	NAD	C3N-C2N-N1N	3.35	124.22	120.36
2	C	400	NAD	O7N-C7N-C3N	4.02	123.98	119.59
2	A	400	NAD	O7N-C7N-C3N	4.46	124.45	119.59
2	B	400	NAD	O4D-C1D-N1N	5.06	113.69	108.13
2	B	400	NAD	C3N-C7N-N7N	5.50	123.83	117.82
2	A	400	NAD	O4D-C1D-N1N	5.67	114.37	108.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	400	NAD	1	0
2	C	400	NAD	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	318/324 (98%)	-0.21	13 (4%)	41 50	13, 25, 48, 72	0
1	B	313/324 (96%)	-0.30	10 (3%)	51 60	13, 23, 47, 72	0
1	C	313/324 (96%)	-0.23	10 (3%)	51 60	12, 23, 47, 70	0
1	D	313/324 (96%)	-0.30	9 (2%)	55 64	12, 22, 47, 73	0
All	All	1257/1296 (96%)	-0.26	42 (3%)	50 59	12, 23, 47, 73	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	85	PRO	7.4
1	C	82	PRO	4.8
1	B	81	VAL	4.4
1	A	86	GLY	4.2
1	A	82	PRO	3.9
1	B	37	GLU	3.5
1	B	82	PRO	3.5
1	C	88	SER	3.4
1	D	88	SER	3.4
1	A	228	THR	3.4
1	B	2	ALA	3.2
1	A	87	MET	3.2
1	D	81	VAL	3.2
1	B	314	GLY	3.1
1	D	91	ASP	3.1
1	B	88	SER	3.0
1	C	81	VAL	3.0
1	D	228	THR	3.0
1	A	84	LYS	2.9
1	A	317	PRO	2.8
1	C	228	THR	2.8

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	201	GLY	2.8
1	C	37	GLU	2.7
1	A	293	ASP	2.6
1	D	317	PRO	2.6
1	D	90	ASP	2.6
1	B	293	ASP	2.5
1	D	2	ALA	2.5
1	A	2	ALA	2.5
1	A	95	ILE	2.3
1	B	317	PRO	2.3
1	D	314	GLY	2.3
1	C	317	PRO	2.2
1	B	228	THR	2.2
1	A	83	ARG	2.2
1	A	318	SER	2.1
1	D	178	ASP	2.1
1	A	303	ALA	2.1
1	C	89	ARG	2.1
1	C	293	ASP	2.0
1	C	202	TRP	2.0
1	B	89	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAD	B	400	44/44	0.62	0.33	4.01	26,34,36,37	44

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAD	D	400	44/44	0.70	0.29	3.52	34,42,44,45	44
2	NAD	C	400	44/44	0.69	0.28	2.38	31,35,37,39	44
2	NAD	A	400	44/44	0.81	0.23	1.89	25,29,31,32	44

## 6.5 Other polymers [i](#)

There are no such residues in this entry.