



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:00 AM GMT

PDB ID : 3GVJ
Title : Crystal structure of an endo-neuraminidaseNF mutant
Authors : Schulz, E.C.; Dickmanns, A.; Ficner, R.
Deposited on : 2009-03-31
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3129 (1.50-1.46)
Clashscore	102246	3380 (1.50-1.46)
Ramachandran outliers	100387	3310 (1.50-1.46)
Sidechain outliers	100360	3308 (1.50-1.46)
RSRZ outliers	91569	3133 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	670	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SIA	A	2002	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SLB	A	2009	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6457 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-N-acetylneuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	670	Total	C	N	O	S	0	24	0
			5425	3420	946	1040	19			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	241	VAL	-	expression tag	UNP Q858B1
A	242	PRO	-	expression tag	UNP Q858B1
A	243	ARG	-	expression tag	UNP Q858B1
A	244	GLY	-	expression tag	UNP Q858B1
A	245	SER	-	expression tag	UNP Q858B1
A	647	ALA	ARG	engineered	UNP Q858B1

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	5	Total	C	N	O	0	0
			101	55	5	41		

- Molecule 3 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	4	Total	C	N	O	0	0
			81	44	4	33		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	850	Total	O	0	0
			850	850		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endo-N-acetylneuraminidase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	119.09Å 119.09Å 175.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	12.00 – 1.48 19.85 – 1.48	Depositor EDS
% Data completeness (in resolution range)	100.0 (12.00-1.48) 99.9 (19.85-1.48)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 1.48Å)	Xtriage
Refinement program	REFMAC 5.5.0039	Depositor
R, R_{free}	0.194 , 0.216 0.196 , 0.219	Depositor DCC
R_{free} test set	7739 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	12.3	Xtriage
Anisotropy	0.147	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.2	EDS
Estimated twinning fraction	0.043 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 155002 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6457	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, SLB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.65	37/5632 (0.7%)	0.92	7/7671 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	373	GLU	CD-OE1	-7.27	1.17	1.25
1	A	661	TYR	CE2-CZ	-6.77	1.29	1.38
1	A	873	GLU	CD-OE2	-6.49	1.18	1.25
1	A	344	TYR	CD2-CE2	-6.47	1.29	1.39
1	A	504[A]	GLY	C-N	6.46	1.49	1.34
1	A	504[B]	GLY	C-N	6.46	1.49	1.34
1	A	344	TYR	CE1-CZ	-6.25	1.30	1.38
1	A	576	GLU	CD-OE2	-6.15	1.18	1.25
1	A	297	GLU	CD-OE2	-6.08	1.19	1.25
1	A	296	TYR	CD1-CE1	-6.02	1.30	1.39
1	A	851	GLY	C-O	-6.01	1.14	1.23
1	A	574[A]	GLU	N-CA	-5.97	1.34	1.46
1	A	574[B]	GLU	N-CA	-5.97	1.34	1.46
1	A	767	GLY	C-O	-5.85	1.14	1.23
1	A	818	GLU	CD-OE2	-5.84	1.19	1.25
1	A	857	TYR	CD1-CE1	-5.82	1.30	1.39
1	A	646	GLU	CD-OE2	-5.78	1.19	1.25
1	A	649	GLU	CD-OE2	-5.64	1.19	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	366	GLY	C-O	-5.64	1.14	1.23
1	A	716	TYR	CE1-CZ	-5.58	1.31	1.38
1	A	755	GLY	C-O	-5.52	1.14	1.23
1	A	592	TYR	CE1-CZ	-5.49	1.31	1.38
1	A	865	ARG	CZ-NH2	-5.48	1.25	1.33
1	A	725	ASN	C-O	-5.42	1.13	1.23
1	A	592	TYR	CD2-CE2	-5.34	1.31	1.39
1	A	604	GLY	C-O	-5.28	1.15	1.23
1	A	586	TYR	CE1-CZ	-5.27	1.31	1.38
1	A	387	TYR	CD1-CE1	-5.26	1.31	1.39
1	A	644	GLY	C-O	-5.19	1.15	1.23
1	A	404	GLU	CD-OE1	-5.18	1.20	1.25
1	A	309	GLU	CG-CD	5.12	1.59	1.51
1	A	585	LYS	CG-CD	-5.11	1.35	1.52
1	A	387	TYR	CE2-CZ	-5.10	1.31	1.38
1	A	540	GLY	C-O	-5.07	1.15	1.23
1	A	814	GLU	CD-OE1	-5.04	1.20	1.25
1	A	857	TYR	CG-CD2	-5.03	1.32	1.39
1	A	546	VAL	CB-CG2	-5.02	1.42	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	503	MET	O-C-N	-16.65	94.89	123.20
1	A	503	MET	CA-C-N	15.04	146.29	116.20
1	A	865	ARG	NE-CZ-NH2	-11.43	114.59	120.30
1	A	620	LEU	CA-CB-CG	7.73	133.08	115.30
1	A	784	ARG	NE-CZ-NH1	-5.78	117.41	120.30
1	A	503	MET	C-N-CA	-5.69	110.35	122.30
1	A	470	ASP	CB-CG-OD2	-5.49	113.36	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	243	ARG	Peptide
1	A	330	GLN	Peptide
1	A	503	MET	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5425	0	5199	54	0
2	A	101	0	82	5	0
3	A	81	0	66	2	0
4	A	850	0	0	5	0
All	All	6457	0	5347	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:SER:HB3	1:A:355[B]:LEU:HD23	1.38	1.04
1:A:345:MET:HB3	1:A:355[B]:LEU:HD21	1.47	0.96
1:A:449:HIS:HD2	1:A:451:LEU:H	1.17	0.88
1:A:345:MET:HB3	1:A:355[A]:LEU:HD11	1.57	0.84
1:A:503:MET:HG2	1:A:504[A]:GLY:HA3	1.60	0.83
1:A:503:MET:HG2	1:A:504[B]:GLY:HA3	1.68	0.72
1:A:373:GLU:OE2	1:A:508:HIS:HE1	1.72	0.72
1:A:449:HIS:CD2	1:A:451:LEU:H	2.05	0.72
1:A:322[A]:THR:HG22	1:A:324:TYR:H	1.59	0.68
1:A:508:HIS:HD2	4:A:984:HOH:O	1.78	0.65
1:A:347:SER:HB3	1:A:355[B]:LEU:CD2	2.22	0.65
1:A:584:ILE:O	1:A:585:LYS:HD2	1.95	0.65
1:A:499[C]:LYS:NZ	4:A:1281:HOH:O	2.30	0.64
1:A:544:GLY:HA2	1:A:549[B]:ARG:HG2	1.80	0.64
1:A:527:HIS:HD2	1:A:582:PRO:O	1.82	0.62
1:A:347:SER:HB3	1:A:355[A]:LEU:HD13	1.81	0.62
2:A:2001:SLB:O7	2:A:2001:SLB:H32	2.01	0.61
1:A:336:TYR:OH	1:A:425[B]:ARG:HD3	2.02	0.59
1:A:499[C]:LYS:CE	2:A:2005:SIA:O7	2.52	0.58
1:A:425[A]:ARG:HD2	1:A:427:LEU:HD21	1.86	0.58
1:A:499[C]:LYS:HE2	2:A:2005:SIA:O7	2.04	0.56
1:A:330:GLN:HB2	1:A:527:HIS:HE1	1.71	0.55
1:A:553:LEU:CD2	1:A:591[A]:LEU:HD21	2.36	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:TYR:OH	1:A:350:HIS:HE1	1.89	0.54
3:A:2006:SIA:O6	3:A:2006:SIA:H92	2.08	0.54
1:A:425[B]:ARG:HH11	1:A:425[B]:ARG:HG3	1.71	0.53
1:A:457:VAL:HA	1:A:504[B]:GLY:O	2.09	0.53
1:A:584:ILE:C	1:A:585:LYS:HD2	2.29	0.53
1:A:330:GLN:HB2	1:A:527:HIS:CE1	2.44	0.52
1:A:355[A]:LEU:HD12	1:A:356:HIS:N	2.25	0.51
1:A:628:HIS:HE1	1:A:658:ASP:OD1	1.94	0.50
1:A:328:TRP:HD1	1:A:330:GLN:HE21	1.57	0.50
1:A:322[A]:THR:HG21	4:A:107:HOH:O	2.11	0.49
1:A:364:ASP:OD2	1:A:368[A]:THR:OG1	2.27	0.48
1:A:542:HIS:HD2	4:A:916:HOH:O	1.95	0.48
1:A:445:HIS:HD2	1:A:481:ASN:HD21	1.62	0.47
1:A:544:GLY:HA2	1:A:549[A]:ARG:HG2	1.96	0.46
2:A:2001:SLB:C3	2:A:2001:SLB:O7	2.64	0.46
1:A:553:LEU:HD21	1:A:591[A]:LEU:HD21	1.97	0.45
1:A:425[B]:ARG:NH1	1:A:425[B]:ARG:HG3	2.32	0.45
1:A:265[C]:THR:HG23	1:A:266:PRO:N	2.32	0.45
1:A:325:TYR:OH	1:A:350:HIS:CE1	2.70	0.45
3:A:2009:SLB:H92	3:A:2009:SLB:O6	2.17	0.44
1:A:243:ARG:HB3	1:A:244:GLY:HA3	1.99	0.44
1:A:444:ILE:HG22	1:A:446:VAL:HG23	2.01	0.43
1:A:499[C]:LYS:HE3	2:A:2005:SIA:O7	2.17	0.43
1:A:440[A]:ARG:HE	1:A:489:GLN:HE21	1.64	0.43
1:A:679[B]:ASN:ND2	1:A:681:ASP:H	2.17	0.43
1:A:445:HIS:HD2	1:A:481:ASN:ND2	2.16	0.43
1:A:279:LYS:HE2	4:A:979:HOH:O	2.18	0.43
1:A:553:LEU:HD22	1:A:591[A]:LEU:HD21	2.01	0.42
1:A:735:LYS:O	1:A:743:HIS:HE1	2.03	0.41
1:A:547:ALA:HA	1:A:548:PRO:HA	1.64	0.41
1:A:247:LYS:HD3	1:A:247:LYS:HA	1.75	0.41
1:A:576:GLU:N	1:A:577:PRO:CD	2.84	0.41
1:A:458:ASN:O	1:A:504[A]:GLY:HA2	2.20	0.40
1:A:322[A]:THR:HB	1:A:326:ASN:OD1	2.20	0.40
1:A:421:ARG:HD2	1:A:512:TRP:CD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	696/670 (104%)	673 (97%)	22 (3%)	1 (0%)	56 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	704	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	593/566 (105%)	590 (100%)	3 (0%)	92 80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	359	TRP
1	A	620	LEU
1	A	762	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	330	GLN
1	A	338	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	350	HIS
1	A	397	ASN
1	A	445	HIS
1	A	449	HIS
1	A	461	ASN
1	A	481	ASN
1	A	489	GLN
1	A	508	HIS
1	A	527	HIS
1	A	542	HIS
1	A	561	ASN
1	A	625	ASN
1	A	628	HIS
1	A	743	HIS
1	A	832	ASN
1	A	906	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SLB	A	2001	2	17,21,21	1.24	2 (11%)	19,31,31	5.34	6 (31%)
2	SIA	A	2002	2	16,20,21	0.35	0	18,28,31	1.19	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SIA	A	2003	2	16,20,21	0.34	0	18,28,31	1.07	1 (5%)
2	SIA	A	2004	2	16,20,21	0.41	0	18,28,31	1.14	2 (11%)
2	SIA	A	2005	2	16,20,21	0.40	0	18,28,31	1.04	1 (5%)
3	SIA	A	2006	3	16,20,21	1.18	2 (12%)	18,28,31	2.83	6 (33%)
3	SIA	A	2007	3	16,20,21	0.38	0	18,28,31	1.02	1 (5%)
3	SIA	A	2008	3	16,20,21	0.38	0	18,28,31	0.98	1 (5%)
3	SLB	A	2009	3	17,21,21	1.30	3 (17%)	19,31,31	5.27	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SLB	A	2001	2	-	0/14/38/38	0/1/1/1
2	SIA	A	2002	2	-	0/14/34/38	0/1/1/1
2	SIA	A	2003	2	-	0/14/34/38	0/1/1/1
2	SIA	A	2004	2	-	0/14/34/38	0/1/1/1
2	SIA	A	2005	2	-	0/14/34/38	0/1/1/1
3	SIA	A	2006	3	-	0/14/34/38	0/1/1/1
3	SIA	A	2007	3	-	0/14/34/38	0/1/1/1
3	SIA	A	2008	3	-	0/14/34/38	0/1/1/1
3	SLB	A	2009	3	-	0/14/38/38	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2006	SIA	C5-N5	-2.39	1.41	1.45
3	A	2006	SIA	C9-C8	-2.36	1.45	1.52
3	A	2009	SLB	O6-C2	2.05	1.44	1.42
3	A	2009	SLB	C3-C2	2.09	1.54	1.51
2	A	2001	SLB	O10-C10	2.33	1.28	1.23
2	A	2001	SLB	O2-C2	3.66	1.43	1.40
3	A	2009	SLB	O2-C2	4.28	1.44	1.40

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2009	SLB	O2-C2-C3	-18.05	86.87	109.41
2	A	2001	SLB	O2-C2-C3	-17.42	87.66	109.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	SLB	O2-C2-O6	-13.93	87.49	110.22
3	A	2009	SLB	O2-C2-O6	-13.71	87.85	110.22
3	A	2006	SIA	O9-C9-C8	-9.22	91.05	111.10
3	A	2006	SIA	C3-C4-C5	-4.77	106.16	111.47
2	A	2002	SIA	C6-C5-N5	-3.25	105.41	111.07
2	A	2004	SIA	C6-C5-N5	-3.24	105.43	111.07
2	A	2003	SIA	C6-C5-N5	-3.23	105.45	111.07
3	A	2006	SIA	C7-C6-C5	-3.03	109.74	114.32
3	A	2008	SIA	C6-C5-N5	-2.88	106.05	111.07
2	A	2005	SIA	C4-C5-N5	-2.37	105.25	110.41
2	A	2001	SLB	C7-C6-C5	-2.36	110.74	114.32
2	A	2004	SIA	C4-C5-N5	-2.29	105.43	110.41
2	A	2001	SLB	C11-C10-N5	-2.28	111.75	116.11
2	A	2002	SIA	C7-C6-C5	-2.22	110.97	114.32
2	A	2001	SLB	C5-N5-C10	-2.14	117.62	123.10
3	A	2006	SIA	C11-C10-N5	-2.01	112.27	116.11
3	A	2007	SIA	C7-C6-C5	2.04	117.41	114.32
3	A	2006	SIA	O6-C6-C5	2.18	112.06	108.48
3	A	2006	SIA	O8-C8-C7	2.28	114.74	109.02
2	A	2001	SLB	O6-C6-C7	4.72	114.39	107.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	SLB	2	0
2	A	2005	SIA	3	0
3	A	2006	SIA	1	0
3	A	2009	SLB	1	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	670/670 (100%)	-0.12	11 (1%) 74 78	8, 13, 21, 37	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	GLY	4.4
1	A	243	ARG	3.8
1	A	546	VAL	3.7
1	A	504[A]	GLY	3.2
1	A	549[A]	ARG	3.1
1	A	547	ALA	3.0
1	A	910	THR	2.9
1	A	491	THR	2.6
1	A	682	ASP	2.5
1	A	548	PRO	2.4
1	A	492	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SLB	A	2009	21/21	0.85	0.19	5.97	20,26,32,35	0
2	SIA	A	2002	20/21	0.89	0.13	2.45	19,22,29,30	0
3	SIA	A	2006	20/21	0.91	0.13	1.92	18,21,30,34	0
2	SIA	A	2003	20/21	0.93	0.11	1.87	18,20,26,27	0
3	SIA	A	2008	20/21	0.94	0.09	1.72	13,18,25,28	0
2	SIA	A	2005	20/21	0.81	0.21	1.53	32,36,38,39	0
2	SIA	A	2004	20/21	0.81	0.23	-	30,32,37,38	0
2	SLB	A	2001	21/21	0.91	0.23	-	32,36,36,36	16
3	SIA	A	2007	20/21	0.92	0.13	-	16,24,28,29	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.