



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:01 AM GMT

PDB ID : 3GY1
Title : CRYSTAL STRUCTURE OF putative mandelate racemase/muconate lactonizing protein from *Clostridium beijerinckii* NCIMB 8052
Authors : Malashkevich, V.N.; Toro, R.; Morano, C.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2009-04-03
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	408	 2% 79% 14% • 5%
1	B	408	 1% 81% 13% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	A	500	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mandelate racemase/muconate lactonizing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	1	0
			3108	1988	531	575	14			
1	B	388	Total	C	N	O	S	0	1	0
			3120	1993	536	577	14			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP A6M2W4
A	2	SER	-	expression tag	UNP A6M2W4
A	3	LEU	-	expression tag	UNP A6M2W4
A	401	GLU	-	expression tag	UNP A6M2W4
A	402	GLY	-	expression tag	UNP A6M2W4
A	403	HIS	-	expression tag	UNP A6M2W4
A	404	HIS	-	expression tag	UNP A6M2W4
A	405	HIS	-	expression tag	UNP A6M2W4
A	406	HIS	-	expression tag	UNP A6M2W4
A	407	HIS	-	expression tag	UNP A6M2W4
A	408	HIS	-	expression tag	UNP A6M2W4
B	1	MET	-	expression tag	UNP A6M2W4
B	2	SER	-	expression tag	UNP A6M2W4
B	3	LEU	-	expression tag	UNP A6M2W4
B	401	GLU	-	expression tag	UNP A6M2W4
B	402	GLY	-	expression tag	UNP A6M2W4
B	403	HIS	-	expression tag	UNP A6M2W4
B	404	HIS	-	expression tag	UNP A6M2W4
B	405	HIS	-	expression tag	UNP A6M2W4
B	406	HIS	-	expression tag	UNP A6M2W4
B	407	HIS	-	expression tag	UNP A6M2W4
B	408	HIS	-	expression tag	UNP A6M2W4

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Mg 1	0	0
2	A	1	Total 1	Mg 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	405	Total 405	O 405	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mandelate racemase/muconate lactonizing protein



- Molecule 1: Mandelate racemase/muconate lactonizing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	111.31Å 111.31Å 141.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.89 – 1.60 19.89 – 1.60	Depositor EDS
% Data completeness (in resolution range)	90.5 (19.89-1.60) 90.5 (19.89-1.60)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, R_{free}	0.179 , 0.212 0.179 , 0.211	Depositor DCC
R_{free} test set	5347 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	6 of 105785 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6635	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 72.98 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.9976e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.40	16/3186 (0.5%)	1.28	23/4330 (0.5%)
1	B	1.44	19/3198 (0.6%)	1.20	14/4345 (0.3%)
All	All	1.42	35/6384 (0.5%)	1.24	37/8675 (0.4%)

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	383	TYR	CD2-CE2	-10.22	1.24	1.39
1	A	383	TYR	CD2-CE2	-9.47	1.25	1.39
1	B	383	TYR	CD1-CE1	-8.78	1.26	1.39
1	A	383	TYR	CD1-CE1	-8.56	1.26	1.39
1	A	211	ARG	CB-CG	-7.36	1.32	1.52
1	A	76	MET	CB-CG	7.29	1.74	1.51
1	B	130	VAL	CB-CG2	6.58	1.66	1.52
1	B	188	VAL	CB-CG2	6.47	1.66	1.52
1	A	308	TRP	CE3-CZ3	6.44	1.49	1.38
1	B	249	SER	CB-OG	-6.42	1.33	1.42
1	B	348	PHE	CE1-CZ	6.34	1.49	1.37
1	B	35	TYR	CE1-CZ	6.23	1.46	1.38
1	A	305	ARG	CZ-NH1	6.14	1.41	1.33
1	A	100	TRP	CG-CD1	6.01	1.45	1.36
1	A	302	PHE	CD1-CE1	5.96	1.51	1.39
1	B	2	SER	N-CA	5.93	1.58	1.46
1	A	95	VAL	CB-CG2	5.89	1.65	1.52
1	B	383	TYR	CE2-CZ	-5.86	1.30	1.38
1	B	76	MET	CB-CG	5.85	1.70	1.51
1	A	224	VAL	CB-CG2	-5.75	1.40	1.52
1	B	135	GLU	CG-CD	5.73	1.60	1.51
1	B	301	ALA	CA-CB	5.63	1.64	1.52
1	B	278	PHE	CE2-CZ	5.54	1.47	1.37
1	A	149	TYR	CG-CD1	5.45	1.46	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	115	PHE	CD1-CE1	5.40	1.50	1.39
1	B	341	ASN	CB-CG	5.36	1.63	1.51
1	B	333	ALA	CA-CB	5.33	1.63	1.52
1	A	191	PHE	CG-CD2	5.30	1.46	1.38
1	B	26	VAL	CB-CG2	5.30	1.64	1.52
1	A	98	ALA	CA-CB	5.26	1.63	1.52
1	B	28	VAL	CB-CG1	5.13	1.63	1.52
1	A	383	TYR	CE2-CZ	-5.07	1.31	1.38
1	A	220	PHE	CG-CD1	5.02	1.46	1.38
1	B	112	TYR	CE2-CZ	5.01	1.45	1.38
1	B	183	TYR	CG-CD2	5.00	1.45	1.39

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	305	ARG	NE-CZ-NH2	-20.51	110.04	120.30
1	A	305	ARG	NE-CZ-NH1	19.06	129.83	120.30
1	B	280	ARG	NE-CZ-NH2	15.36	127.98	120.30
1	A	280	ARG	NE-CZ-NH2	13.97	127.29	120.30
1	B	195	ARG	NE-CZ-NH2	-11.33	114.64	120.30
1	B	280	ARG	NE-CZ-NH1	-10.98	114.81	120.30
1	A	239	ASP	CB-CG-OD1	8.58	126.02	118.30
1	A	280	ARG	NE-CZ-NH1	-8.29	116.16	120.30
1	A	207	ASP	CB-CG-OD1	8.16	125.64	118.30
1	B	239	ASP	CB-CG-OD1	7.66	125.20	118.30
1	A	239	ASP	CB-CG-OD2	-7.66	111.41	118.30
1	A	111	LEU	CB-CG-CD1	-7.26	98.66	111.00
1	A	270	LEU	CB-CG-CD1	-7.15	98.85	111.00
1	A	109	MET	CA-CB-CG	7.02	125.23	113.30
1	A	211	ARG	CG-CD-NE	-6.87	97.38	111.80
1	A	207	ASP	CB-CG-OD2	-6.53	112.42	118.30
1	B	77	MET	CG-SD-CE	-6.49	89.81	100.20
1	B	115	PHE	CB-CG-CD2	-6.40	116.32	120.80
1	B	109	MET	CA-CB-CG	6.38	124.15	113.30
1	B	118	LYS	CD-CE-NZ	-5.97	97.98	111.70
1	A	77	MET	CB-CG-SD	5.86	129.98	112.40
1	A	115	PHE	CB-CG-CD2	-5.70	116.81	120.80
1	A	211	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	305	ARG	CD-NE-CZ	5.56	131.38	123.60
1	B	305[A]	ARG	NE-CZ-NH1	-5.54	117.53	120.30
1	B	305[B]	ARG	NE-CZ-NH1	-5.54	117.53	120.30
1	B	293	LEU	CB-CG-CD1	-5.33	101.93	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	111	LEU	CB-CG-CD2	-5.25	102.08	111.00
1	A	261	PHE	CB-CG-CD2	-5.22	117.14	120.80
1	A	45	ARG	NE-CZ-NH1	5.19	122.90	120.30
1	A	370	ASP	CB-CG-OD1	5.19	122.97	118.30
1	A	234	ASP	CB-CG-OD1	5.16	122.94	118.30
1	A	270	LEU	CB-CG-CD2	5.14	119.73	111.00
1	B	153	ARG	NE-CZ-NH2	5.12	122.86	120.30
1	B	280	ARG	CD-NE-CZ	5.11	130.76	123.60
1	A	325	LEU	CB-CG-CD2	5.08	119.64	111.00
1	B	278	PHE	CB-CG-CD2	-5.07	117.25	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3108	0	3051	41	0
1	B	3120	0	3061	36	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	405	0	0	22	1
All	All	6635	0	6112	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:MET:CG	1:A:76:MET:CB	1.74	1.64
1:B:268:LYS:HD3	3:B:596:HOH:O	1.41	1.21
1:B:219:GLN:HG2	3:B:557:HOH:O	1.61	1.01
1:A:268:LYS:HD3	3:B:569:HOH:O	1.62	1.00
1:A:224:VAL:HB	3:B:650:HOH:O	0.80	0.98

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:135:GLU:HG3	3:B:670:HOH:O	1.70	0.91
1:A:401:GLU:HG2	1:B:81:TYR:CD1	2.14	0.83
1:B:383:TYR:HE2	3:B:752:HOH:O	1.67	0.76
1:A:401:GLU:HG2	1:B:81:TYR:HD1	1.52	0.74
1:A:383:TYR:HE2	3:B:539:HOH:O	1.71	0.74
1:B:388:TRP:CD1	3:B:625:HOH:O	2.41	0.72
1:A:305:ARG:HD3	3:B:517:HOH:O	1.89	0.72
1:B:388:TRP:NE1	3:B:625:HOH:O	2.23	0.71
1:A:208:VAL:O	1:A:211:ARG:HG2	1.95	0.66
1:A:81:TYR:CD1	1:B:401:GLU:HG2	2.30	0.66
1:B:341:ASN:OD1	1:B:343:ASN:HB2	1.95	0.65
1:A:145:ARG:NH2	1:A:201:LYS:HB3	2.12	0.64
1:A:200:TYR:OH	3:B:758:HOH:O	1.66	0.64
1:B:383:TYR:HD1	3:B:431:HOH:O	1.80	0.63
1:A:296:GLY:HA2	1:A:306:ILE:HD11	1.79	0.63
1:B:384:ARG:O	1:B:387:GLU:HG2	1.99	0.63
1:A:341:ASN:OD1	1:A:343:ASN:HB2	2.00	0.61
1:B:305[A]:ARG:HD3	3:B:562:HOH:O	2.02	0.60
1:B:7:ILE:H	1:B:31:ASN:ND2	1.99	0.60
1:A:268:LYS:CD	3:B:569:HOH:O	2.34	0.59
1:B:341:ASN:ND2	1:B:344:THR:H	2.01	0.58
1:A:342:ASP:HA	1:A:345:ARG:HD2	1.84	0.57
1:A:383:TYR:HD1	3:B:468:HOH:O	1.88	0.56
1:A:341:ASN:ND2	1:A:344:THR:H	2.04	0.56
1:B:282:HIS:ND1	1:B:309:HIS:HD2	2.03	0.56
1:A:341:ASN:HD21	1:A:344:THR:H	1.52	0.56
1:A:143:GLU:HG2	3:B:679:HOH:O	2.06	0.54
1:A:76:MET:CB	1:A:76:MET:SD	2.91	0.54
1:B:305[B]:ARG:HD2	1:B:333:ALA:HB2	1.88	0.53
1:A:312:SER:HB3	1:A:338:ILE:HG21	1.91	0.52
1:A:81:TYR:HD1	1:B:401:GLU:HG2	1.72	0.52
1:A:7:ILE:H	1:A:31:ASN:ND2	2.08	0.51
1:A:85:GLY:HA3	1:B:85:GLY:HA3	1.92	0.51
1:A:75:MET:HG3	1:B:390:GLN:OE1	2.11	0.50
1:A:226:LYS:HD2	1:A:227:TYR:CZ	2.47	0.50
1:A:224:VAL:HG22	1:A:224:VAL:O	2.12	0.49
1:A:208:VAL:O	1:A:211:ARG:CG	2.61	0.49
1:A:383:TYR:CE2	3:B:539:HOH:O	2.54	0.49
1:A:393:ILE:HB	1:A:394:PRO:CD	2.42	0.48
1:A:56:TYR:CE1	1:B:383:TYR:HE1	2.31	0.48
1:B:210:GLU:OE1	1:B:238:PRO:HD3	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:ARG:HD2	1:B:64:ARG:HA	1.86	0.47
1:B:2:SER:HB2	3:B:789:HOH:O	2.15	0.46
1:B:218:VAL:HA	1:B:247:ILE:HG12	1.96	0.46
1:B:83:ARG:NH1	3:B:757:HOH:O	2.49	0.45
1:B:341:ASN:HD21	1:B:344:THR:H	1.63	0.45
1:A:64:ARG:NH2	1:A:75:MET:SD	2.83	0.45
1:A:224:VAL:HG22	1:A:229:PRO:HG3	1.99	0.44
1:B:279:ILE:HG23	1:B:279:ILE:HD12	1.76	0.44
1:B:274:ARG:HD3	3:B:510:HOH:O	2.16	0.44
1:A:224:VAL:CB	3:B:650:HOH:O	1.70	0.44
1:A:128:HIS:CD2	1:A:128:HIS:H	2.34	0.44
1:A:259:GLU:OE2	1:B:81:TYR:OH	2.30	0.43
1:B:209:HIS:NE2	1:B:401:GLU:OE2	2.31	0.42
1:B:248:ARG:HA	1:B:248:ARG:HD2	1.78	0.42
1:A:209:HIS:NE2	1:A:401:GLU:OE2	2.41	0.42
1:A:274:ARG:HD3	3:B:452:HOH:O	2.21	0.41
1:B:43:GLN:HE21	1:B:43:GLN:HB2	1.63	0.41
1:A:390:GLN:OE1	1:B:75:MET:HG3	2.20	0.41
1:B:51:LEU:HD23	1:B:51:LEU:C	2.41	0.41
1:A:264:PRO:HA	1:A:267:TRP:CE2	2.54	0.41
1:A:306:ILE:HB	1:A:332:ALA:HA	2.03	0.40
1:B:20:ARG:HD2	1:B:20:ARG:HA	2.00	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:770:HOH:O	3:B:770:HOH:O[2_555]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/408 (94%)	374 (97%)	10 (3%)	0	100	100
1	B	385/408 (94%)	377 (98%)	8 (2%)	0	100	100
All	All	769/816 (94%)	751 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/357 (95%)	333 (98%)	6 (2%)	66	41
1	B	340/357 (95%)	335 (98%)	5 (2%)	72	50
All	All	679/714 (95%)	668 (98%)	11 (2%)	70	47

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	42	PHE
1	A	189	SER
1	A	211	ARG
1	A	341	ASN
1	A	403	HIS
1	B	31	ASN
1	B	42	PHE
1	B	111	LEU
1	B	341	ASN
1	B	403	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	43	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	67	ASN
1	A	128	HIS
1	A	215	ASN
1	A	219	GLN
1	A	326	ASN
1	A	328	ASN
1	A	386	HIS
1	B	31	ASN
1	B	43	GLN
1	B	67	ASN
1	B	215	ASN
1	B	219	GLN
1	B	309	HIS
1	B	326	ASN
1	B	328	ASN
1	B	341	ASN
1	B	386	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/408 (94%)	-0.46	8 (2%) 67 65	8, 14, 32, 64	0
1	B	388/408 (95%)	-0.47	6 (1%) 76 75	8, 13, 32, 67	0
All	All	775/816 (94%)	-0.46	14 (1%) 71 70	8, 14, 32, 67	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	158	PHE	9.8
1	A	158	PHE	7.6
1	B	404	HIS	6.1
1	A	404	HIS	5.1
1	B	175	GLY	3.7
1	A	403	HIS	3.7
1	A	175	GLY	3.5
1	A	2	SER	2.9
1	B	403	HIS	2.8
1	B	174	GLN	2.8
1	A	383	TYR	2.2
1	A	20	ARG	2.0
1	A	177	TYR	2.0
1	B	20	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MG	A	500	1/1	0.98	0.15	5.05	24,24,24,24	0
2	MG	B	500	1/1	0.99	0.09	1.34	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.