



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:25 AM GMT

PDB ID : 2GZ3
Title : Structure of Aspartate Semialdehyde Dehydrogenase (ASADH) from Streptococcus pneumoniae complexed with NADP and aspartate-semialdehyde
Authors : Faehnle, C.R.; Le Coq, J.; Liu, X.; Viola, R.E.
Deposited on : 2006-05-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>••</div> </div> </div>
1	B	366	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>• 5%</div> </div> </div>
1	C	366	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>13%</div> <div>••</div> </div> </div>
1	D	366	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>10%</div> <div>••</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAP	B	367	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11791 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate beta-semialdehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	357	Total	C	N	O	S	0	0	0
			2720	1720	458	529	13			
1	B	347	Total	C	N	O	S	0	0	0
			2622	1653	443	513	13			
1	C	355	Total	C	N	O	S	0	0	0
			2697	1706	454	524	13			
1	D	357	Total	C	N	O	S	0	0	0
			2720	1719	458	530	13			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	359	LEU	-	CLONING ARTIFACT	UNP Q8DQ00
A	360	GLU	-	CLONING ARTIFACT	UNP Q8DQ00
A	361	HIS	-	EXPRESSION TAG	UNP Q8DQ00
A	362	HIS	-	EXPRESSION TAG	UNP Q8DQ00
A	363	HIS	-	EXPRESSION TAG	UNP Q8DQ00
A	364	HIS	-	EXPRESSION TAG	UNP Q8DQ00
A	365	HIS	-	EXPRESSION TAG	UNP Q8DQ00
A	366	HIS	-	EXPRESSION TAG	UNP Q8DQ00
B	359	LEU	-	CLONING ARTIFACT	UNP Q8DQ00
B	360	GLU	-	CLONING ARTIFACT	UNP Q8DQ00
B	361	HIS	-	EXPRESSION TAG	UNP Q8DQ00
B	362	HIS	-	EXPRESSION TAG	UNP Q8DQ00
B	363	HIS	-	EXPRESSION TAG	UNP Q8DQ00
B	364	HIS	-	EXPRESSION TAG	UNP Q8DQ00
B	365	HIS	-	EXPRESSION TAG	UNP Q8DQ00
B	366	HIS	-	EXPRESSION TAG	UNP Q8DQ00
C	359	LEU	-	CLONING ARTIFACT	UNP Q8DQ00
C	360	GLU	-	CLONING ARTIFACT	UNP Q8DQ00
C	361	HIS	-	EXPRESSION TAG	UNP Q8DQ00
C	362	HIS	-	EXPRESSION TAG	UNP Q8DQ00
C	363	HIS	-	EXPRESSION TAG	UNP Q8DQ00

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	364	HIS	-	EXPRESSION TAG	UNP Q8DQ00
C	365	HIS	-	EXPRESSION TAG	UNP Q8DQ00
C	366	HIS	-	EXPRESSION TAG	UNP Q8DQ00
D	359	LEU	-	CLONING ARTIFACT	UNP Q8DQ00
D	360	GLU	-	CLONING ARTIFACT	UNP Q8DQ00
D	361	HIS	-	EXPRESSION TAG	UNP Q8DQ00
D	362	HIS	-	EXPRESSION TAG	UNP Q8DQ00
D	363	HIS	-	EXPRESSION TAG	UNP Q8DQ00
D	364	HIS	-	EXPRESSION TAG	UNP Q8DQ00
D	365	HIS	-	EXPRESSION TAG	UNP Q8DQ00
D	366	HIS	-	EXPRESSION TAG	UNP Q8DQ00

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is (2R)-2-AMINO-4-OXOBUTANOIC ACID (three-letter code: AS2) (formula: $C_4H_7NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	B	1	Total	C	N	O	0	0
			8	4	1	3		
3	C	1	Total	C	N	O	0	0
			8	4	1	3		
3	D	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	232	Total	O	0	0
			232	232		
4	B	207	Total	O	0	0
			207	207		
4	C	156	Total	O	0	0
			156	156		
4	D	213	Total	O	0	0
			213	213		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aspartate beta-semialdehyde dehydrogenase




- Molecule 1: Aspartate beta-semialdehyde dehydrogenase



- Molecule 1: Aspartate beta-semialdehyde dehydrogenase



- Molecule 1: Aspartate beta-semialdehyde dehydrogenase



M232	S236	S250	M251	M252	S253	K309	E314	M320	S323	R348	E352	L353	K354	K358	LEU	GLU	HIS	HIS	HIS	HIS	HIS	HIS	HIS
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	59.35Å 64.44Å 96.74Å 89.98° 81.06° 82.00°	Depositor
Resolution (Å)	30.01 – 2.10 29.78 – 2.10	Depositor EDS
% Data completeness (in resolution range)	91.7 (30.01-2.10) 91.0 (29.78-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.211 , 0.277 0.211 , 0.274	Depositor DCC
R_{free} test set	3769 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	24.0	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 75008 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11791	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AS2, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/2769	0.64	1/3762 (0.0%)
1	B	0.50	0/2669	0.62	0/3633
1	C	0.49	0/2744	0.61	0/3728
1	D	0.52	0/2769	0.61	0/3763
All	All	0.52	0/10951	0.62	1/14886 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	45	LEU	CA-CB-CG	5.08	126.98	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2720	0	2723	31	0
1	B	2622	0	2593	23	0
1	C	2697	0	2697	32	0
1	D	2720	0	2718	21	0
2	A	48	0	25	4	0
2	B	48	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	48	0	25	1	0
2	D	48	0	25	4	0
3	A	8	0	5	1	0
3	B	8	0	5	0	0
3	C	8	0	5	0	0
3	D	8	0	5	0	0
4	A	232	0	0	9	0
4	B	207	0	0	6	0
4	C	156	0	0	3	0
4	D	213	0	0	5	0
All	All	11791	0	10851	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:367:NAP:H5N	4:A:479:HOH:O	1.50	1.07
1:B:154:TYR:HA	4:B:581:HOH:O	1.70	0.90
1:C:327:LEU:HD22	4:C:552:HOH:O	1.83	0.79
1:C:332:TRP:HB2	4:C:552:HOH:O	1.88	0.72
1:D:352:GLU:HG2	1:D:354:LYS:HE2	1.75	0.69
1:C:7:VAL:HG13	1:C:18:MET:HE3	1.77	0.67
1:A:18:MET:CE	1:A:94:ASN:OD1	2.46	0.63
1:A:300:ARG:HD3	1:A:332:TRP:CE2	2.33	0.62
1:C:250:SER:HB2	4:C:401:HOH:O	1.99	0.62
1:C:73:GLY:HA2	2:C:367:NAP:O3D	2.00	0.61
1:C:219:TYR:O	1:C:223:LYS:HG3	2.01	0.61
1:B:7:VAL:HG13	1:B:18:MET:HE3	1.83	0.61
1:A:29:ILE:HG21	1:A:32:ILE:HG13	1.84	0.59
1:D:18:MET:HE2	4:D:611:HOH:O	2.03	0.58
1:A:18:MET:HE3	1:A:94:ASN:OD1	2.04	0.57
1:B:7:VAL:HG13	1:B:18:MET:CE	2.35	0.57
1:B:262:GLU:HB2	4:B:603:HOH:O	2.04	0.57
1:C:40:SER:HA	1:C:43:LYS:HD3	1.86	0.57
1:C:32:ILE:HG23	1:C:52:ILE:CD1	2.35	0.56
1:A:183:CYS:HB2	4:A:631:HOH:O	2.04	0.56
1:C:7:VAL:HG13	1:C:18:MET:CE	2.35	0.56
1:A:309:LYS:HG3	4:A:596:HOH:O	2.05	0.56
1:C:59:GLU:N	1:C:59:GLU:OE1	2.38	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:ILE:HD12	1:A:32:ILE:N	2.21	0.55
1:C:300:ARG:HD3	1:C:332:TRP:CE2	2.42	0.54
1:A:106:LEU:HD21	1:A:227:GLU:HG2	1.90	0.53
1:C:32:ILE:HG23	1:C:52:ILE:HD13	1.91	0.53
1:D:170:ARG:NH2	1:D:186:HIS:O	2.41	0.53
1:A:215:ASN:O	1:A:216:ASP:HB2	2.09	0.53
1:A:143:LYS:HE2	4:A:414:HOH:O	2.09	0.53
1:C:45:LEU:HD13	1:C:54:ILE:HD11	1.91	0.52
1:D:236:SER:HB2	4:D:533:HOH:O	2.10	0.52
1:A:300:ARG:HD3	1:A:332:TRP:CD2	2.45	0.52
1:B:33:ARG:NH1	1:B:61:ALA:O	2.42	0.52
1:A:220:GLU:HA	1:A:223:LYS:HD2	1.92	0.52
1:C:138:GLU:N	1:C:139:PRO:HD2	2.24	0.52
1:C:213:THR:HG22	1:C:219:TYR:CD1	2.45	0.51
1:B:253:SER:HB3	1:B:320:TRP:CZ2	2.46	0.51
1:D:40:SER:OG	2:D:367:NAP:O3X	2.28	0.51
1:A:339:GLU:O	1:A:343:GLU:HG3	2.11	0.51
1:B:300:ARG:HD3	1:B:332:TRP:CE2	2.45	0.50
1:A:138:GLU:N	1:A:139:PRO:HD2	2.27	0.50
1:A:153:THR:OG1	1:A:243:CYS:HA	2.11	0.50
1:A:33:ARG:NH1	4:A:496:HOH:O	2.43	0.49
1:A:94:ASN:HB3	2:A:367:NAP:H6N	1.95	0.49
1:D:94:ASN:O	2:D:367:NAP:H6N	2.13	0.49
1:C:154:TYR:HB2	1:C:253:SER:HB2	1.94	0.49
1:C:253:SER:HB3	1:C:320:TRP:CZ2	2.47	0.49
1:A:212:PHE:HA	1:A:218:THR:HA	1.95	0.48
1:D:215:ASN:O	1:D:216:ASP:HB2	2.13	0.48
1:A:2:GLY:N	4:A:626:HOH:O	2.46	0.48
1:D:138:GLU:N	1:D:139:PRO:HD2	2.28	0.47
1:A:249:LEU:HB2	4:A:432:HOH:O	2.15	0.47
1:D:173:ARG:O	1:D:177:ASN:HB2	2.14	0.47
1:C:106:LEU:HD21	1:C:227:GLU:HG2	1.96	0.47
1:B:90:VAL:HG11	1:B:341:LEU:HD13	1.96	0.47
1:A:79:LYS:HG3	4:A:512:HOH:O	2.14	0.47
1:C:20:LYS:O	1:C:24:GLU:HG2	2.15	0.47
1:D:18:MET:SD	1:D:69:LEU:HD13	2.55	0.46
1:D:188:GLU:HG3	4:D:437:HOH:O	2.15	0.46
1:B:266:ILE:HG22	1:B:270:LYS:HE3	1.97	0.46
1:D:49:ASP:HB2	4:D:480:HOH:O	2.15	0.46
1:C:213:THR:HG22	1:C:219:TYR:CE1	2.51	0.46
1:D:20:LYS:HA	1:D:20:LYS:HD2	1.71	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:98:PHE:HB3	1:C:104:VAL:HG21	1.98	0.45
1:D:253:SER:HB3	1:D:320:TRP:CZ2	2.51	0.45
1:B:141:ARG:HB2	1:B:146:LEU:HD12	1.99	0.45
1:B:173:ARG:O	1:B:177:ASN:HB2	2.16	0.45
1:C:123:ILE:HD12	1:C:123:ILE:N	2.31	0.45
1:A:330:ALA:HB2	2:A:367:NAP:C4N	2.47	0.44
1:B:169:GLN:HE21	1:B:249:LEU:HD11	1.82	0.44
1:A:127:ASN:HB3	1:A:227:GLU:OE1	2.17	0.44
1:C:261:LYS:O	1:C:315:LYS:NZ	2.48	0.44
1:A:253:SER:HB3	1:A:320:TRP:CZ2	2.51	0.44
1:D:159:GLY:HA2	2:D:367:NAP:N7N	2.32	0.44
1:D:32:ILE:HG22	4:D:595:HOH:O	2.17	0.44
1:B:114:HIS:HD2	4:B:604:HOH:O	1.99	0.44
1:B:250:SER:CB	4:B:480:HOH:O	2.66	0.44
1:B:276:PHE:HA	1:B:277:PRO:HD3	1.86	0.44
1:B:244:VAL:HB	4:B:581:HOH:O	2.17	0.44
1:D:175:VAL:HG11	1:D:203:PHE:CE2	2.53	0.43
1:A:46:LYS:HD2	1:A:48:LYS:O	2.19	0.43
1:B:280:VAL:HG23	1:B:301:ASP:HB3	2.01	0.43
1:B:250:SER:HB3	4:B:480:HOH:O	2.18	0.43
1:A:276:PHE:HA	1:A:277:PRO:HD3	1.87	0.43
1:C:250:SER:O	1:C:251:ALA:HB3	2.19	0.42
1:B:328:LYS:O	1:B:333:ASN:HB2	2.18	0.42
1:A:18:MET:HE1	1:A:94:ASN:OD1	2.17	0.42
1:C:7:VAL:HG22	1:C:69:LEU:HD12	2.02	0.42
1:C:165:ILE:HG23	1:C:249:LEU:HD22	2.00	0.42
1:D:250:SER:O	1:D:251:ALA:HB3	2.19	0.42
1:D:73:GLY:HA2	2:D:367:NAP:O3D	2.19	0.42
1:B:211:VAL:HG23	1:B:219:TYR:HD1	1.85	0.42
1:A:182:PRO:HA	1:A:185:LEU:HD12	2.02	0.41
1:C:212:PHE:HA	1:C:218:THR:HA	2.02	0.41
1:C:170:ARG:NH2	1:C:186:HIS:O	2.53	0.41
1:C:206:LEU:HA	1:C:207:PRO:HD2	1.91	0.41
1:B:149:ILE:HG12	1:B:258:ILE:HG22	2.02	0.41
1:B:74:SER:HB2	1:B:97:TYR:HB2	2.02	0.41
1:C:32:ILE:HG23	1:C:52:ILE:HD11	2.03	0.41
1:A:2:GLY:HA2	4:A:626:HOH:O	2.20	0.41
1:D:141:ARG:NH1	1:D:232:MET:HB3	2.36	0.41
1:A:125:CYS:HA	1:A:126:PRO:HD3	1.97	0.41
1:C:276:PHE:HA	1:C:277:PRO:HD3	1.94	0.41
2:A:367:NAP:C7N	3:A:400:AS2:HBA	2.51	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:98:PHE:HB3	1:D:104:VAL:HG21	2.03	0.40
1:A:154:TYR:CD2	1:A:154:TYR:N	2.90	0.40
1:C:43:LYS:HE2	1:C:43:LYS:HB3	1.90	0.40
1:B:154:TYR:HB2	1:B:253:SER:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/366 (97%)	342 (96%)	11 (3%)	2 (1%)	30	24
1	B	345/366 (94%)	329 (95%)	14 (4%)	2 (1%)	30	24
1	C	351/366 (96%)	337 (96%)	12 (3%)	2 (1%)	30	24
1	D	355/366 (97%)	336 (95%)	18 (5%)	1 (0%)	46	45
All	All	1406/1464 (96%)	1344 (96%)	55 (4%)	7 (0%)	34	30

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	ALA
1	B	251	ALA
1	C	251	ALA
1	D	251	ALA
1	B	330	ALA
1	C	330	ALA
1	A	330	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/302 (97%)	282 (97%)	10 (3%)	44	45
1	B	279/302 (92%)	270 (97%)	9 (3%)	46	48
1	C	289/302 (96%)	277 (96%)	12 (4%)	36	35
1	D	292/302 (97%)	278 (95%)	14 (5%)	31	29
All	All	1152/1208 (95%)	1107 (96%)	45 (4%)	39	39

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	LYS
1	A	74	SER
1	A	129	SER
1	A	154	TYR
1	A	170	ARG
1	A	173	ARG
1	A	201	ILE
1	A	210	ASP
1	A	348	ARG
1	A	352	GLU
1	B	18	MET
1	B	37	SER
1	B	45	LEU
1	B	59	GLU
1	B	63	GLU
1	B	170	ARG
1	B	250	SER
1	B	262	GLU
1	B	348	ARG
1	C	18	MET
1	C	49	ASP
1	C	50	GLN
1	C	69	LEU
1	C	94	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	129	SER
1	C	170	ARG
1	C	186	HIS
1	C	267	GLU
1	C	290	TYR
1	C	348	ARG
1	C	350	THR
1	D	20	LYS
1	D	48	LYS
1	D	51	ASP
1	D	63	GLU
1	D	94	ASN
1	D	129	SER
1	D	170	ARG
1	D	201	ILE
1	D	214	ASP
1	D	309	LYS
1	D	314	GLU
1	D	323	SER
1	D	348	ARG
1	D	358	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	198	HIS
1	A	292	GLN
1	B	94	ASN
1	B	114	HIS
1	B	336	GLN
1	C	198	HIS
1	D	114	HIS
1	D	198	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAP	A	367	-	42,52,52	1.73	4 (9%)	54,80,80	2.13	10 (18%)
3	AS2	A	400	1	4,7,7	0.70	0	1,8,8	1.04	0
2	NAP	B	367	-	42,52,52	1.93	8 (19%)	54,80,80	1.90	7 (12%)
3	AS2	B	400	1	4,7,7	0.54	0	1,8,8	1.20	0
2	NAP	C	367	-	42,52,52	1.83	5 (11%)	54,80,80	1.91	6 (11%)
3	AS2	C	400	1	4,7,7	0.75	0	1,8,8	1.11	0
2	NAP	D	367	-	42,52,52	1.87	8 (19%)	54,80,80	2.07	14 (25%)
3	AS2	D	400	1	4,7,7	0.63	0	1,8,8	1.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	367	-	-	0/27/67/67	0/5/5/5
3	AS2	A	400	1	-	0/3/7/7	0/0/0/0
2	NAP	B	367	-	-	0/27/67/67	0/5/5/5
3	AS2	B	400	1	-	0/3/7/7	0/0/0/0
2	NAP	C	367	-	-	0/27/67/67	0/5/5/5
3	AS2	C	400	1	-	0/3/7/7	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	D	367	-	-	0/27/67/67	0/5/5/5
3	AS2	D	400	1	-	0/3/7/7	0/0/0/0

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	367	NAP	P2B-O3X	-2.14	1.47	1.54
2	B	367	NAP	P2B-O3X	-2.11	1.47	1.54
2	B	367	NAP	O4B-C1B	2.11	1.43	1.41
2	B	367	NAP	P2B-O2B	2.14	1.66	1.60
2	B	367	NAP	C3N-C7N	2.31	1.54	1.50
2	D	367	NAP	P2B-O2X	2.34	1.63	1.54
2	C	367	NAP	O4B-C1B	2.47	1.44	1.41
2	D	367	NAP	P2B-O1X	2.62	1.59	1.51
2	D	367	NAP	C3N-C7N	2.72	1.54	1.50
2	B	367	NAP	P2B-O1X	2.76	1.60	1.51
2	D	367	NAP	O4B-C1B	2.76	1.44	1.41
2	A	367	NAP	P2B-O1X	2.79	1.60	1.51
2	C	367	NAP	P2B-O1X	2.93	1.60	1.51
2	A	367	NAP	O4D-C1D	3.51	1.45	1.41
2	D	367	NAP	O4D-C1D	3.88	1.46	1.41
2	C	367	NAP	O4D-C1D	4.18	1.46	1.41
2	A	367	NAP	C6N-N1N	5.19	1.49	1.35
2	B	367	NAP	C6N-N1N	5.27	1.49	1.35
2	C	367	NAP	C6N-N1N	5.34	1.49	1.35
2	B	367	NAP	O4D-C1D	5.44	1.48	1.41
2	D	367	NAP	C6N-N1N	5.52	1.50	1.35
2	A	367	NAP	C6N-C5N	6.90	1.54	1.38
2	B	367	NAP	C6N-C5N	7.10	1.54	1.38
2	D	367	NAP	C6N-C5N	7.24	1.54	1.38
2	C	367	NAP	C6N-C5N	7.36	1.55	1.38

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	367	NAP	N3A-C2A-N1A	-9.53	121.60	128.89
2	B	367	NAP	N3A-C2A-N1A	-9.09	121.93	128.89
2	A	367	NAP	N3A-C2A-N1A	-8.97	122.03	128.89
2	D	367	NAP	N3A-C2A-N1A	-8.92	122.06	128.89
2	A	367	NAP	C3N-C7N-N7N	-4.07	113.36	117.82
2	D	367	NAP	C6N-C5N-C4N	-3.47	114.21	119.44
2	C	367	NAP	C5N-C6N-N1N	-3.42	114.55	120.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	367	NAP	C3N-C7N-N7N	-3.22	114.29	117.82
2	A	367	NAP	C5N-C6N-N1N	-3.08	115.15	120.47
2	D	367	NAP	C3N-C7N-N7N	-3.03	114.50	117.82
2	A	367	NAP	C4D-O4D-C1D	-2.98	106.44	109.72
2	B	367	NAP	C5N-C6N-N1N	-2.96	115.34	120.47
2	B	367	NAP	C6N-C5N-C4N	-2.92	115.04	119.44
2	D	367	NAP	C5N-C6N-N1N	-2.84	115.55	120.47
2	A	367	NAP	C6N-C5N-C4N	-2.83	115.17	119.44
2	D	367	NAP	O2X-P2B-O1X	-2.52	102.47	110.58
2	C	367	NAP	C6N-C5N-C4N	-2.52	115.64	119.44
2	C	367	NAP	PN-O3-PA	-2.34	126.16	132.73
2	D	367	NAP	C4A-C5A-N7A	-2.29	107.37	109.48
2	B	367	NAP	O2X-P2B-O1X	-2.23	103.40	110.58
2	D	367	NAP	C4B-O4B-C1B	-2.22	107.28	109.72
2	D	367	NAP	C4D-O4D-C1D	-2.18	107.32	109.72
2	D	367	NAP	C1B-N9A-C4A	-2.12	123.74	126.94
2	D	367	NAP	O4B-C4B-C5B	-2.06	101.94	109.32
2	A	367	NAP	O4B-C1B-N9A	2.05	112.39	108.10
2	A	367	NAP	O7N-C7N-N7N	2.11	125.57	122.59
2	B	367	NAP	O4D-C1D-N1N	2.12	110.46	108.13
2	D	367	NAP	O7N-C7N-C3N	2.18	121.97	119.59
2	A	367	NAP	O3X-P2B-O2X	2.36	116.36	107.38
2	D	367	NAP	O3X-P2B-O2X	2.50	116.91	107.38
2	D	367	NAP	O4D-C1D-N1N	2.58	110.96	108.13
2	C	367	NAP	O4D-C1D-N1N	3.09	111.53	108.13
2	C	367	NAP	C5N-C4N-C3N	5.48	127.23	120.33
2	A	367	NAP	O4D-C1D-N1N	5.49	114.17	108.13
2	A	367	NAP	C5N-C4N-C3N	5.56	127.32	120.33
2	B	367	NAP	C5N-C4N-C3N	5.94	127.80	120.33
2	D	367	NAP	C5N-C4N-C3N	6.59	128.62	120.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	367	NAP	4	0
3	A	400	AS2	1	0
2	C	367	NAP	1	0
2	D	367	NAP	4	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/366 (97%)	0.05	5 (1%) 78 82	11, 18, 28, 34	0
1	B	347/366 (94%)	0.22	10 (2%) 55 63	10, 21, 32, 42	0
1	C	355/366 (96%)	0.26	6 (1%) 73 78	14, 24, 37, 50	0
1	D	357/366 (97%)	0.17	4 (1%) 82 86	11, 21, 35, 45	0
All	All	1416/1464 (96%)	0.18	25 (1%) 71 76	10, 21, 34, 50	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	219	TYR	4.2
1	D	44	SER	4.0
1	A	219	TYR	3.4
1	A	186	HIS	3.3
1	B	49	ASP	3.3
1	A	49	ASP	3.3
1	D	49	ASP	3.3
1	B	83	TYR	3.0
1	C	235	ASP	2.8
1	B	256	VAL	2.8
1	C	353	LEU	2.7
1	B	60	THR	2.6
1	B	246	ILE	2.6
1	C	49	ASP	2.6
1	A	92	VAL	2.5
1	B	321	VAL	2.4
1	C	193	GLY	2.4
1	B	186	HIS	2.3
1	B	154	TYR	2.3
1	B	255	SER	2.3
1	A	194	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	198	HIS	2.1
1	B	286	ALA	2.1
1	D	51	ASP	2.1
1	C	186	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAP	B	367	48/48	0.90	0.18	2.47	28,40,59,59	0
2	NAP	D	367	48/48	0.86	0.21	1.66	30,41,63,64	0
2	NAP	A	367	48/48	0.89	0.17	1.63	23,32,52,52	0
2	NAP	C	367	48/48	0.88	0.18	1.47	28,41,56,58	0
3	AS2	D	400	8/8	0.91	0.16	0.67	22,22,23,24	0
3	AS2	B	400	8/8	0.90	0.14	0.54	24,24,25,26	0
3	AS2	A	400	8/8	0.91	0.14	0.51	21,23,23,24	0
3	AS2	C	400	8/8	0.93	0.13	-0.02	25,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.