



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:02 AM GMT

PDB ID : 3GZS
Title : Crystal structure of a susd superfamily protein (bf3413) from bacteroides fragilis nctc 9343 at 2.10 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2009-04-07
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	<div> <div>15%</div> <div>92%</div> <div>5%</div> <div>.</div> </div>
1	B	515	<div> <div>13%</div> <div>90%</div> <div>5%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	1[A]	-	-	-	X
2	GOL	A	1[B]	-	-	-	X
2	GOL	A	5	-	-	-	X
2	GOL	A	6	-	-	-	X
2	GOL	B	2	-	-	-	X
2	GOL	B	3	-	-	-	X
3	ACT	A	11	-	-	-	X
3	ACT	B	9	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8328 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized SusD superfamily protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	495	Total	C	N	O	S	Se	0	2	0
			3879	2457	659	748	2	13			
1	B	495	Total	C	N	O	S	Se	0	3	0
			3881	2457	660	749	2	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q5L9X2
B	0	GLY	-	LEADER SEQUENCE	UNP Q5L9X2

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	1
			7	3	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	243	Total	O	0	0
			243	243		
4	B	260	Total	O	0	0
			260	260		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized SusD superfamily protein



- Molecule 1: Uncharacterized SusD superfamily protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.92Å 50.38Å 117.95Å 90.00° 108.05° 90.00°	Depositor
Resolution (Å)	29.63 – 2.09 29.63 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.3 (29.63-2.09) 97.3 (29.63-2.09)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.156 , 0.198 0.166 , 0.207	Depositor DCC
R_{free} test set	2934 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 57418 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8328	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.67 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.2811e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OCS, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	2/3953 (0.1%)	0.58	2/5337 (0.0%)
1	B	0.65	1/3958 (0.0%)	0.58	2/5345 (0.0%)
All	All	0.64	3/7911 (0.0%)	0.58	4/10682 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	73	GLU	CB-CG	5.92	1.63	1.52
1	A	73	GLU	CB-CG	5.35	1.62	1.52
1	A	111	MSE	SE-CE	-5.15	1.65	1.95

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	244	ARG	NE-CZ-NH2	-5.61	117.50	120.30
1	A	244	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	B	244	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	244	ARG	NE-CZ-NH2	-5.34	117.63	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3879	0	3681	13	0
1	B	3881	0	3681	20	0
2	A	25	0	30	0	0
2	B	12	0	16	0	0
3	A	12	0	9	0	0
3	B	16	0	12	2	0
4	A	243	0	0	0	0
4	B	260	0	0	1	0
All	All	8328	0	7429	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:72:THR:HG23	1:B:296:TYR:OH	1.81	0.80
1:A:72:THR:HG23	1:A:296:TYR:OH	1.97	0.65
1:B:202:VAL:HG13	1:B:231:MSE:HE3	1.82	0.62
1:A:166:MSE:SE	1:A:503:LEU:HD13	2.54	0.57
1:B:215[A]:VAL:HG23	1:B:227:VAL:HG22	1.88	0.55
1:B:454:LYS:NZ	1:B:468:GLU:OE2	2.29	0.54
1:B:166:MSE:SE	1:B:503:LEU:HD13	2.60	0.52
1:A:120:TYR:CG	1:A:506:PRO:HG3	2.46	0.51
1:A:244:ARG:NH2	1:A:464:GLU:OE2	2.44	0.50
1:A:202:VAL:HG13	1:A:231:MSE:HE3	1.93	0.50
1:B:276:GLN:HB3	1:B:356:TYR:CD1	2.48	0.49
1:A:123:MSE:SE	1:A:166:MSE:HE3	2.63	0.48
1:B:215[A]:VAL:HG23	1:B:227:VAL:CG2	2.43	0.48
1:A:508:THR:O	1:A:512:GLN:HG2	2.13	0.47
1:A:231:MSE:HE2	1:A:265:ALA:CB	2.45	0.47
1:B:446:LYS:NZ	3:B:9:ACT:H1	2.30	0.46
1:A:231:MSE:HE2	1:A:265:ALA:HB2	1.97	0.46
1:B:243:ILE:O	1:B:246:ARG:HG3	2.17	0.45
1:A:243:ILE:O	1:A:246:ARG:HG3	2.17	0.45
1:B:244:ARG:NH2	1:B:464:GLU:OE2	2.49	0.44
1:A:139:GLN:O	1:B:343:GLY:HA3	2.17	0.44
1:B:206:ASP:OD2	1:B:231:MSE:HE1	2.19	0.43
1:A:72:THR:H	1:A:75:VAL:HG23	1.83	0.43
1:B:202:VAL:HG13	1:B:231:MSE:CE	2.46	0.43
1:B:301:MSE:HE2	1:B:306:TYR:HA	2.01	0.42
1:B:312:TYR:HB3	1:B:435:ILE:HD11	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:109:TYR:HA	1:B:111:MSE:CE	2.50	0.41
1:B:467:THR:HG23	1:B:537:TRP:CZ2	2.55	0.41
1:B:359:LYS:NZ	4:B:853:HOH:O	2.53	0.41
1:B:109:TYR:O	1:B:111:MSE:HE3	2.22	0.40
1:B:446:LYS:HZ2	3:B:9:ACT:H1	1.86	0.40
1:A:505:TYR:CD2	1:A:517:LEU:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	494/515 (96%)	478 (97%)	16 (3%)	0	100	100
1	B	495/515 (96%)	477 (96%)	18 (4%)	0	100	100
All	All	989/1030 (96%)	955 (97%)	34 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/409 (95%)	389 (100%)	0	100	100
1	B	390/409 (95%)	390 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	779/818 (95%)	779 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	428	ASN
1	B	210	ASN
1	B	428	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	OCS	A	364	1	7,8,9	1.28	1 (14%)	7,11,13	1.70	3 (42%)
1	OCS	B	364	1	7,8,9	0.89	0	7,11,13	1.53	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	A	364	1	-	0/4/7/9	0/0/0/0
1	OCS	B	364	1	-	0/4/7/9	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	364	OCS	CB-SG	2.82	1.82	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	364	OCS	O-C-CA	-2.10	120.01	125.49
1	B	364	OCS	O-C-CA	-2.09	120.06	125.49
1	A	364	OCS	OD1-SG-CB	2.43	108.99	106.94
1	A	364	OCS	OD3-SG-CB	2.62	109.15	106.94
1	B	364	OCS	OD1-SG-CB	2.91	109.40	106.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	1[A]	-	5,5,5	0.65	0	5,5,5	1.61	1 (20%)
2	GOL	A	1[B]	-	5,5,5	0.65	0	5,5,5	1.49	1 (20%)
3	ACT	A	11	-	1,3,3	1.61	0	0,3,3	0.00	-
3	ACT	A	13	-	1,3,3	1.73	0	0,3,3	0.00	-
2	GOL	A	4	-	5,5,5	0.50	0	5,5,5	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	5	-	5,5,5	0.45	0	5,5,5	0.25	0
2	GOL	A	6	-	5,5,5	0.32	0	5,5,5	0.35	0
3	ACT	A	8	-	1,3,3	1.22	0	0,3,3	0.00	-
3	ACT	B	10	-	1,3,3	1.77	0	0,3,3	0.00	-
3	ACT	B	12	-	1,3,3	1.49	0	0,3,3	0.00	-
2	GOL	B	2	-	5,5,5	0.37	0	5,5,5	0.24	0
2	GOL	B	3	-	5,5,5	0.25	0	5,5,5	0.35	0
3	ACT	B	7	-	1,3,3	2.73	1 (100%)	0,3,3	0.00	-
3	ACT	B	9	-	1,3,3	1.97	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1[A]	-	-	0/4/4/4	0/0/0/0
2	GOL	A	1[B]	-	-	0/4/4/4	0/0/0/0
3	ACT	A	11	-	-	0/0/0/0	0/0/0/0
3	ACT	A	13	-	-	0/0/0/0	0/0/0/0
2	GOL	A	4	-	-	0/4/4/4	0/0/0/0
2	GOL	A	5	-	-	0/4/4/4	0/0/0/0
2	GOL	A	6	-	-	0/4/4/4	0/0/0/0
3	ACT	A	8	-	-	0/0/0/0	0/0/0/0
3	ACT	B	10	-	-	0/0/0/0	0/0/0/0
3	ACT	B	12	-	-	0/0/0/0	0/0/0/0
2	GOL	B	2	-	-	0/4/4/4	0/0/0/0
2	GOL	B	3	-	-	0/4/4/4	0/0/0/0
3	ACT	B	7	-	-	0/0/0/0	0/0/0/0
3	ACT	B	9	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	7	ACT	CH3-C	2.73	1.52	1.48

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1[A]	GOL	O1-C1-C2	-3.48	93.33	110.18
2	A	1[B]	GOL	O1-C1-C2	-3.20	94.68	110.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	9	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	481/515 (93%)	0.91	75 (15%) 3 4	43, 50, 61, 86	0
1	B	481/515 (93%)	0.81	69 (14%) 4 5	43, 50, 62, 88	0
All	All	962/1030 (93%)	0.86	144 (14%) 3 5	43, 50, 62, 88	0

All (144) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	VAL	10.7
1	B	178	GLY	10.2
1	A	48	GLY	9.5
1	B	179	SER	7.0
1	B	142	GLY	6.0
1	B	49	VAL	5.7
1	A	411	THR	5.6
1	A	179	SER	5.4
1	B	465	ALA	5.1
1	A	401	SER	5.1
1	A	451	ILE	5.1
1	A	465	ALA	5.0
1	B	180	PHE	4.9
1	A	92	TYR	4.8
1	B	411	THR	4.8
1	A	458	ILE	4.7
1	A	405	THR	4.7
1	A	178	GLY	4.6
1	A	450	ILE	4.6
1	A	455	TYR	4.5
1	B	455	TYR	4.5
1	A	52	GLY	4.4
1	A	142	GLY	4.4
1	A	467	THR	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	141	GLU	4.4
1	A	440	GLY	4.3
1	A	180	PHE	4.3
1	B	264	PRO	4.2
1	B	177	LYS	4.1
1	A	237	LEU	4.1
1	B	451	ILE	4.0
1	B	143	THR	4.0
1	B	462	GLY	4.0
1	A	441	ALA	4.0
1	A	454	LYS	3.9
1	A	251	GLY	3.8
1	B	158	ALA	3.8
1	B	458	ILE	3.8
1	B	452	THR	3.8
1	A	210	ASN	3.8
1	A	82	VAL	3.7
1	A	305	ILE	3.7
1	B	50	ILE	3.7
1	A	53	ALA	3.7
1	B	72	THR	3.6
1	A	72	THR	3.6
1	B	365	PHE	3.5
1	B	159	ALA	3.5
1	A	402	SER	3.4
1	B	473	GLY	3.4
1	B	212	SER	3.4
1	B	541	SER	3.3
1	A	141	GLU	3.3
1	A	386	ASN	3.3
1	A	219	PHE	3.3
1	B	459	PHE	3.3
1	B	137	ILE	3.2
1	B	88	ALA	3.2
1	A	368	ALA	3.2
1	B	443	GLU	3.2
1	B	136	ARG	3.2
1	A	464	GLU	3.2
1	B	401	SER	3.2
1	A	62	VAL	3.2
1	A	212	SER	3.1
1	A	397	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	467	THR	3.1
1	A	461	ASN	3.0
1	B	400	ASN	3.0
1	A	90	ALA	3.0
1	B	466	TRP	3.0
1	A	457	ALA	3.0
1	A	442	THR	2.9
1	B	48	GLY	2.9
1	A	159	ALA	2.9
1	B	475	PRO	2.9
1	B	181	LYS	2.8
1	A	282	GLY	2.8
1	A	253	ALA	2.8
1	A	308	TYR	2.8
1	A	475	PRO	2.8
1	A	50	ILE	2.8
1	A	463	GLN	2.8
1	A	375	PHE	2.8
1	A	466	TRP	2.8
1	B	62	VAL	2.7
1	A	474	TYR	2.7
1	B	162[A]	ARG	2.7
1	B	155	ILE	2.7
1	A	472	THR	2.7
1	A	158	ALA	2.7
1	B	399	ASP	2.6
1	A	209	SER	2.6
1	A	400	ASN	2.6
1	A	468	GLU	2.6
1	A	424	GLU	2.6
1	A	206	ASP	2.6
1	B	269	ASN	2.6
1	A	350	HIS	2.5
1	B	82	VAL	2.5
1	B	207	ASP	2.5
1	A	363	VAL	2.5
1	B	450	ILE	2.5
1	A	460	PRO	2.5
1	B	324	ASN	2.4
1	A	438	GLU	2.4
1	B	426	SER	2.4
1	B	144	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	122	PHE	2.4
1	B	469	TRP	2.4
1	B	456	LEU	2.4
1	B	461	ASN	2.3
1	A	393	PHE	2.3
1	B	460	PRO	2.3
1	A	409	ASP	2.3
1	B	216	LEU	2.3
1	B	92	TYR	2.3
1	B	463	GLN	2.3
1	B	219	PHE	2.3
1	A	469	TRP	2.3
1	A	262	LYS	2.3
1	B	140	ASP	2.3
1	B	437	TRP	2.3
1	A	177	LYS	2.2
1	B	157	ILE	2.2
1	A	462	GLY	2.2
1	A	459	PHE	2.2
1	B	176	GLY	2.2
1	A	250	ALA	2.2
1	A	157	ILE	2.2
1	B	160	LEU	2.1
1	B	403	ALA	2.1
1	A	215	VAL	2.1
1	B	357	TRP	2.1
1	A	410	SER	2.1
1	B	398	LEU	2.1
1	A	439	ASN	2.1
1	B	378	GLY	2.1
1	B	438	GLU	2.1
1	B	167	PHE	2.1
1	A	199	GLU	2.0
1	A	374	GLY	2.0
1	B	90	ALA	2.0
1	B	255	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	OCS	A	364	9/10	0.94	0.12	-	51,52,60,61	0
1	OCS	B	364	9/10	0.95	0.11	-	50,51,60,64	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ACT	A	11	4/4	0.62	0.24	9.94	82,83,84,84	0
2	GOL	A	6	6/6	0.84	0.31	8.65	57,70,72,73	0
2	GOL	B	3	6/6	0.91	0.26	6.83	46,61,70,70	0
2	GOL	A	5	6/6	0.73	0.38	5.89	73,76,79,81	0
2	GOL	A	1[A]	6/6	0.60	0.35	4.76	53,71,77,79	1
2	GOL	A	1[B]	6/6	0.60	0.35	4.76	59,71,77,79	1
2	GOL	B	2	6/6	0.90	0.33	3.32	45,55,64,68	0
3	ACT	B	9	4/4	0.84	0.21	0.61	66,66,67,68	0
2	GOL	A	4	6/6	0.94	0.15	0.38	49,55,68,74	0
3	ACT	B	7	4/4	0.98	0.10	0.14	36,36,38,40	0
3	ACT	A	8	4/4	0.95	0.23	0.11	49,51,54,56	0
3	ACT	A	13	4/4	0.98	0.10	-1.30	43,43,45,46	0
3	ACT	B	10	4/4	0.94	0.15	-1.47	44,52,52,52	0
3	ACT	B	12	4/4	0.86	0.15	-	68,73,74,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.