



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:03 PM GMT

PDB ID : 4GZS  
Title : N2 neuraminidase D151G mutant of a/Tanzania/205/2010 H3N2 in complex with hepes  
Authors : Zhu, X.; Wilson, I.A.  
Deposited on : 2012-09-06  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	<div> <div>4%</div> <div>83% 15% ..</div> </div>
1	B	393	<div> <div>6%</div> <div>82% 16% ..</div> </div>
1	C	393	<div> <div>6%</div> <div>80% 16% ..</div> </div>
1	D	393	<div> <div>2%</div> <div>80% 18% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EPE	A	502	-	-	-	X
3	EPE	C	503	-	-	-	X
3	EPE	D	502	-	-	-	X
4	NAG	A	503	-	-	-	X
4	NAG	B	503	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	0	0	0
			2998	1858	531	587	22			
1	B	388	Total	C	N	O	S	0	0	0
			2998	1858	531	587	22			
1	C	388	Total	C	N	O	S	0	0	0
			2998	1858	531	587	22			
1	D	388	Total	C	N	O	S	0	0	0
			2998	1858	531	587	22			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		
2	D	1	Total	Ca	0	0
			1	1		
2	C	2	Total	Ca	0	0
			2	2		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	D	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 4 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	5	Total	C	N	O	0	0
			61	34	2	25		
4	B	5	Total	C	N	O	0	0
			61	34	2	25		
4	C	5	Total	C	N	O	0	0
			61	34	2	25		
4	D	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	2	Total	C	N	O	0	0
			24	14	1	9		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	2	Total	C	N	O	0	0
			24	14	1	9		
5	C	2	Total	C	N	O	0	0
			24	14	1	9		

- Molecule 6 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	C	1	Total	C	N	O	0	0
			14	8	1	5		
6	C	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	91	Total	O	0	0
			91	91		
8	B	89	Total	O	0	0
			89	89		
8	C	93	Total	O	0	0
			93	93		
8	D	90	Total	O	0	0
			90	90		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Neuraminidase




#### • Molecule 1: Neuraminidase





#### • Molecule 1: Neuraminidase







● Molecule 1: Neuraminidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.81Å 110.20Å 110.59Å 90.00° 97.46° 90.00°	Depositor
Resolution (Å)	50.00 – 2.35 44.21 – 2.34	Depositor EDS
% Data completeness (in resolution range)	92.4 (50.00-2.35) 91.8 (44.21-2.34)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.02 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.225 , 0.287 0.225 , 0.283	Depositor DCC
$R_{free}$ test set	3706 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.4	Xtriage
Anisotropy	1.073	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 33.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 74279 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	12862	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CA, FUC, EPE, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.56	0/3065	0.70	1/4156 (0.0%)
1	B	0.57	0/3065	0.68	0/4156
1	C	0.59	0/3065	0.71	1/4156 (0.0%)
1	D	0.58	0/3065	0.70	1/4156 (0.0%)
All	All	0.58	0/12260	0.70	3/16624 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	VAL	CB-CA-C	-7.38	97.37	111.40
1	C	216	VAL	CB-CA-C	-5.48	100.98	111.40
1	D	216	VAL	CB-CA-C	-5.43	101.09	111.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2998	0	2856	37	0
1	B	2998	0	2855	43	0
1	C	2998	0	2855	50	0
1	D	2998	0	2855	42	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	1	0	0	0	0
3	A	15	0	17	2	0
3	B	15	0	17	2	0
3	C	15	0	17	1	0
3	D	15	0	17	2	0
4	A	61	0	52	0	0
4	B	61	0	52	1	0
4	C	61	0	52	0	0
4	D	61	0	52	2	0
5	A	24	0	22	1	0
5	B	24	0	22	0	0
5	C	24	0	22	0	0
6	A	14	0	13	0	0
6	B	28	0	26	0	0
6	C	28	0	26	0	0
6	D	28	0	26	0	0
7	D	28	0	25	0	0
8	A	91	0	0	1	0
8	B	89	0	0	2	0
8	C	93	0	0	0	0
8	D	90	0	0	3	0
All	All	12862	0	11879	160	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:THR:HB	1:C:216:VAL:CG2	2.02	0.88
1:A:177:ALA:HB2	1:A:193:CYS:HB3	1.60	0.84
1:C:452:THR:HB	1:D:216:VAL:HG22	1.68	0.73
1:B:452:THR:HB	1:C:216:VAL:HG22	1.69	0.73
1:A:216:VAL:HG13	1:D:453:TYR:C	2.13	0.69

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/393 (98%)	361 (94%)	21 (5%)	4 (1%)	19	20
1	B	386/393 (98%)	358 (93%)	24 (6%)	4 (1%)	19	20
1	C	386/393 (98%)	354 (92%)	27 (7%)	5 (1%)	15	13
1	D	386/393 (98%)	357 (92%)	25 (6%)	4 (1%)	19	20
All	All	1544/1572 (98%)	1430 (93%)	97 (6%)	17 (1%)	17	17

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	468	PRO
1	B	333	SER
1	C	234	ASN
1	D	468	PRO
1	A	319	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/341 (99%)	323 (96%)	14 (4%)	36	46
1	B	337/341 (99%)	328 (97%)	9 (3%)	52	67
1	C	337/341 (99%)	324 (96%)	13 (4%)	39	51
1	D	337/341 (99%)	324 (96%)	13 (4%)	39	51
All	All	1348/1364 (99%)	1299 (96%)	49 (4%)	42	55

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	463	ASP
1	C	267	THR
1	D	338	LEU
1	C	211	LEU
1	C	276	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	136	GLN
1	C	168	HIS
1	D	168	HIS
1	B	393	ASN
1	D	385	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

28 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	503	1,4	14,14,15	0.63	0	15,19,21	1.26	2 (13%)
4	NAG	A	504	4	14,14,15	0.40	0	15,19,21	1.38	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	BMA	A	505	4	11,11,12	0.69	0	14,15,17	0.82	0
4	MAN	A	506	4	11,11,12	0.66	0	14,15,17	1.64	3 (21%)
4	MAN	A	507	4	11,11,12	0.74	0	14,15,17	2.56	4 (28%)
5	NAG	A	508	1,5	14,14,15	0.61	0	15,19,21	1.53	2 (13%)
5	FUC	A	509	5	10,10,11	0.57	0	14,14,16	1.09	2 (14%)
4	NAG	B	503	1,4	14,14,15	0.42	0	15,19,21	1.41	1 (6%)
4	NAG	B	504	4	14,14,15	0.52	0	15,19,21	0.85	0
4	BMA	B	505	4	11,11,12	0.94	0	14,15,17	1.56	1 (7%)
4	MAN	B	506	4	11,11,12	0.65	0	14,15,17	1.65	3 (21%)
4	MAN	B	507	4	11,11,12	0.60	0	14,15,17	2.34	4 (28%)
5	NAG	B	509	1,5	14,14,15	0.59	0	15,19,21	0.85	0
5	FUC	B	510	5	10,10,11	0.67	0	14,14,16	1.01	1 (7%)
4	NAG	C	504	1,4	14,14,15	0.76	1 (7%)	15,19,21	1.06	1 (6%)
4	NAG	C	505	4	14,14,15	0.57	0	15,19,21	1.19	1 (6%)
4	BMA	C	506	4	11,11,12	0.51	0	14,15,17	1.21	1 (7%)
4	MAN	C	507	4	11,11,12	0.56	0	14,15,17	2.05	3 (21%)
4	MAN	C	508	4	11,11,12	0.63	0	14,15,17	2.24	3 (21%)
5	NAG	C	510	1,5	14,14,15	0.51	0	15,19,21	1.06	1 (6%)
5	FUC	C	511	5	10,10,11	0.84	0	14,14,16	2.21	5 (35%)
4	NAG	D	503	1,4	14,14,15	0.68	0	15,19,21	1.38	2 (13%)
4	NAG	D	504	4	14,14,15	0.50	0	15,19,21	1.08	1 (6%)
4	BMA	D	505	4	11,11,12	0.72	0	14,15,17	1.09	1 (7%)
4	MAN	D	506	4	11,11,12	0.58	0	14,15,17	1.45	1 (7%)
4	MAN	D	507	4	11,11,12	0.62	0	14,15,17	1.96	5 (35%)
7	NAG	D	509	1,7	14,14,15	0.45	0	15,19,21	1.68	2 (13%)
7	NAG	D	510	7	14,14,15	0.55	0	15,19,21	1.17	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	503	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	504	4	-	0/6/23/26	0/1/1/1
4	BMA	A	505	4	-	0/2/19/22	0/1/1/1
4	MAN	A	506	4	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	507	4	-	0/2/19/22	0/1/1/1
5	NAG	A	508	1,5	-	0/6/23/26	0/1/1/1
5	FUC	A	509	5	-	0/0/17/20	0/1/1/1
4	NAG	B	503	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	504	4	-	0/6/23/26	0/1/1/1
4	BMA	B	505	4	-	0/2/19/22	0/1/1/1
4	MAN	B	506	4	-	0/2/19/22	0/1/1/1
4	MAN	B	507	4	-	0/2/19/22	0/1/1/1
5	NAG	B	509	1,5	-	0/6/23/26	0/1/1/1
5	FUC	B	510	5	-	0/0/17/20	0/1/1/1
4	NAG	C	504	1,4	-	0/6/23/26	0/1/1/1
4	NAG	C	505	4	-	0/6/23/26	0/1/1/1
4	BMA	C	506	4	-	0/2/19/22	0/1/1/1
4	MAN	C	507	4	-	0/2/19/22	0/1/1/1
4	MAN	C	508	4	-	0/2/19/22	0/1/1/1
5	NAG	C	510	1,5	-	0/6/23/26	0/1/1/1
5	FUC	C	511	5	-	0/0/17/20	0/1/1/1
4	NAG	D	503	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	504	4	-	0/6/23/26	0/1/1/1
4	BMA	D	505	4	-	0/2/19/22	0/1/1/1
4	MAN	D	506	4	-	0/2/19/22	0/1/1/1
4	MAN	D	507	4	-	0/2/19/22	0/1/1/1
7	NAG	D	509	1,7	-	0/6/23/26	0/1/1/1
7	NAG	D	510	7	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	504	NAG	O5-C1	-2.01	1.40	1.43

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	511	FUC	O5-C1-C2	-3.90	104.53	110.86
4	C	507	MAN	C3-C4-C5	-3.35	104.36	110.20
4	C	504	NAG	C2-N2-C7	-2.99	119.20	123.04
7	D	509	NAG	O3-C3-C2	-2.67	103.83	109.11
4	B	506	MAN	C3-C4-C5	-2.58	105.70	110.20

There are no chirality outliers.

There are no torsion outliers.



There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	508	NAG	1	0
4	B	503	NAG	1	0
4	D	503	NAG	1	0
4	D	505	BMA	1	0

## 5.6 Ligand geometry

Of 16 ligands modelled in this entry, 5 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	EPE	A	502	-	14,15,15	0.67	0	18,20,20	3.07	6 (33%)
6	NAG	A	510	1	14,14,15	0.67	1 (7%)	15,19,21	1.35	2 (13%)
3	EPE	B	502	-	14,15,15	0.53	0	18,20,20	2.70	8 (44%)
6	NAG	B	508	1	14,14,15	0.44	0	15,19,21	1.88	1 (6%)
6	NAG	B	511	1	14,14,15	0.58	0	15,19,21	2.36	1 (6%)
3	EPE	C	503	-	14,15,15	0.59	0	18,20,20	2.16	5 (27%)
6	NAG	C	509	1	14,14,15	0.54	0	15,19,21	1.62	3 (20%)
6	NAG	C	512	1	14,14,15	0.68	1 (7%)	15,19,21	1.78	1 (6%)
3	EPE	D	502	-	14,15,15	0.74	0	18,20,20	3.09	7 (38%)
6	NAG	D	508	1	14,14,15	0.48	0	15,19,21	0.93	1 (6%)
6	NAG	D	511	1	14,14,15	0.45	0	15,19,21	1.66	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EPE	A	502	-	-	0/9/19/19	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	510	1	-	0/6/23/26	0/1/1/1
3	EPE	B	502	-	-	0/9/19/19	0/1/1/1
6	NAG	B	508	1	-	0/6/23/26	0/1/1/1
6	NAG	B	511	1	-	0/6/23/26	0/1/1/1
3	EPE	C	503	-	-	0/9/19/19	0/1/1/1
6	NAG	C	509	1	-	0/6/23/26	0/1/1/1
6	NAG	C	512	1	-	0/6/23/26	0/1/1/1
3	EPE	D	502	-	-	0/9/19/19	0/1/1/1
6	NAG	D	508	1	-	0/6/23/26	0/1/1/1
6	NAG	D	511	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	510	NAG	C1-C2	2.04	1.55	1.52
6	C	512	NAG	C1-C2	2.12	1.55	1.52

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	511	NAG	C4-C3-C2	-4.15	104.78	111.23
6	D	511	NAG	C2-N2-C7	-2.73	119.54	123.04
6	A	510	NAG	C3-C2-N2	-2.20	105.28	110.56
3	A	502	EPE	O3S-S-O2S	-2.06	106.81	111.61
3	B	502	EPE	C9-N1-C6	2.03	116.48	111.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	EPE	2	0
3	B	502	EPE	2	0
3	C	503	EPE	1	0
3	D	502	EPE	2	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/393 (98%)	0.55	15 (3%) 43 57	14, 28, 42, 61	0
1	B	388/393 (98%)	0.63	22 (5%) 27 41	16, 28, 43, 57	0
1	C	388/393 (98%)	0.61	25 (6%) 23 34	11, 28, 42, 59	0
1	D	388/393 (98%)	0.45	8 (2%) 67 79	12, 27, 42, 61	0
All	All	1552/1572 (98%)	0.56	70 (4%) 37 52	11, 28, 42, 61	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	469	ILE	10.7
1	A	469	ILE	9.1
1	D	468	PRO	7.6
1	D	469	ILE	6.8
1	B	469	ILE	5.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	503	14/15	0.90	0.20	7.12	26,32,38,38	0
4	NAG	B	503	14/15	0.89	0.19	2.35	22,34,37,38	0
4	NAG	D	503	14/15	0.84	0.18	1.06	30,36,42,43	0
4	NAG	C	504	14/15	0.88	0.19	0.66	27,32,38,39	0
4	MAN	C	507	11/12	0.74	0.23	-	48,52,54,56	0
4	NAG	C	505	14/15	0.90	0.15	-	29,31,34,37	0
5	FUC	B	510	10/11	0.73	0.39	-	70,71,71,72	0
5	NAG	A	508	14/15	0.72	0.21	-	58,62,65,69	0
5	NAG	B	509	14/15	0.72	0.26	-	58,63,68,71	0
4	NAG	A	504	14/15	0.89	0.18	-	31,32,35,36	0
5	FUC	A	509	10/11	0.73	0.35	-	66,67,67,68	0
7	NAG	D	510	14/15	0.57	0.28	-	63,64,66,66	0
4	MAN	B	506	11/12	0.79	0.22	-	43,46,47,49	0
4	MAN	D	506	11/12	0.84	0.23	-	49,51,53,55	0
4	BMA	D	505	11/12	0.89	0.15	-	42,46,48,51	0
4	NAG	B	504	14/15	0.90	0.14	-	35,37,40,40	0
4	BMA	B	505	11/12	0.92	0.23	-	42,44,47,50	0
4	MAN	C	508	11/12	0.85	0.25	-	50,51,52,53	0
4	BMA	C	506	11/12	0.84	0.19	-	42,46,48,50	0
5	NAG	C	510	14/15	0.81	0.21	-	52,57,62,63	0
4	BMA	A	505	11/12	0.84	0.16	-	40,44,47,50	0
4	MAN	B	507	11/12	0.69	0.20	-	47,50,51,52	0
4	MAN	D	507	11/12	0.76	0.26	-	54,54,55,56	0
4	NAG	D	504	14/15	0.89	0.13	-	35,36,39,41	0
7	NAG	D	509	14/15	0.70	0.22	-	54,58,61,62	0
4	MAN	A	507	11/12	0.76	0.28	-	52,54,54,55	0
5	FUC	C	511	10/11	0.69	0.32	-	63,64,65,65	0
4	MAN	A	506	11/12	0.78	0.22	-	46,48,50,51	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EPE	D	502	15/15	0.83	0.25	5.86	38,41,45,45	0
3	EPE	C	503	15/15	0.86	0.26	5.27	42,48,51,52	0
3	EPE	A	502	15/15	0.91	0.20	2.04	37,43,45,46	0
6	NAG	A	510	14/15	0.79	0.28	1.56	50,54,55,56	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EPE	B	502	15/15	0.92	0.17	1.47	35,38,39,40	0
2	CA	A	501	1/1	0.94	0.08	-1.87	34,34,34,34	0
2	CA	B	501	1/1	0.97	0.07	-1.94	30,30,30,30	0
2	CA	C	502	1/1	0.86	0.09	-2.31	34,34,34,34	0
2	CA	D	501	1/1	1.00	0.03	-2.59	32,32,32,32	0
2	CA	C	501	1/1	0.87	0.15	-	64,64,64,64	0
6	NAG	C	512	14/15	0.71	0.31	-	54,56,59,59	0
6	NAG	B	508	14/15	0.71	0.29	-	50,53,55,55	0
6	NAG	B	511	14/15	0.76	0.33	-	50,53,54,54	0
6	NAG	C	509	14/15	0.69	0.32	-	60,64,64,64	0
6	NAG	D	508	14/15	0.69	0.31	-	59,63,65,65	0
6	NAG	D	511	14/15	0.76	0.25	-	53,56,57,58	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.