



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 02:25 AM GMT

PDB ID : 2H0Q
Title : Crystal Structure of the PGM domain of the Suppressor of T-Cell receptor (Sts-1)
Authors : Nassar, N.; Ford, B.; Carpino, N.
Deposited on : 2006-05-15
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5422 (1.84-1.80)
Clashscore	102246	6347 (1.84-1.80)
Ramachandran outliers	100387	6276 (1.84-1.80)
Sidechain outliers	100360	6276 (1.84-1.80)
RSRZ outliers	91569	5439 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 6472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Suppressor of T-cell receptor signaling 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	261	Total	C 2066	N 1311	O 361	S 375	19	0	6	0
1	B	256	Total	C 2032	N 1287	O 356	S 372	17	0	9	0
1	C	261	Total	C 2071	N 1314	O 363	S 377	17	0	2	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	132	Total O 132 132	0	0
2	B	109	Total O 109 109	0	0
2	C	62	Total O 62 62	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Suppressor of T-cell receptor signaling 1



- Molecule 1: Suppressor of T-cell receptor signaling 1



- Molecule 1: Suppressor of T-cell receptor signaling 1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.23Å 74.34Å 100.11Å 90.00° 101.46° 90.00°	Depositor
Resolution (Å)	98.06 – 1.82 45.49 – 1.82	Depositor EDS
% Data completeness (in resolution range)	96.2 (98.06-1.82) 96.2 (45.49-1.82)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) >$ ¹	2.80 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R , R_{free}	0.196 , 0.236 0.232 , 0.268	Depositor DCC
R_{free} test set	3640 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.713	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 57.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.34$	Xtriage
Outliers	0 of 72282 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6472	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	0/2154	0.70	1/2924 (0.0%)
1	B	0.64	1/2128 (0.0%)	0.68	1/2889 (0.0%)
1	C	0.49	0/2140	0.63	2/2905 (0.1%)
All	All	0.60	1/6422 (0.0%)	0.67	4/8718 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	602	CYS	CB-SG	-6.48	1.71	1.82

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	597	LEU	CA-CB-CG	-7.45	98.17	115.30
1	B	602	CYS	CB-CA-C	-5.39	99.63	110.40
1	C	405	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	A	474	LEU	CA-CB-CG	5.04	126.90	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2066	0	2038	25	0
1	B	2032	0	2023	40	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2071	0	2051	24	1
2	A	132	0	0	0	0
2	B	109	0	0	3	0
2	C	62	0	0	3	0
All	All	6472	0	6112	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:436[B]:MET:CE	1:B:436[B]:MET:HE2	1.55	1.35
1:B:570:GLU:HG2	2:B:738:HOH:O	1.36	1.22
1:A:436[B]:MET:CE	1:B:436[B]:MET:CE	2.16	1.15
1:C:462[B]:ARG:HH12	1:C:565:HIS:CE1	1.74	1.04
1:A:436[B]:MET:SD	1:B:436[B]:MET:HE3	1.67	1.02
1:A:436[B]:MET:SD	1:B:436[B]:MET:CE	0.92	0.97
1:A:622:THR:HA	1:B:626[B]:THR:HG21	1.47	0.96
1:A:436[B]:MET:SD	1:B:436[B]:MET:HE1	0.58	0.94
1:C:405:ARG:NH2	2:C:692:HOH:O	2.07	0.86
1:B:626[B]:THR:HG23	2:B:663:HOH:O	1.78	0.82
1:B:607:GLU:HG3	1:B:608:THR:N	1.98	0.76
1:A:436[B]:MET:HE3	1:B:436[B]:MET:HE2	1.65	0.75
1:A:468:HIS:NE2	1:A:472:LYS:HE3	2.05	0.72
1:C:462[B]:ARG:NH1	1:C:565:HIS:CE1	2.56	0.72
1:A:582:GLN:NE2	1:A:590[A]:MET:SD	2.64	0.70
1:A:436[B]:MET:SD	1:B:436[B]:MET:HE2	1.13	0.68
1:A:468:HIS:CD2	1:A:472:LYS:HE3	2.30	0.67
1:B:374[B]:CYS:SG	1:B:376:PHE:CE1	2.83	0.67
1:C:439:ARG:NH1	1:C:443:GLU:OE2	2.33	0.62
1:B:426:LYS:NZ	1:B:499:THR:HG23	2.16	0.59
1:C:384:MET:HG3	1:C:429:PRO:HG2	1.85	0.58
1:A:523:HIS:HE1	1:A:552:GLU:OE2	1.87	0.57
1:C:583:ASN:HD21	1:C:585:LYS:HD2	1.71	0.56
1:A:384:MET:HG3	1:A:429:PRO:HG2	1.89	0.55
1:C:462[B]:ARG:HH12	1:C:565:HIS:HE1	1.49	0.54
1:C:626:THR:HA	2:C:685:HOH:O	2.08	0.54
1:C:380:HIS:CE1	1:C:462[B]:ARG:HD2	2.45	0.51
1:B:459:PRO:HD3	1:B:486:PRO:HA	1.93	0.51
1:A:554:LYS:NZ	1:A:604:GLU:OE1	2.42	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:603:GLU:HB2	1:B:614:THR:HG21	1.93	0.49
1:B:587:PHE:O	1:B:591:VAL:HG23	2.11	0.49
1:B:607:GLU:HG3	1:B:608:THR:H	1.74	0.48
1:C:486:PRO:HB2	1:C:524:ILE:O	2.14	0.48
1:A:491:TRP:CE2	1:A:493:LYS:HG3	2.49	0.48
1:A:621[A]:LEU:HD11	1:B:433:PHE:HE1	1.79	0.47
1:B:375:LEU:HD11	1:B:562:ILE:HD12	1.96	0.47
1:B:446:LEU:HD12	2:B:679:HOH:O	2.14	0.46
1:C:405:ARG:HD3	1:C:412:HIS:HA	1.97	0.46
1:C:555:SER:HB3	1:C:556:LYS:HG3	1.97	0.46
1:B:373:ARG:N	1:B:558:ASN:O	2.49	0.45
1:A:626:THR:HG23	1:B:623:HIS:CD2	2.52	0.45
1:B:607:GLU:O	1:B:609:GLY:N	2.44	0.45
1:C:462[A]:ARG:NH2	2:C:660:HOH:O	2.48	0.44
1:B:426:LYS:HZ2	1:B:499:THR:HG23	1.80	0.44
1:B:509:LEU:HB2	1:B:516[A]:VAL:HG21	1.97	0.44
1:C:426:LYS:NZ	1:C:496:ALA:HB3	2.32	0.44
1:B:384:MET:HG3	1:B:429:PRO:HG2	1.99	0.44
1:A:517:ASP:OD1	1:A:519:THR:HB	2.17	0.44
1:C:590:MET:CE	1:C:594:ILE:HD11	2.47	0.44
1:C:580:SER:HA	1:C:581:PRO:HD3	1.91	0.43
1:A:623:HIS:CD2	1:B:626[A]:THR:HG23	2.54	0.43
1:C:383:ARG:HG2	1:C:462[B]:ARG:HD3	2.01	0.43
1:A:486:PRO:HB2	1:A:524:ILE:O	2.19	0.43
1:B:490:GLU:HB3	1:B:565:HIS:CG	2.54	0.43
1:B:582:GLN:NE2	1:B:590[B]:MET:SD	2.91	0.42
1:A:374:CYS:HB2	1:A:559:ASN:OD1	2.18	0.42
1:C:433:PHE:O	1:C:437:GLN:HG3	2.19	0.42
1:C:462[B]:ARG:HH12	1:C:565:HIS:HD1	1.60	0.42
1:B:605:LEU:HD21	1:B:612:GLN:NE2	2.34	0.42
1:C:541:ARG:O	1:C:545:VAL:HG23	2.19	0.42
1:B:426:LYS:NZ	1:B:499:THR:CG2	2.83	0.42
1:A:488:LEU:HD13	1:A:562:ILE:HG23	2.02	0.41
1:B:374[B]:CYS:SG	1:B:376:PHE:CZ	3.10	0.41
1:C:536:ASP:O	1:C:540:ASN:ND2	2.54	0.41
1:B:551:SER:HA	1:B:554:LYS:HG3	2.02	0.41
1:A:468:HIS:HE2	1:A:472:LYS:HE3	1.82	0.41
1:A:550:ILE:O	1:A:554:LYS:HG2	2.20	0.41
1:C:589:GLN:HG3	1:C:592:ARG:NH1	2.35	0.41
1:C:462[B]:ARG:NH1	1:C:565:HIS:ND1	2.60	0.41
1:B:602:CYS:HB3	1:B:611:TRP:HB3	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:574:CYS:SG	1:B:581:PRO:HA	2.61	0.41
1:B:513:ASN:HB2	1:C:513:ASN:CG	2.41	0.40
1:B:607:GLU:C	1:B:609:GLY:H	2.25	0.40
1:B:584:SER:O	1:B:588:VAL:HG23	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:391:TYR:OH	1:C:540:ASN:ND2[3_555]	2.16	0.04

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	265/261 (102%)	259 (98%)	6 (2%)	0	100 100
1	B	263/261 (101%)	252 (96%)	10 (4%)	1 (0%)	39 23
1	C	262/261 (100%)	252 (96%)	10 (4%)	0	100 100
All	All	790/783 (101%)	763 (97%)	26 (3%)	1 (0%)	56 38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	608	THR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	234/229 (102%)	228 (97%)	6 (3%)	54 37
1	B	233/229 (102%)	226 (97%)	7 (3%)	48 31
1	C	232/229 (101%)	225 (97%)	7 (3%)	48 31
All	All	699/687 (102%)	679 (97%)	20 (3%)	54 33

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	373	ARG
1	A	390	LYS
1	A	396[A]	CYS
1	A	396[B]	CYS
1	A	516	VAL
1	A	605	LEU
1	B	394	SER
1	B	405[A]	ARG
1	B	405[B]	ARG
1	B	531	ILE
1	B	532	SER
1	B	582	GLN
1	B	589	GLN
1	C	377	VAL
1	C	405	ARG
1	C	531	ILE
1	C	558	ASN
1	C	583	ASN
1	C	597	LEU
1	C	630	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	523	HIS
1	B	412	HIS
1	B	577	GLN
1	B	582	GLN
1	B	612	GLN
1	C	412	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	558	ASN
1	C	630	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/261 (100%)	0.52	15 (5%) 27 22	20, 30, 43, 50	0
1	B	256/261 (98%)	0.72	27 (10%) 8 6	23, 31, 45, 51	0
1	C	261/261 (100%)	1.23	66 (25%) 1 0	26, 38, 49, 55	0
All	All	778/783 (99%)	0.83	108 (13%) 4 3	20, 33, 46, 55	0

All (108) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	496	ALA	9.5
1	C	605	LEU	7.9
1	C	610	ILE	7.2
1	C	531	ILE	6.8
1	C	606	GLY	6.1
1	C	555	SER	5.9
1	C	496	ALA	5.7
1	C	578	GLY	5.6
1	C	537	THR	5.6
1	C	579	LEU	5.5
1	C	631	TRP	5.2
1	C	543	PHE	5.1
1	C	536	ASP	5.0
1	B	628	GLY	4.9
1	A	605	LEU	4.9
1	C	607	GLU	4.7
1	B	605	LEU	4.4
1	C	530	ALA	4.3
1	B	555	SER	4.3
1	C	498	SER	4.3
1	B	531	ILE	4.2
1	C	556	LYS	4.2
1	A	498	SER	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	611	TRP	4.1
1	C	576	LEU	4.1
1	C	632	ARG	4.1
1	A	478	ASN	3.9
1	B	498	SER	3.9
1	C	491	TRP	3.8
1	C	608	THR	3.8
1	C	534	SER	3.8
1	A	479	HIS	3.7
1	C	521	ARG	3.6
1	B	606	GLY	3.6
1	C	585	LYS	3.5
1	C	558	ASN	3.5
1	C	557	GLY	3.5
1	C	602	CYS	3.4
1	C	478	ASN	3.3
1	C	373	ARG	3.3
1	B	499	THR	3.2
1	C	580	SER	3.2
1	C	538	TYR	3.2
1	B	607	GLU	3.1
1	B	391	TYR	3.1
1	B	579	LEU	3.1
1	C	495	VAL	3.1
1	C	609	GLY	3.0
1	C	554	LYS	3.0
1	C	590	MET	3.0
1	C	527	SER	3.0
1	C	540	ASN	3.0
1	C	584	SER	3.0
1	B	435	CYS	2.9
1	C	391	TYR	2.9
1	B	578	GLY	2.9
1	C	535	TYR	2.8
1	B	581	PRO	2.8
1	C	539	ILE	2.8
1	C	581	PRO	2.8
1	C	574	CYS	2.8
1	B	556	LYS	2.7
1	B	433	PHE	2.7
1	C	553	CYS	2.6
1	A	496	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	433	PHE	2.6
1	C	586	ASP	2.6
1	C	603	GLU	2.6
1	C	547	LYS	2.6
1	B	585	LYS	2.6
1	C	435	CYS	2.5
1	A	381	GLY	2.5
1	A	435	CYS	2.5
1	A	387	VAL	2.5
1	B	554	LYS	2.5
1	C	612	GLN	2.5
1	C	592	ARG	2.5
1	C	582	GLN	2.5
1	C	519	THR	2.4
1	C	614	THR	2.4
1	C	529	LEU	2.4
1	C	493	LYS	2.4
1	C	604	GLU	2.4
1	B	495	VAL	2.4
1	A	590[A]	MET	2.4
1	B	592	ARG	2.3
1	B	612	GLN	2.3
1	A	414	LEU	2.3
1	C	589	GLN	2.3
1	C	532	SER	2.3
1	B	422	ARG	2.2
1	C	544	GLN	2.2
1	A	411	PRO	2.2
1	B	621	LEU	2.1
1	C	374	CYS	2.1
1	B	558	ASN	2.1
1	B	519[A]	THR	2.1
1	C	629	PHE	2.1
1	A	430	ILE	2.1
1	A	521	ARG	2.1
1	C	633	GLU	2.1
1	C	436[A]	MET	2.1
1	C	593	LYS	2.0
1	C	583	ASN	2.0
1	B	438	ALA	2.0
1	B	441	VAL	2.0
1	C	390	LYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	436[A]	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.