



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:07 PM GMT

PDB ID : 4H6C  
Title : Crystal Structure of the Allene Oxide Cyclase 1 from *Physcomitrella patens*  
Authors : Neumann, P.; Ficner, R.  
Deposited on : 2012-09-19  
Resolution : 1.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2199 (1.40-1.32)
Clashscore	102246	2337 (1.40-1.32)
Ramachandran outliers	100387	2280 (1.40-1.32)
Sidechain outliers	100360	2279 (1.40-1.32)
RSRZ outliers	91569	2199 (1.40-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	<div> <div>4%</div> <div>84% 9% 8%</div> </div>
1	B	195	<div> <div>5%</div> <div>80% 11% 9%</div> </div>
1	C	195	<div> <div>7%</div> <div>76% 14% 10%</div> </div>
1	D	195	<div> <div>2%</div> <div>85% 7% 8%</div> </div>
1	E	195	<div> <div>2%</div> <div>83% 9% 8%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	195	
1	G	195	
1	H	195	
1	I	195	
1	J	195	
1	K	195	
1	L	195	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	203	-	-	-	X
2	PO4	G	203	-	-	-	X
2	PO4	H	202	-	-	-	X
3	HEZ	A	202	-	-	X	X
3	HEZ	B	201	-	-	-	X
3	HEZ	D	301	-	-	-	X
3	HEZ	E	201	-	-	-	X
3	HEZ	F	201	-	-	-	X
3	HEZ	G	202	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 19586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Allene oxide cyclase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	4	0
			1437	932	233	267	5			
1	D	180	Total	C	N	O	S	0	3	0
			1422	924	231	262	5			
1	C	175	Total	C	N	O	S	0	2	0
			1381	899	224	253	5			
1	B	177	Total	C	N	O	S	0	3	0
			1408	916	229	258	5			
1	E	180	Total	C	N	O	S	0	4	0
			1431	930	233	263	5			
1	F	175	Total	C	N	O	S	0	2	0
			1379	897	224	253	5			
1	G	175	Total	C	N	O	S	0	5	0
			1412	921	229	257	5			
1	H	175	Total	C	N	O	S	0	1	0
			1375	895	224	251	5			
1	I	180	Total	C	N	O	S	0	1	0
			1405	912	228	260	5			
1	J	175	Total	C	N	O	S	0	2	0
			1383	899	225	254	5			
1	K	180	Total	C	N	O	S	0	5	0
			1442	937	234	266	5			
1	L	175	Total	C	N	O	S	0	2	0
			1384	901	225	253	5			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
A	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
A	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
A	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
A	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP Q8GS38
D	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
D	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
D	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
D	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
D	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
D	0	SER	-	EXPRESSION TAG	UNP Q8GS38
C	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
C	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
C	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
C	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
C	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
C	0	SER	-	EXPRESSION TAG	UNP Q8GS38
B	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
B	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
B	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
B	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
B	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
B	0	SER	-	EXPRESSION TAG	UNP Q8GS38
E	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
E	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
E	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
E	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
E	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
E	0	SER	-	EXPRESSION TAG	UNP Q8GS38
F	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
F	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
F	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
F	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
F	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
F	0	SER	-	EXPRESSION TAG	UNP Q8GS38
G	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
G	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
G	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
G	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
G	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
G	0	SER	-	EXPRESSION TAG	UNP Q8GS38
H	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
H	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
H	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
H	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
H	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
H	0	SER	-	EXPRESSION TAG	UNP Q8GS38
I	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
I	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
I	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
I	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
I	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
I	0	SER	-	EXPRESSION TAG	UNP Q8GS38
J	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
J	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
J	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
J	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
J	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
J	0	SER	-	EXPRESSION TAG	UNP Q8GS38
K	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
K	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
K	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
K	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
K	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
K	0	SER	-	EXPRESSION TAG	UNP Q8GS38
L	-5	GLY	-	EXPRESSION TAG	UNP Q8GS38
L	-4	GLY	-	EXPRESSION TAG	UNP Q8GS38
L	-3	PRO	-	EXPRESSION TAG	UNP Q8GS38
L	-2	LEU	-	EXPRESSION TAG	UNP Q8GS38
L	-1	GLY	-	EXPRESSION TAG	UNP Q8GS38
L	0	SER	-	EXPRESSION TAG	UNP Q8GS38

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		
2	G	1	Total	O	P	0	0
			5	4	1		
2	G	1	Total	O	P	0	0
			5	4	1		
2	H	1	Total	O	P	0	0
			5	4	1		
2	H	1	Total	O	P	0	0
			5	4	1		
2	I	1	Total	O	P	0	0
			5	4	1		
2	I	1	Total	O	P	0	0
			5	4	1		
2	J	1	Total	O	P	0	0
			5	4	1		
2	J	1	Total	O	P	0	0
			5	4	1		
2	K	1	Total	O	P	0	0
			5	4	1		
2	L	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	D	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		
3	E	1	Total	C	O	0	0
			8	6	2		
3	F	1	Total	C	O	0	0
			8	6	2		
3	G	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	238	Total	O	0	0
			238	238		
4	D	302	Total	O	0	0
			302	302		
4	C	197	Total	O	0	0
			197	197		
4	B	245	Total	O	0	0
			245	245		
4	E	280	Total	O	0	0
			280	280		
4	F	209	Total	O	0	0
			209	209		

*Continued on next page...*





*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	192	Total 192	O 192	0	0
4	H	196	Total 196	O 196	0	0
4	I	211	Total 211	O 211	0	0
4	J	162	Total 163	O 163	0	1
4	K	238	Total 239	O 239	0	1
4	L	121	Total 122	O 122	0	1



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Allene oxide cyclase





- Molecule 1: Allene oxide cyclase




- Molecule 1: Allene oxide cyclase



- Molecule 1: Allene oxide cyclase





- Molecule 1: Allene oxide cyclase









- Molecule 1: Allene oxide cyclase





- Molecule 1: Allene oxide cyclase





- Molecule 1: Allene oxide cyclase



- Molecule 1: Allene oxide cyclase





- Molecule 1: Allene oxide cyclase








● Molecule 1: Allene oxide cyclase



● Molecule 1: Allene oxide cyclase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.32Å 67.43Å 161.78Å 84.61° 79.32° 61.99°	Depositor
Resolution (Å)	19.80 – 1.35 19.80 – 1.35	Depositor EDS
% Data completeness (in resolution range)	82.4 (19.80-1.35) 82.3 (19.80-1.35)	Depositor EDS
$R_{merge}$	0.02	Depositor
$R_{sym}$	0.00	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.138 , 0.174 0.139 , 0.175	Depositor DCC
$R_{free}$ test set	22342 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.4	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 62.5	EDS
Estimated twinning fraction	0.042 for h,h-k,h-l 0.024 for -h,-h+k,-l 0.022 for -h,-k,-h+l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 446818 reflections	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	19586	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, HEZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.82	0/1481	0.82	3/2011 (0.1%)
1	B	0.74	0/1449	0.73	1/1965 (0.1%)
1	C	0.75	0/1421	0.71	0/1927
1	D	0.80	0/1465	0.81	0/1991
1	E	0.75	0/1474	0.77	1/2002 (0.0%)
1	F	0.79	0/1419	0.72	0/1924
1	G	0.73	0/1454	0.71	0/1972
1	H	0.72	0/1416	0.71	0/1920
1	I	0.73	0/1447	0.69	0/1965
1	J	0.69	0/1424	0.68	0/1931
1	K	0.72	0/1485	0.70	0/2017
1	L	0.67	0/1424	0.66	0/1929
All	All	0.74	0/17359	0.73	5/23554 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	86	ARG	NE-CZ-NH1	8.16	124.38	120.30
1	A	173	ASP	CB-CG-OD2	-5.63	113.24	118.30
1	A	24	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	A	173	ASP	CB-CG-OD1	5.09	122.88	118.30
1	B	173	ASP	CB-CG-OD1	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1437	0	1388	17	0
1	B	1408	0	1373	21	0
1	C	1381	0	1350	23	0
1	D	1422	0	1380	14	0
1	E	1431	0	1392	17	0
1	F	1379	0	1343	16	0
1	G	1412	0	1376	17	0
1	H	1375	0	1341	13	0
1	I	1405	0	1366	6	0
1	J	1383	0	1344	13	0
1	K	1442	0	1404	8	0
1	L	1384	0	1353	18	0
2	A	5	0	0	0	0
2	B	10	0	0	1	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
2	G	10	0	0	0	0
2	H	10	0	0	0	0
2	I	10	0	0	0	0
2	J	10	0	0	0	0
2	K	5	0	0	0	0
2	L	5	0	0	0	0
3	A	8	0	14	10	0
3	B	8	0	14	4	0
3	D	8	0	13	1	0
3	E	8	0	14	0	0
3	F	8	0	14	3	0
3	G	8	0	14	1	0
4	A	238	0	0	4	0
4	B	245	0	0	9	2
4	C	197	0	0	2	0
4	D	302	0	0	4	0
4	E	280	0	0	6	2
4	F	209	0	0	2	0
4	G	192	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	196	0	0	2	2
4	I	211	0	0	4	3
4	J	163	0	0	2	0
4	K	239	0	0	1	1
4	L	122	0	0	4	0
All	All	19586	0	16493	183	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 183 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:90:LEU:HD13	1:G:104[B]:GLN:HG2	1.34	1.08
1:A:107[B]:TYR:CE2	3:A:202:HEZ:C5	2.48	0.97
1:A:107[B]:TYR:CE2	3:A:202:HEZ:H51	2.06	0.91
1:G:90:LEU:CD1	1:G:104[B]:GLN:HG2	2.02	0.87
3:B:201:HEZ:H42	4:B:475:HOH:O	1.78	0.83

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:537:HOH:O	4:E:579:HOH:O[1_565]	1.81	0.39
4:H:493:HOH:O	4:I:508:HOH:O[1_565]	2.00	0.20
4:B:432:HOH:O	4:E:468:HOH:O[1_565]	2.03	0.17
4:H:493:HOH:O	4:I:509:HOH:O[1_565]	2.05	0.15
4:I:500:HOH:O	4:K:524:HOH:O[1_656]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/195 (93%)	180 (99%)	2 (1%)	0	100	100
1	B	176/195 (90%)	176 (100%)	0	0	100	100
1	C	173/195 (89%)	172 (99%)	1 (1%)	0	100	100
1	D	181/195 (93%)	179 (99%)	2 (1%)	0	100	100
1	E	182/195 (93%)	181 (100%)	1 (0%)	0	100	100
1	F	173/195 (89%)	172 (99%)	1 (1%)	0	100	100
1	G	176/195 (90%)	175 (99%)	1 (1%)	0	100	100
1	H	172/195 (88%)	170 (99%)	1 (1%)	1 (1%)	30	8
1	I	179/195 (92%)	178 (99%)	1 (1%)	0	100	100
1	J	173/195 (89%)	172 (99%)	1 (1%)	0	100	100
1	K	183/195 (94%)	181 (99%)	2 (1%)	0	100	100
1	L	173/195 (89%)	170 (98%)	3 (2%)	0	100	100
All	All	2123/2340 (91%)	2106 (99%)	16 (1%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	180	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/157 (98%)	153 (99%)	1 (1%)	90	73
1	B	151/157 (96%)	151 (100%)	0	100	100
1	C	148/157 (94%)	148 (100%)	0	100	100
1	D	152/157 (97%)	151 (99%)	1 (1%)	88	68
1	E	153/157 (98%)	153 (100%)	0	100	100
1	F	147/157 (94%)	145 (99%)	2 (1%)	74	43
1	G	151/157 (96%)	150 (99%)	1 (1%)	88	68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	147/157 (94%)	146 (99%)	1 (1%)	88	68
1	I	150/157 (96%)	148 (99%)	2 (1%)	76	45
1	J	148/157 (94%)	147 (99%)	1 (1%)	88	68
1	K	155/157 (99%)	155 (100%)	0	100	100
1	L	148/157 (94%)	146 (99%)	2 (1%)	74	43
All	All	1804/1884 (96%)	1793 (99%)	11 (1%)	90	73

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	137	ILE
1	H	179	LEU
1	J	153	LYS
1	F	135	GLN
1	I	179	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	177	GLN
1	I	43	HIS
1	H	135	GLN
1	F	135	GLN
1	H	182	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	201	-	4,4,4	0.92	0	6,6,6	0.31	0
3	HEZ	A	202	-	7,7,7	0.42	0	6,6,6	1.11	0
3	HEZ	B	201	-	7,7,7	0.44	0	6,6,6	0.87	0
2	PO4	B	202	-	4,4,4	1.18	1 (25%)	6,6,6	0.35	0
2	PO4	B	203	-	4,4,4	0.44	0	6,6,6	0.29	0
2	PO4	C	201	-	4,4,4	0.25	0	6,6,6	0.33	0
3	HEZ	D	301	-	7,7,7	1.32	2 (28%)	6,6,6	0.85	0
2	PO4	D	302	-	4,4,4	0.56	0	6,6,6	0.29	0
3	HEZ	E	201	-	7,7,7	0.63	0	6,6,6	0.53	0
2	PO4	E	202	-	4,4,4	0.55	0	6,6,6	0.39	0
3	HEZ	F	201	-	7,7,7	0.33	0	6,6,6	0.49	0
2	PO4	F	202	-	4,4,4	0.51	0	6,6,6	0.28	0
2	PO4	G	201	-	4,4,4	0.32	0	6,6,6	0.33	0
3	HEZ	G	202	-	7,7,7	0.55	0	6,6,6	0.52	0
2	PO4	G	203	-	4,4,4	0.28	0	6,6,6	0.31	0
2	PO4	H	201	-	4,4,4	0.24	0	6,6,6	0.34	0
2	PO4	H	202	-	4,4,4	0.64	0	6,6,6	0.29	0
2	PO4	I	201	-	4,4,4	0.63	0	6,6,6	0.28	0
2	PO4	I	202	-	4,4,4	0.27	0	6,6,6	0.27	0
2	PO4	J	201	-	4,4,4	0.70	0	6,6,6	0.40	0
2	PO4	J	202	-	4,4,4	0.41	0	6,6,6	0.28	0
2	PO4	K	201	-	4,4,4	0.71	0	6,6,6	0.29	0
2	PO4	L	201	-	4,4,4	0.46	0	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	201	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEZ	A	202	-	-	0/5/5/5	0/0/0/0
3	HEZ	B	201	-	-	0/5/5/5	0/0/0/0
2	PO4	B	202	-	-	0/0/0/0	0/0/0/0
2	PO4	B	203	-	-	0/0/0/0	0/0/0/0
2	PO4	C	201	-	-	0/0/0/0	0/0/0/0
3	HEZ	D	301	-	-	0/5/5/5	0/0/0/0
2	PO4	D	302	-	-	0/0/0/0	0/0/0/0
3	HEZ	E	201	-	-	0/5/5/5	0/0/0/0
2	PO4	E	202	-	-	0/0/0/0	0/0/0/0
3	HEZ	F	201	-	-	0/5/5/5	0/0/0/0
2	PO4	F	202	-	-	0/0/0/0	0/0/0/0
2	PO4	G	201	-	-	0/0/0/0	0/0/0/0
3	HEZ	G	202	-	-	0/5/5/5	0/0/0/0
2	PO4	G	203	-	-	0/0/0/0	0/0/0/0
2	PO4	H	201	-	-	0/0/0/0	0/0/0/0
2	PO4	H	202	-	-	0/0/0/0	0/0/0/0
2	PO4	I	201	-	-	0/0/0/0	0/0/0/0
2	PO4	I	202	-	-	0/0/0/0	0/0/0/0
2	PO4	J	201	-	-	0/0/0/0	0/0/0/0
2	PO4	J	202	-	-	0/0/0/0	0/0/0/0
2	PO4	K	201	-	-	0/0/0/0	0/0/0/0
2	PO4	L	201	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	202	PO4	P-O1	2.10	1.61	1.52
3	D	301	HEZ	C5-C6	2.13	1.63	1.50
3	D	301	HEZ	O6-C6	2.31	1.54	1.42

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	202	HEZ	10	0
3	B	201	HEZ	4	0
2	B	203	PO4	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	301	HEZ	1	0
3	F	201	HEZ	3	0
3	G	202	HEZ	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	180/195 (92%)	0.17	8 (4%) 38 38	12, 23, 39, 51	2 (1%)
1	B	177/195 (90%)	0.08	10 (5%) 28 27	12, 19, 39, 59	3 (1%)
1	C	175/195 (89%)	0.33	14 (8%) 15 15	12, 24, 39, 55	2 (1%)
1	D	180/195 (92%)	-0.12	4 (2%) 65 67	10, 16, 26, 45	2 (1%)
1	E	180/195 (92%)	-0.07	4 (2%) 65 67	12, 18, 29, 45	1 (0%)
1	F	175/195 (89%)	0.16	8 (4%) 36 36	13, 23, 41, 49	2 (1%)
1	G	175/195 (89%)	0.15	12 (6%) 20 20	16, 23, 40, 51	3 (1%)
1	H	175/195 (89%)	0.21	16 (9%) 11 12	16, 25, 43, 52	4 (2%)
1	I	180/195 (92%)	-0.03	5 (2%) 56 58	17, 25, 38, 48	3 (1%)
1	J	175/195 (89%)	0.31	12 (6%) 20 20	17, 27, 48, 58	3 (1%)
1	K	180/195 (92%)	-0.07	4 (2%) 65 67	16, 22, 34, 47	2 (1%)
1	L	175/195 (89%)	0.56	29 (16%) 2 2	19, 32, 49, 57	1 (0%)
All	All	2127/2340 (90%)	0.14	126 (5%) 26 25	10, 23, 42, 59	28 (1%)

The worst 5 of 126 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	42	ALA	7.5
1	C	42	ALA	6.8
1	F	42	ALA	6.0
1	J	36	LYS	5.4
1	J	42	ALA	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	HEZ	D	301	8/8	0.90	0.23	14.80	21,31,33,35	6
3	HEZ	E	201	8/8	0.92	0.23	11.63	20,34,37,38	6
3	HEZ	G	202	8/8	0.86	0.14	7.06	34,36,38,38	2
2	PO4	B	203	5/5	0.82	0.18	6.12	21,31,38,43	5
3	HEZ	B	201	8/8	0.87	0.22	5.41	40,42,47,50	2
2	PO4	G	203	5/5	0.80	0.18	5.06	26,34,43,43	5
3	HEZ	A	202	8/8	0.92	0.17	4.85	31,33,39,42	5
2	PO4	H	202	5/5	0.85	0.20	3.76	29,41,43,44	5
3	HEZ	F	201	8/8	0.91	0.17	2.41	39,41,42,43	5
2	PO4	I	202	5/5	0.92	0.12	1.45	29,32,39,41	5
2	PO4	C	201	5/5	0.97	0.08	1.06	26,26,28,29	5
2	PO4	A	201	5/5	0.98	0.07	0.57	23,24,26,27	5
2	PO4	J	201	5/5	0.98	0.07	0.37	22,24,25,26	5
2	PO4	J	202	5/5	0.92	0.10	0.25	35,43,47,50	5
2	PO4	B	202	5/5	0.97	0.07	0.06	19,20,23,23	5
2	PO4	I	201	5/5	0.99	0.05	-0.72	23,24,27,27	5
2	PO4	H	201	5/5	0.99	0.05	-0.80	28,28,29,29	5
2	PO4	D	302	5/5	0.99	0.06	-0.94	17,17,19,20	5
2	PO4	K	201	5/5	0.99	0.05	-0.98	26,26,27,27	5
2	PO4	F	202	5/5	0.99	0.05	-1.00	21,23,25,27	5
2	PO4	L	201	5/5	0.99	0.05	-1.10	28,29,29,30	5
2	PO4	E	202	5/5	0.99	0.05	-1.12	19,20,22,23	5
2	PO4	G	201	5/5	0.99	0.04	-1.99	22,23,25,26	5

### 6.5 Other polymers ⓘ

There are no such residues in this entry.