



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:47 PM GMT

PDB ID : 1H80  
Title : 1,3-ALPHA-1,4-BETA-D-GALACTOSE-4-SULFATE-3,6-ANHYDRO-D-GALACTOSE-2-SULFATE 4 GALACTOXYDROLASE  
Authors : Michel, G.; Chantalat, L.; Dideberg, O.  
Deposited on : 2001-01-22  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	
1	B	464	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	1499	-	-	-	X
5	GOL	A	1500	-	-	-	X
5	GOL	A	1501	-	-	-	X
5	GOL	B	1499	-	-	-	X
5	GOL	B	1500	-	-	-	X
5	GOL	B	1501	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IOTA-CARRAGEENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	433	Total	C	N	O	S	55	11	0
			3431	2143	612	657	19			
1	B	430	Total	C	N	O	S	85	9	0
			3401	2125	605	652	19			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Ca	0	0
			3	3		
2	A	3	Total	Ca	0	0
			3	3		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Na	0	0
			3	3		
4	A	3	Total	Na	0	0
			3	3		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	531	Total	O	0	0
			531	531		
6	B	579	Total	O	0	0
			579	579		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: IOTA-CARRAGEENASE



#### • Molecule 1: IOTA-CARRAGEENASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.87Å 90.07Å 124.12Å 90.00° 93.53° 90.00°	Depositor
Resolution (Å)	20.00 – 1.60 8.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	97.4 (20.00-1.60) 97.6 (8.00-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.35 (at 1.60Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.207 , 0.223 0.210 , 0.226	Depositor DCC
$R_{free}$ test set	7885 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.9	Xtriage
Anisotropy	0.204	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.46 , 55.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 156301 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7992	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, CL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/3554	0.65	1/4796 (0.0%)
1	B	0.33	0/3515	0.65	1/4743 (0.0%)
All	All	0.33	0/7069	0.65	2/9539 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	174	ASN	N-CA-C	5.71	126.41	111.00
1	B	174	ASN	N-CA-C	5.50	125.86	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3431	0	3336	60	0
1	B	3401	0	3298	50	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	1	0	0	1	0
3	B	1	0	0	1	0
4	A	3	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	3	0	0	0	0
5	A	18	0	24	3	0
5	B	18	0	24	3	0
6	A	531	0	0	4	0
6	B	579	0	0	1	0
All	All	7992	0	6682	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:VAL:HG12	1:A:370:ILE:HD12	1.24	1.16
1:A:408[B]:CYS:SG	1:A:475:VAL:HG23	2.03	0.98
1:A:309:VAL:CG1	1:A:370:ILE:HD12	1.97	0.94
1:B:269[B]:CYS:SG	1:B:295:SER:HB2	2.14	0.87
1:A:269[B]:CYS:SG	1:A:295:SER:HB2	2.15	0.86
1:B:408[B]:CYS:SG	1:B:475:VAL:HG23	2.14	0.86
1:A:309:VAL:HG12	1:A:370:ILE:CD1	2.06	0.84
1:A:269[B]:CYS:SG	1:A:295:SER:CB	2.72	0.78
1:A:202:ARG:HD2	5:A:1501:GOL:O3	1.83	0.78
1:B:269[B]:CYS:SG	1:B:295:SER:CB	2.72	0.77
1:A:412:CYS:SG	1:A:427:ILE:HB	2.28	0.72
1:A:309:VAL:CG1	1:A:370:ILE:CD1	2.65	0.72
1:B:125:ARG:HG3	1:B:160:ALA:HB3	1.70	0.72
1:B:412:CYS:SG	1:B:427:ILE:HB	2.31	0.71
1:A:44[B]:GLN:HG3	1:B:208:ARG:NH1	2.06	0.70
1:A:131:VAL:HA	1:A:192:THR:HG21	1.73	0.69
1:B:200:TRP:CE2	5:B:1501:GOL:H2	2.27	0.69
1:B:484[B]:CYS:SG	1:B:488:ARG:HB3	2.34	0.68
1:A:309:VAL:HG11	1:A:370:ILE:HG13	1.74	0.67
1:B:304[B]:SER:OG	1:B:448:VAL:HG21	1.98	0.64
1:B:412:CYS:SG	1:B:484[B]:CYS:HB2	2.40	0.60
1:B:455:ARG:HD3	1:B:464:LYS:HA	1.84	0.60
1:A:416:LYS:H	1:A:416:LYS:CD	2.16	0.58
1:A:484[B]:CYS:SG	1:A:488:ARG:HB3	2.44	0.57
1:B:202:ARG:NH2	5:B:1501:GOL:O3	2.37	0.57
1:A:416:LYS:HD3	1:A:416:LYS:N	2.19	0.57
1:B:106:ARG:NH1	1:B:141[B]:GLN:HG3	2.20	0.57
1:B:438:THR:HB	1:B:445:ASP:OD1	2.05	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:ARG:HG2	1:B:163:LYS:HB2	1.87	0.56
1:B:416:LYS:H	1:B:416:LYS:CE	2.19	0.55
1:B:416:LYS:H	1:B:416:LYS:CD	2.18	0.55
1:B:416:LYS:HD3	1:B:416:LYS:N	2.22	0.55
1:A:309:VAL:HG11	1:A:370:ILE:CG1	2.37	0.54
1:A:416:LYS:H	1:A:416:LYS:CE	2.20	0.54
1:A:240:ILE:CD1	1:A:277[B]:MET:SD	2.96	0.54
1:A:44[B]:GLN:HG3	1:B:208:ARG:HH12	1.73	0.53
1:A:412:CYS:SG	1:A:484[B]:CYS:HB2	2.47	0.53
1:A:416:LYS:HD3	1:A:416:LYS:H	1.72	0.53
1:B:240:ILE:CD1	1:B:277[B]:MET:SD	2.97	0.53
1:B:200:TRP:NE1	5:B:1501:GOL:H2	2.26	0.51
1:A:372:PRO:HG2	1:A:444:TYR:HB3	1.92	0.51
1:A:452:ASN:OD1	1:A:454:LYS:HE2	2.10	0.51
1:B:416:LYS:HD3	1:B:416:LYS:H	1.76	0.51
1:A:408[B]:CYS:SG	1:A:475:VAL:C	2.89	0.50
1:A:183:ILE:O	1:A:184:PHE:HB2	2.12	0.49
1:A:416:LYS:N	1:A:416:LYS:CD	2.75	0.49
1:A:455:ARG:HD3	1:A:464:LYS:HA	1.95	0.49
1:A:261:ASN:HB3	6:A:2298:HOH:O	2.13	0.48
1:B:183:ILE:O	1:B:184:PHE:HB2	2.13	0.48
1:B:416:LYS:N	1:B:416:LYS:CD	2.77	0.48
1:A:231:PHE:O	1:A:264:ALA:HA	2.14	0.47
1:A:222:GLN:NE2	3:A:1495:CL:CL	2.84	0.47
1:B:100:LYS:HD3	1:B:133:ASN:HD21	1.78	0.47
1:A:125[B]:ARG:NH1	6:A:2192:HOH:O	2.47	0.47
1:B:187:ILE:HB	1:B:221:ILE:HD13	1.97	0.47
1:B:408[B]:CYS:SG	1:B:475:VAL:C	2.94	0.46
1:B:231:PHE:O	1:B:264:ALA:HA	2.15	0.46
1:A:372:PRO:CG	1:A:444:TYR:HB3	2.46	0.46
1:B:222:GLN:NE2	3:B:1495:CL:CL	2.86	0.46
1:A:269[B]:CYS:HB3	1:A:275:ALA:CA	2.45	0.46
1:A:202:ARG:HA	1:A:227:ASP:O	2.16	0.46
1:B:174:ASN:HA	1:B:208:ARG:O	2.15	0.46
1:B:202:ARG:HA	1:B:227:ASP:O	2.17	0.45
1:B:365:LYS:O	1:B:369:GLY:HA2	2.17	0.45
1:A:416:LYS:HE2	6:A:2437:HOH:O	2.16	0.45
1:A:208[B]:ARG:HG3	5:A:1499:GOL:O3	2.16	0.45
1:A:408[B]:CYS:SG	1:A:476:CYS:N	2.89	0.45
1:A:269[B]:CYS:HB3	1:A:275:ALA:HA	1.98	0.45
1:A:135:VAL:O	1:A:167:VAL:HA	2.17	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:245:GLU:HA	1:B:279:GLY:O	2.17	0.44
1:B:135:VAL:O	1:B:167:VAL:HA	2.17	0.44
1:A:365:LYS:O	1:A:369:GLY:HA2	2.17	0.44
1:A:125[B]:ARG:NH1	6:A:2191:HOH:O	2.51	0.44
1:A:245:GLU:HA	1:A:279:GLY:O	2.18	0.44
1:A:187:ILE:HB	1:A:221:ILE:HD13	1.99	0.44
1:B:297:SER:HB2	1:B:388:GLY:HA3	1.99	0.43
1:B:261:ASN:HB3	6:B:2352:HOH:O	2.19	0.43
1:B:106:ARG:NH2	1:B:141[B]:GLN:OE1	2.52	0.43
1:B:297:SER:CB	1:B:388:GLY:HA3	2.48	0.43
1:B:300:SER:HA	1:B:432:ALA:O	2.19	0.43
1:A:138:PHE:O	1:A:170:TYR:HA	2.19	0.43
1:B:309:VAL:HG13	1:B:370:ILE:CD1	2.50	0.42
1:B:168:ARG:HA	1:B:202:ARG:O	2.19	0.42
1:A:240:ILE:HD13	1:A:277[B]:MET:SD	2.59	0.42
1:B:274:ALA:HA	1:B:300:SER:O	2.20	0.42
1:A:247:ASP:HB3	1:A:281:HIS:CG	2.54	0.42
1:A:304[A]:SER:OG	1:A:448:VAL:HG21	2.20	0.42
1:B:309:VAL:HG13	1:B:370:ILE:HD12	2.02	0.42
1:B:45:GLN:HB3	1:B:83:GLY:HA2	2.02	0.42
1:B:416:LYS:H	1:B:416:LYS:HE2	1.85	0.41
1:A:254:TYR:O	1:A:256:GLN:HG3	2.20	0.41
1:A:438:THR:HB	1:A:445:ASP:OD1	2.20	0.41
1:B:452:ASN:OD1	1:B:454:LYS:HE3	2.20	0.41
1:A:269[B]:CYS:SG	1:A:295:SER:HB3	2.60	0.41
1:A:95:LEU:HD22	1:A:116:THR:O	2.20	0.41
1:A:168:ARG:HA	1:A:202:ARG:O	2.20	0.41
1:B:173:SER:HA	1:B:207:GLU:O	2.20	0.41
1:A:309:VAL:CG1	1:A:370:ILE:CG1	2.97	0.41
1:B:138:PHE:O	1:B:170:TYR:HA	2.20	0.41
1:A:312:PHE:CD1	1:A:312:PHE:N	2.89	0.41
1:A:416:LYS:HE2	1:A:416:LYS:H	1.84	0.40
1:B:368:TYR:HB2	1:B:370:ILE:HG12	2.02	0.40
1:A:274:ALA:HA	1:A:300:SER:O	2.21	0.40
1:A:297:SER:CB	1:A:388:GLY:HA3	2.52	0.40
1:A:37:ASP:HB3	5:A:1500:GOL:H12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/464 (94%)	417 (95%)	21 (5%)	0	100	100
1	B	433/464 (93%)	412 (95%)	21 (5%)	0	100	100
All	All	871/928 (94%)	829 (95%)	42 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/399 (96%)	380 (99%)	5 (1%)	76	56
1	B	381/399 (96%)	377 (99%)	4 (1%)	82	67
All	All	766/798 (96%)	757 (99%)	9 (1%)	84	60

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125[A]	ARG
1	A	125[B]	ARG
1	A	190	ASP
1	A	309	VAL
1	A	366	LEU
1	B	125	ARG
1	B	190	ASP
1	B	277[A]	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	277[B]	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	222	GLN
1	B	60	ASN
1	B	195	ASN
1	B	222	GLN
1	B	261	ASN
1	B	422	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 14 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	1499	-	5,5,5	0.87	0	5,5,5	0.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	1500	-	5,5,5	0.90	0	5,5,5	0.28	0
5	GOL	A	1501	-	5,5,5	0.93	0	5,5,5	0.34	0
5	GOL	B	1499	-	5,5,5	0.88	0	5,5,5	0.34	0
5	GOL	B	1500	-	5,5,5	0.91	0	5,5,5	0.28	0
5	GOL	B	1501	-	5,5,5	0.76	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	1499	-	-	0/4/4/4	0/0/0/0
5	GOL	A	1500	-	-	0/4/4/4	0/0/0/0
5	GOL	A	1501	-	-	0/4/4/4	0/0/0/0
5	GOL	B	1499	-	-	0/4/4/4	0/0/0/0
5	GOL	B	1500	-	-	0/4/4/4	0/0/0/0
5	GOL	B	1501	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1499	GOL	1	0
5	A	1500	GOL	1	0
5	A	1501	GOL	1	0
5	B	1501	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	433/464 (93%)	0.37	31 (7%) 18 16	10, 18, 40, 48	19 (4%)
1	B	430/464 (92%)	0.42	35 (8%) 15 13	10, 16, 41, 49	36 (8%)
All	All	863/928 (92%)	0.40	66 (7%) 17 15	10, 17, 41, 49	55 (6%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	312	PHE	12.8
1	B	335	GLY	10.7
1	A	359	ALA	8.5
1	B	441	THR	7.8
1	B	366	LEU	7.8
1	B	359	ALA	7.5
1	A	312	PHE	6.6
1	B	370	ILE	6.6
1	B	336	CYS	6.5
1	A	366	LEU	6.5
1	B	28	VAL	5.9
1	B	444	TYR	5.8
1	B	369	GLY	5.4
1	B	365	LYS	5.2
1	A	368	TYR	5.0
1	A	313	SER	5.0
1	A	336	CYS	4.9
1	B	368	TYR	4.9
1	A	441	THR	4.6
1	B	352	ALA	4.6
1	A	412	CYS	4.4
1	A	369	GLY	4.3
1	A	360	CYS	4.3
1	B	339	THR	4.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	335	GLY	4.0
1	A	28	VAL	4.0
1	A	351	ALA	4.0
1	B	484[A]	CYS	3.9
1	B	361	LEU	3.7
1	A	370	ILE	3.7
1	B	29	SER	3.6
1	B	337	ALA	3.6
1	B	269[A]	CYS	3.5
1	B	338	GLN	3.5
1	A	362	ASP	3.5
1	B	412	CYS	3.5
1	B	362	ASP	3.4
1	B	358	ASP	3.4
1	B	364	ALA	3.2
1	B	310	GLU	3.2
1	B	407	MET	3.0
1	A	365	LYS	3.0
1	B	119	GLY	2.9
1	B	357	LYS	2.9
1	A	29	SER	2.8
1	A	444	TYR	2.8
1	A	310	GLU	2.8
1	B	367	GLU	2.7
1	A	407	MET	2.5
1	B	354	VAL	2.5
1	B	62	THR	2.5
1	B	360	CYS	2.5
1	B	439	PRO	2.5
1	A	364	ALA	2.5
1	A	153	ASP	2.4
1	B	195	ASN	2.4
1	A	122	LYS	2.4
1	A	62	THR	2.4
1	A	358	ASP	2.4
1	A	338	GLN	2.4
1	A	339	THR	2.4
1	B	363	LYS	2.3
1	A	269[A]	CYS	2.3
1	A	439	PRO	2.1
1	A	371	GLU	2.0
1	A	309	VAL	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	B	1501	6/6	0.62	0.32	10.71	51,53,53,53	0
5	GOL	A	1501	6/6	0.36	0.31	6.66	51,53,53,53	0
5	GOL	B	1499	6/6	0.76	0.24	5.41	52,53,53,53	0
5	GOL	B	1500	6/6	0.60	0.29	4.55	51,53,53,54	0
5	GOL	A	1500	6/6	0.50	0.30	4.48	51,53,53,54	0
5	GOL	A	1499	6/6	0.76	0.26	3.80	52,53,53,53	0
3	CL	A	1495	1/1	0.99	0.13	1.95	28,28,28,28	0
4	NA	A	1498	1/1	0.92	0.12	0.81	27,27,27,27	0
3	CL	B	1495	1/1	0.98	0.08	-0.09	27,27,27,27	0
2	CA	A	1494	1/1	0.95	0.13	-0.43	27,27,27,27	0
4	NA	B	1497	1/1	0.99	0.07	-1.00	13,13,13,13	0
2	CA	A	1493	1/1	0.99	0.04	-1.62	16,16,16,16	0
2	CA	B	1493	1/1	0.98	0.04	-1.62	15,15,15,15	0
4	NA	A	1497	1/1	0.99	0.05	-2.03	16,16,16,16	0
2	CA	B	1494	1/1	0.96	0.08	-2.14	26,26,26,26	0
2	CA	A	1492	1/1	0.99	0.04	-	15,15,15,15	0
4	NA	A	1496	1/1	0.97	0.10	-	15,15,15,15	0
2	CA	B	1492	1/1	0.99	0.04	-	15,15,15,15	0
4	NA	B	1498	1/1	0.96	0.18	-	25,25,25,25	0
4	NA	B	1496	1/1	0.99	0.04	-	15,15,15,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.