



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:08 AM GMT

PDB ID : 3H9H
Title : Human Class I MHC HLA-A2(A150P) in complex with the Tel1p peptide
Authors : Borbulevych, O.Y.; Baker, B.M.
Deposited on : 2009-04-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>9%</div> <div>93%</div> <div>7%</div> </div>
1	D	275	<div> <div>9%</div> <div>93%</div> <div>6%</div> </div>
2	B	100	<div> <div>4%</div> <div>92%</div> <div>7%</div> </div>
2	E	100	<div> <div>4%</div> <div>96%</div> <div></div> </div>
3	C	9	<div> <div>11%</div> <div>78%</div> <div>22%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: a red segment on the left labeled '11%', a green segment in the middle labeled '89%', and a yellow segment on the right labeled '11%'.

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	6	0
			2282	1428	415	430	9			
1	D	275	Total	C	N	O	S	0	9	0
			2292	1436	414	433	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	150	PRO	ALA	ENGINEERED	UNP P01892
D	150	PRO	ALA	ENGINEERED	UNP P01892

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	1	0
			842	536	141	161	4			
2	E	100	Total	C	N	O	S	0	1	0
			840	535	141	160	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	EXPRESSION TAG	UNP P61769
E	0	MET	-	EXPRESSION TAG	UNP P61769

- Molecule 3 is a protein called Tel1p peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	0	0
			83	58	11	13	1			
3	F	9	Total	C	N	O	S	0	0	0
			83	58	11	13	1			


- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	176	Total	O	0	0
			176	176		
5	B	92	Total	O	0	0
			92	92		
5	C	3	Total	O	0	0
			3	3		
5	D	171	Total	O	0	0
			171	171		
5	E	85	Total	O	0	2
			87	87		
5	F	6	Total	O	0	0
			6	6		



- Molecule 3: Tel1p peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.16Å 86.89Å 79.41Å 90.00° 89.97° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 19.89 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.1 (20.00-2.00) 97.1 (19.89-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.171 , 0.226 0.176 , 0.229	Depositor DCC
R_{free} test set	2841 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.9	EDS
Estimated twinning fraction	0.155 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 56055 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6987	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/2366	0.85	4/3210 (0.1%)
1	D	0.76	0/2385	0.82	2/3236 (0.1%)
2	B	0.89	3/868 (0.3%)	0.86	3/1173 (0.3%)
2	E	0.80	1/866 (0.1%)	0.78	0/1170
3	C	0.82	0/86	0.96	0/115
3	F	0.80	0/86	0.79	0/115
All	All	0.79	4/6657 (0.1%)	0.83	9/9019 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	80	CYS	CB-SG	-6.83	1.70	1.82
2	E	80	CYS	CB-SG	-5.33	1.73	1.81
2	B	9	VAL	CB-CG2	-5.12	1.42	1.52
2	B	82	VAL	CB-CG2	5.08	1.63	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	97	ARG	NE-CZ-NH1	6.77	123.69	120.30
1	A	14	ARG	NE-CZ-NH2	-6.13	117.24	120.30
1	D	106	ASP	CB-CG-OD1	6.02	123.72	118.30
2	B	97	ARG	NE-CZ-NH2	-5.95	117.32	120.30
1	A	181	ARG	NE-CZ-NH2	5.79	123.19	120.30
1	A	37	ASP	CB-CG-OD1	5.70	123.43	118.30
2	B	12	ARG	NE-CZ-NH1	-5.35	117.63	120.30
2	B	81	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	97	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2282	0	2151	6	0
1	D	2292	0	2169	5	0
2	B	842	0	807	2	0
2	E	840	0	808	1	0
3	C	83	0	81	0	0
3	F	83	0	81	0	0
4	B	12	0	16	1	0
4	D	12	0	16	0	0
4	E	6	0	8	0	0
5	A	176	0	0	1	0
5	B	92	0	0	0	0
5	C	3	0	0	0	0
5	D	171	0	0	0	0
5	E	87	0	0	0	0
5	F	6	0	0	0	0
All	All	6987	0	6137	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:186[A]:LYS:HD2	1:D:207[A]:SER:HB2	1.81	0.61
1:A:19:GLU:OE1	1:A:75[A]:ARG:NH1	2.35	0.59
1:D:14:ARG:HB2	1:D:17:ARG:HB2	1.95	0.48
1:D:74:HIS:CE1	1:D:97:ARG:HE	2.32	0.47
1:A:111:ARG:NH1	5:A:379:HOH:O	2.40	0.47
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.50	0.47
1:A:6:ARG:NE	1:A:113:TYR:OH	2.46	0.47
1:A:20:PRO:O	1:A:75[A]:ARG:NH2	2.48	0.46
1:D:133:TRP:HB2	1:D:144:LYS:HG3	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.53	0.43
2:B:97:ARG:HD2	4:B:101:GOL:H2	2.01	0.41
1:A:51:TRP:CZ2	1:A:179:LEU:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/275 (102%)	274 (98%)	4 (1%)	1 (0%)	39	33
1	D	282/275 (102%)	276 (98%)	5 (2%)	1 (0%)	39	33
2	B	99/100 (99%)	98 (99%)	1 (1%)	0	100	100
2	E	99/100 (99%)	98 (99%)	1 (1%)	0	100	100
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	F	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	773/768 (101%)	758 (98%)	13 (2%)	2 (0%)	46	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	150	PRO
1	A	150	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/232 (103%)	229 (96%)	9 (4%)	40	36
1	D	241/232 (104%)	234 (97%)	7 (3%)	50	49
2	B	96/95 (101%)	95 (99%)	1 (1%)	82	85
2	E	96/95 (101%)	94 (98%)	2 (2%)	61	63
3	C	8/8 (100%)	6 (75%)	2 (25%)	1	0
3	F	8/8 (100%)	7 (88%)	1 (12%)	6	3
All	All	687/670 (102%)	665 (97%)	22 (3%)	47	44

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ARG
1	A	75[A]	ARG
1	A	75[B]	ARG
1	A	88	SER
1	A	113	TYR
1	A	152	VAL
1	A	157	ARG
1	A	196	ASP
1	A	225	THR
2	B	70	PHE
3	C	1	MET
3	C	6	LEU
1	D	35	ARG
1	D	111	ARG
1	D	121	LYS
1	D	128	GLU
1	D	141	GLN
1	D	223	ASP
1	D	225	THR
2	E	0	MET
2	E	70	PHE
3	F	1	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	141	GLN
1	D	145	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	100	-	5,5,5	0.35	0	5,5,5	1.09	0
4	GOL	B	101	-	5,5,5	0.55	0	5,5,5	1.13	0
4	GOL	D	276	-	5,5,5	0.56	0	5,5,5	0.24	0
4	GOL	D	277	-	5,5,5	0.37	0	5,5,5	0.70	0
4	GOL	E	100	-	5,5,5	0.19	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	100	-	-	0/4/4/4	0/0/0/0
4	GOL	B	101	-	-	0/4/4/4	0/0/0/0
4	GOL	D	276	-	-	0/4/4/4	0/0/0/0
4	GOL	D	277	-	-	0/4/4/4	0/0/0/0
4	GOL	E	100	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	101	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.52	24 (8%)	13 13	26, 34, 48, 57	0
1	D	275/275 (100%)	0.57	26 (9%)	10 11	25, 34, 49, 67	0
2	B	100/100 (100%)	0.31	4 (4%)	42 44	28, 35, 43, 50	0
2	E	100/100 (100%)	0.35	4 (4%)	42 44	30, 34, 43, 49	0
3	C	9/9 (100%)	1.29	1 (11%)	7 8	33, 40, 52, 53	0
3	F	9/9 (100%)	1.44	1 (11%)	7 8	32, 39, 53, 55	0
All	All	768/768 (100%)	0.51	60 (7%)	16 17	25, 34, 48, 67	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	151	HIS	6.5
1	D	151	HIS	5.8
3	F	5	TYR	5.8
3	C	5	TYR	5.2
1	A	226	GLN	4.2
1	A	150	PRO	4.1
1	D	17	ARG	4.1
1	D	150	PRO	4.0
1	A	18	GLY	3.6
1	D	197	HIS	3.5
1	D	275	GLU	3.5
1	A	225	THR	3.3
1	D	196	ASP	3.2
1	D	54	GLN	3.2
2	E	48	LYS	3.2
1	D	217	TRP	3.1
1	D	225	THR	3.1
1	D	226	GLN	3.0
1	D	41	ALA	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	95[A]	VAL	3.0
1	A	196	ASP	2.9
1	A	197	HIS	2.9
1	A	275	GLU	2.9
2	B	89	GLN	2.8
1	A	136	ALA	2.8
1	A	95[A]	VAL	2.8
1	D	50	PRO	2.7
1	A	268	LYS	2.7
2	B	75	LYS	2.7
1	D	157	ARG	2.7
1	D	136	ALA	2.7
1	D	16	GLY	2.7
2	B	23	LEU	2.6
1	A	86	ASN	2.6
1	A	16	GLY	2.5
1	D	152	VAL	2.5
1	A	54	GLN	2.5
1	A	108	ARG	2.5
1	A	19	GLU	2.4
1	A	28	VAL	2.4
1	A	128	GLU	2.4
1	D	10	THR	2.3
1	D	192	HIS	2.3
1	D	249	VAL	2.3
1	A	195	SER	2.3
1	D	79	GLY	2.3
2	E	23	LEU	2.3
1	A	157	ARG	2.2
1	A	133	TRP	2.2
2	E	75	LYS	2.2
2	B	25	CYS	2.2
1	A	34	VAL	2.2
1	D	193	ALA	2.2
2	E	27	VAL	2.2
1	D	22	PHE	2.1
1	D	113	TYR	2.1
1	A	261	VAL	2.1
1	D	19	GLU	2.1
1	D	201	LEU	2.1
1	A	33	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GOL	B	101	6/6	0.86	0.24	1.56	57,58,59,60	0
4	GOL	D	276	6/6	0.88	0.16	1.49	52,53,54,55	0
4	GOL	E	100	6/6	0.89	0.27	1.32	41,48,49,50	0
4	GOL	B	100	6/6	0.79	0.17	0.94	52,53,53,54	0
4	GOL	D	277	6/6	0.89	0.19	-	64,64,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.