



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:07 PM GMT

PDB ID : 4HA0
Title : Structure of Geobacillus kaustophilus lactonase, mutant R230D with Zn²⁺
Authors : Xue, B.; Chow, J.Y.; Yew, W.S.; Robinson, R.C.
Deposited on : 2012-09-25
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	 19% 80% 17% .
1	B	330	 6% 82% 15% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	OH	A	403	-	-	-	X
4	OH	B	403	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphotriesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2540	1615	432	477	16			
1	B	323	Total	C	N	O	S	0	0	0
			2540	1615	432	477	16			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP Q5KZU5
A	-2	SER	-	EXPRESSION TAG	UNP Q5KZU5
A	-1	HIS	-	EXPRESSION TAG	UNP Q5KZU5
A	0	ASN	-	EXPRESSION TAG	UNP Q5KZU5
A	230	ASP	ARG	ENGINEERED MUTATION	UNP Q5KZU5
B	-3	GLY	-	EXPRESSION TAG	UNP Q5KZU5
B	-2	SER	-	EXPRESSION TAG	UNP Q5KZU5
B	-1	HIS	-	EXPRESSION TAG	UNP Q5KZU5
B	0	ASN	-	EXPRESSION TAG	UNP Q5KZU5
B	230	ASP	ARG	ENGINEERED MUTATION	UNP Q5KZU5

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is HYDROXIDE ION (three-letter code: OH) (formula: HO).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			1	1		
4	B	1	Total	O	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	137	Total	O	0	0
			137	137		
5	B	146	Total	O	0	0
			146	146		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphotriesterase



• Molecule 1: Phosphotriesterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.97Å 76.22Å 134.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.98 – 1.90 29.97 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (19.98-1.90) 97.4 (29.97-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.259 , 0.303 0.267 , 0.303	Depositor DCC
R_{free} test set	2850 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	31.1	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 51.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	3 of 56078 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5369	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 89.56 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.2567e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, OH, KCX, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2586	0.62	1/3500 (0.0%)
1	B	0.62	1/2586 (0.0%)	0.62	0/3500
All	All	0.62	1/5172 (0.0%)	0.62	1/7000 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	100	TYR	CD1-CE1	-5.16	1.31	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	115	LEU	CA-CB-CG	5.37	127.65	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2540	0	2477	33	0
1	B	2540	0	2478	29	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	137	0	0	1	1
5	B	146	0	0	0	0
All	All	5369	0	4955	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:GLN:HG2	1:A:241:THR:HA	1.60	0.82
1:B:259:LYS:HG2	1:B:314:GLN:HE21	1.56	0.71
1:B:100:TYR:CE2	1:B:102:GLY:HA3	2.34	0.62
1:A:78:PRO:HG2	1:A:134:ILE:HG21	1.84	0.59
1:B:78:PRO:HG2	1:B:134:ILE:HG21	1.86	0.57
1:B:112:ARG:HA	1:B:115:LEU:HG	1.89	0.55
1:B:208:CYS:HB3	1:B:228:PHE:CD2	2.42	0.54
1:A:208:CYS:HB3	1:A:228:PHE:CD2	2.43	0.53
1:A:265:HIS:HB2	1:A:267:THR:HG23	1.90	0.53
1:B:265:HIS:HB2	1:B:267:THR:HG23	1.92	0.51
1:A:43:ARG:HA	5:A:593:HOH:O	2.10	0.51
1:B:250:LEU:HD11	1:B:305:GLU:HG3	1.91	0.51
1:A:250:LEU:HD11	1:A:305:GLU:HG3	1.95	0.49
1:B:282:PHE:CD1	1:B:282:PHE:N	2.81	0.48
1:B:210:ASN:O	1:B:216:HIS:HE1	1.97	0.47
1:A:237:VAL:HG12	1:A:237:VAL:O	2.15	0.47
1:A:210:ASN:O	1:A:216:HIS:HE1	1.97	0.47
1:B:106:PRO:O	1:B:110:GLN:HG3	2.14	0.47
1:B:19:LYS:HG2	1:B:63:ILE:HD13	1.96	0.47
1:A:111:PHE:O	1:A:114:LEU:HB2	2.15	0.47
1:A:282:PHE:N	1:A:282:PHE:CD1	2.81	0.47
1:A:19:LYS:HG2	1:A:63:ILE:HD13	1.97	0.47
1:B:152:ARG:HE	1:B:152:ARG:HB2	1.51	0.46
1:A:207:MET:O	1:A:216:HIS:NE2	2.48	0.46
1:B:55:ALA:HB3	1:B:88:THR:HG21	1.98	0.46
1:A:239:ALA:HA	1:A:240:PRO:HD2	1.83	0.45
1:A:55:ALA:HB3	1:A:88:THR:HG21	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:VAL:HG22	1:A:225:TYR:HB2	1.97	0.45
1:B:145:KCX:OQ2	1:B:178:HIS:HB2	2.17	0.45
1:B:203:VAL:HG22	1:B:225:TYR:HB2	1.99	0.45
1:A:57:LYS:HD3	1:A:292:GLU:OE1	2.17	0.45
1:B:57:LYS:HD3	1:B:292:GLU:OE1	2.17	0.44
1:A:39:ARG:NE	1:B:124:ASP:OD2	2.36	0.44
1:A:100:TYR:CE2	1:A:102:GLY:HA3	2.53	0.44
1:B:303:LYS:HE2	1:B:303:LYS:HB3	1.68	0.44
1:A:277:THR:HG22	1:A:278:LEU:N	2.32	0.44
1:A:152:ARG:HE	1:A:152:ARG:HB2	1.51	0.44
1:A:282:PHE:HD1	1:A:282:PHE:N	2.16	0.44
1:B:130:LEU:HD11	1:B:144:ILE:HD11	1.99	0.44
1:A:106:PRO:O	1:A:110:GLN:HG3	2.18	0.44
1:A:41:THR:CG2	1:A:42:PHE:N	2.81	0.43
1:B:296:VAL:HG23	1:B:297:ASN:OD1	2.18	0.43
1:A:207:MET:HG3	1:A:226:ILE:HB	2.01	0.43
1:A:299:ILE:HB	1:A:300:PRO:HD3	2.01	0.43
1:A:277:THR:CG2	1:A:278:LEU:N	2.81	0.43
1:B:8:VAL:HA	1:B:139:ILE:HG23	2.01	0.43
1:B:299:ILE:HB	1:B:300:PRO:HD3	2.01	0.43
1:B:55:ALA:HB1	1:B:90:LEU:HD22	2.00	0.43
1:A:61:HIS:CD2	1:A:295:PHE:HB2	2.53	0.43
1:B:207:MET:O	1:B:216:HIS:NE2	2.52	0.42
1:A:280:GLU:N	1:A:281:PRO:HD2	2.35	0.42
1:B:282:PHE:N	1:B:282:PHE:HD1	2.15	0.41
1:B:207:MET:HG3	1:B:226:ILE:HB	2.02	0.41
1:A:3:GLU:HG3	1:A:4:MET:HG2	2.03	0.41
1:A:60:ARG:C	1:A:61:HIS:ND1	2.74	0.41
1:B:39:ARG:HH11	1:B:39:ARG:HD3	1.76	0.41
1:A:303:LYS:HE2	1:A:303:LYS:HB3	1.70	0.41
1:A:8:VAL:HA	1:A:139:ILE:HG23	2.02	0.41
1:B:41:THR:CG2	1:B:42:PHE:N	2.83	0.40
1:B:3:GLU:HG3	1:B:4:MET:HG2	2.04	0.40
1:A:259:LYS:HG2	1:A:314:GLN:HE21	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:558:HOH:O	5:A:559:HOH:O 3_545	2.08	0.12

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/330 (97%)	313 (98%)	6 (2%)	1 (0%)	46	35
1	B	320/330 (97%)	312 (98%)	7 (2%)	1 (0%)	46	35
All	All	640/660 (97%)	625 (98%)	13 (2%)	2 (0%)	46	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	PRO
1	B	69	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/267 (98%)	259 (98%)	4 (2%)	72	69
1	B	263/267 (98%)	259 (98%)	4 (2%)	72	69
All	All	526/534 (98%)	518 (98%)	8 (2%)	72	69

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	HIS
1	A	99	TYR
1	A	178	HIS
1	A	264	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	41	THR
1	B	99	TYR
1	B	115	LEU
1	B	264	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	314	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	A	145	1,3,2	7,11,12	0.74	0	7,12,14	2.00	1 (14%)
1	KCX	B	145	1,3,2	7,11,12	1.04	0	7,12,14	1.76	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	145	1,3,2	-	0/6/10/12	0/0/0/0
1	KCX	B	145	1,3,2	-	0/6/10/12	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	KCX	CE-NZ-CX	-4.86	117.99	123.49
1	B	145	KCX	CE-NZ-CX	-4.05	118.91	123.49
1	B	145	KCX	O-C-CA	-2.03	120.20	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	145	KCX	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are modelled with single atom and 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/330 (97%)	1.08	63 (19%) 1 1	20, 50, 85, 128	0
1	B	322/330 (97%)	0.54	19 (5%) 26 29	21, 38, 62, 83	0
All	All	644/660 (97%)	0.81	82 (12%) 5 5	20, 43, 82, 128	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	237	VAL	8.6
1	B	238	GLY	6.5
1	A	227	ALA	6.1
1	A	263	LEU	5.6
1	A	176	ILE	5.0
1	A	262	MET	4.9
1	B	280	GLU	4.9
1	A	237	VAL	4.9
1	A	22	ILE	4.8
1	B	236	MET	4.7
1	A	280	GLU	4.6
1	A	203	VAL	4.6
1	A	226	ILE	4.5
1	B	282	PHE	4.5
1	A	177	THR	4.1
1	A	205	GLY	3.9
1	A	204	ILE	3.8
1	B	285	MET	3.7
1	A	229	ASP	3.6
1	A	261	ILE	3.5
1	A	146	LEU	3.5
1	A	236	MET	3.4
1	A	225	TYR	3.4
1	B	283	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	235	GLY	3.3
1	A	264	SER	3.3
1	B	22	ILE	3.3
1	A	289	TRP	3.3
1	A	42	PHE	3.2
1	A	291	VAL	3.2
1	A	231	PHE	3.2
1	B	175	ILE	3.1
1	A	52	VAL	3.0
1	A	238	GLY	3.0
1	A	295	PHE	2.8
1	A	179	THR	2.8
1	A	298	ILE	2.8
1	A	294	LEU	2.8
1	A	67	VAL	2.8
1	A	70	THR	2.7
1	A	178	HIS	2.7
1	A	144	ILE	2.7
1	A	282	PHE	2.7
1	B	42	PHE	2.7
1	A	285	MET	2.7
1	A	243	GLU	2.6
1	A	232	GLY	2.6
1	B	278	LEU	2.6
1	B	16	GLN	2.6
1	A	296	VAL	2.6
1	A	224	VAL	2.5
1	A	21	LEU	2.5
1	A	60	ARG	2.5
1	A	93	ILE	2.5
1	A	16	GLN	2.5
1	B	146	LEU	2.5
1	B	279	PRO	2.4
1	A	265	HIS	2.4
1	A	88	THR	2.4
1	A	74	CYS	2.3
1	A	45	ASP	2.3
1	A	311	VAL	2.3
1	A	228	PHE	2.3
1	A	95	ALA	2.3
1	A	211	THR	2.3
1	A	268	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	147	ALA	2.2
1	A	305	GLU	2.2
1	B	43	ARG	2.2
1	B	176	ILE	2.2
1	A	69	PRO	2.1
1	A	323	LEU	2.1
1	A	287	LYS	2.1
1	A	266	ASP	2.1
1	A	247	ARG	2.1
1	A	143	VAL	2.1
1	A	49	ARG	2.1
1	A	207	MET	2.1
1	B	284	GLU	2.1
1	A	23	HIS	2.1
1	B	144	ILE	2.1
1	A	175	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	KCX	B	145	12/13	0.94	0.21	-	20,23,30,39	0
1	KCX	A	145	12/13	0.93	0.23	-	29,32,44,45	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	OH	A	403	1/1	0.90	0.39	5.02	41,41,41,41	0
4	OH	B	403	1/1	0.96	0.25	3.44	34,34,34,34	0
2	FE	B	401	1/1	1.00	0.11	-1.39	23,23,23,23	0
2	FE	A	401	1/1	0.97	0.11	-3.60	29,29,29,29	0
3	ZN	B	402	1/1	0.99	0.10	-	30,30,30,30	1
3	ZN	A	402	1/1	0.99	0.12	-	33,33,33,33	1

6.5 Other polymers [i](#)

There are no such residues in this entry.