



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:09 AM GMT

PDB ID : 3HCP  
Title : Human ferrochelatase with Mn and deuteroporphyrin bound  
Authors : Medlock, A.E.; Dailey, H.A.; Lanzilotta, W.N.  
Deposited on : 2009-05-06  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 1%, green 88%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>88%</span> <span>11%</span> </div> </div>
1	B	359	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 13%, green 86%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>86%</span> <span>13%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CHD	A	3	-	-	-	X
3	CHD	B	4	-	-	-	X
4	GOL	A	1	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6583 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ferrochelatase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	359	Total	C	N	O	S	0	3	0
			2901	1845	506	532	18			
1	B	359	Total	C	N	O	S	0	9	0
			2930	1871	505	536	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	PHE	VARIANT	UNP P22830
B	610	ALA	PHE	VARIANT	UNP P22830

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 3 is CHOLIC ACID (three-letter code: CHD) (formula:  $C_{24}H_{40}O_5$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			29	24	5		
3	A	1	Total	C	O	0	0
			29	24	5		
3	B	1	Total	C	O	0	0
			29	24	5		
3	B	1	Total	C	O	0	0
			29	24	5		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			5	3	2		
5	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 6 is FE(III) DEUTEROPORPHYRIN IX (three-letter code: FDE) (formula:  $C_{30}H_{28}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	Fe	N	O	0	0
			39	30	1	4	4		
6	B	1	Total	C	Fe	N	O	0	0
			39	30	1	4	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	280	Total	O	0	0
			280	280		
7	B	254	Total	O	0	0
			254	254		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ferrochelatase, mitochondrial



- Molecule 1: Ferrochelatase, mitochondrial





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.57Å 93.69Å 109.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.83 – 2.00 46.42 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.83-2.00) 99.3 (46.42-2.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.90 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.170 , 0.201 0.173 , 0.176	Depositor DCC
$R_{free}$ test set	3077 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.8	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	5 of 60758 reflections (0.008%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6583	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.11 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.0403e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FDE, CHD, IMD, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	0/2979	0.73	4/4035 (0.1%)
1	B	0.72	0/3027	0.71	4/4099 (0.1%)
All	All	0.73	0/6006	0.72	8/8134 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	ARG	NE-CZ-NH1	8.29	124.45	120.30
1	A	226	ARG	NE-CZ-NH2	-7.88	116.36	120.30
1	B	726	ARG	NE-CZ-NH1	6.96	123.78	120.30
1	A	91	ARG	NE-CZ-NH1	6.44	123.52	120.30
1	A	91	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	B	726	ARG	NE-CZ-NH2	-6.01	117.30	120.30
1	B	591	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	B	772	ARG	NE-CZ-NH1	-5.38	117.61	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2901	0	2917	30	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2930	0	2966	37	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	58	0	78	2	0
3	B	58	0	78	3	0
4	A	6	0	8	0	0
5	A	5	0	4	0	0
5	B	5	0	4	0	0
6	A	39	0	25	2	0
6	B	39	0	24	2	0
7	A	280	0	0	4	0
7	B	254	0	0	6	0
All	All	6583	0	6104	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:398:GLN:HE22	1:B:797:TYR:H	1.18	0.91
1:A:297:TYR:H	1:B:898:GLN:HE22	1.23	0.84
1:A:263:HIS:NE2	1:A:343:GLU:HG3	1.93	0.83
1:A:383[B]:ASP:OD2	7:A:505:HOH:O	1.98	0.81
1:B:623:TYR:OH	6:B:925:FDE:O2A	1.97	0.81
1:B:763:HIS:HD2	7:B:84:HOH:O	1.68	0.75
1:B:843:GLU:OE2	7:B:523:HOH:O	2.09	0.70
1:B:823:CYS:SG	1:B:862[B]:VAL:HG12	2.32	0.69
1:A:175:GLU:O	1:A:179:ARG:HG2	1.94	0.67
1:B:922:GLN:O	1:B:923:LEU:HB2	1.97	0.65
1:A:286:LYS:NZ	7:A:632:HOH:O	2.30	0.63
1:B:912:ARG:NH1	1:B:913:GLU:OE2	2.32	0.61
1:A:398:GLN:HE22	1:B:797:TYR:N	1.96	0.58
1:A:422:GLN:O	1:A:423:LEU:HB2	2.04	0.57
1:B:874:ASN:HD22	1:B:874:ASN:C	2.07	0.57
1:A:263:HIS:CE1	1:A:343:GLU:HG3	2.39	0.57
1:B:720[B]:LYS:HE2	1:B:923:LEU:HD13	1.86	0.56
1:A:297:TYR:N	1:B:898:GLN:HE22	2.01	0.56
1:B:720[A]:LYS:HZ1	1:B:894:LEU:CD2	2.18	0.56
3:A:2:CHD:H183	3:A:2:CHD:H212	1.87	0.56
1:A:179:ARG:HH11	1:A:179:ARG:HG3	1.71	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:273:GLY:HA2	1:B:813:PRO:HG3	1.88	0.56
1:A:412:ARG:NH1	1:A:413:GLU:OE2	2.40	0.54
1:A:123:TYR:OH	6:A:425:FDE:O2A	2.21	0.54
1:B:615:ARG:HD2	3:B:1:CHD:H232	1.89	0.53
1:A:288:MET:HG3	1:A:297:TYR:CD2	2.44	0.53
1:A:226:ARG:HD3	1:A:279:GLU:OE2	2.09	0.53
1:A:374:ASN:HD22	1:A:374:ASN:C	2.11	0.53
1:A:374:ASN:ND2	1:A:376:LEU:H	2.07	0.52
1:A:179:ARG:NH1	1:A:179:ARG:HG3	2.24	0.52
1:A:216:LYS:HG3	1:A:217:PRO:HD2	1.92	0.52
1:B:829:ASN:HD22	1:B:864:ASN:HB2	1.75	0.51
1:B:837[B]:PHE:CD2	1:B:841:HIS:HE1	2.28	0.51
1:B:732:LEU:HD22	1:B:876:LEU:HD11	1.93	0.51
1:A:329:ASN:HD22	1:A:364:ASN:HB2	1.76	0.51
1:A:300:VAL:HG12	1:A:313:PRO:HG2	1.93	0.50
1:B:837[B]:PHE:CD2	1:B:841:HIS:CE1	3.00	0.50
1:B:726:ARG:HD3	1:B:779:GLU:OE2	2.12	0.49
1:B:859[A]:GLU:HG3	7:B:496:HOH:O	2.11	0.49
1:B:673:ALA:O	1:B:677:MET:HG3	2.11	0.49
1:B:834:PRO:HG2	7:B:210:HOH:O	2.13	0.49
1:B:857:ALA:HB1	1:B:862[B]:VAL:HG11	1.95	0.48
1:A:374:ASN:HD22	1:A:375:PRO:N	2.11	0.48
1:B:614:ARG:HG3	7:B:358:HOH:O	2.11	0.48
3:B:1:CHD:H192	6:B:925:FDE:HMF1	1.95	0.47
1:B:608:ALA:HB3	1:B:609:PRO:HD3	1.95	0.47
1:B:816:ASP:HB3	1:B:852:TYR:CE1	2.50	0.47
1:B:874:ASN:HD22	1:B:875:PRO:N	2.13	0.47
1:A:374:ASN:HD22	1:A:376:LEU:H	1.63	0.47
1:A:221:TRP:H	1:A:421:GLN:NE2	2.13	0.46
1:B:788:MET:HG3	1:B:797:TYR:CD2	2.53	0.44
1:B:720[B]:LYS:HG2	7:B:482:HOH:O	2.17	0.43
1:A:303:SER:HB3	1:A:347:GLU:OE2	2.18	0.43
1:B:677:MET:HB3	1:B:682:LEU:HD12	2.00	0.43
1:A:288:MET:HG3	1:A:297:TYR:CE2	2.54	0.42
1:B:720[B]:LYS:CE	1:B:923:LEU:HD13	2.50	0.42
7:A:632:HOH:O	1:B:785:GLN:CG	2.67	0.42
1:A:215:ARG:HD2	7:A:582:HOH:O	2.19	0.42
1:A:313:PRO:HG3	1:B:773:GLY:HA2	2.01	0.42
1:B:599:MET:HB2	3:B:1:CHD:H62	2.02	0.41
3:A:2:CHD:H192	6:A:425:FDE:HMF1	2.02	0.41
1:B:820:LYS:O	1:B:824:GLU:HG3	2.21	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:800:VAL:HG12	1:B:813:PRO:HG2	2.04	0.40
1:A:316:ASP:HB3	1:A:352:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/359 (100%)	354 (98%)	6 (2%)	0	100	100
1	B	366/359 (102%)	357 (98%)	8 (2%)	1 (0%)	46	41
All	All	726/718 (101%)	711 (98%)	14 (2%)	1 (0%)	56	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	603	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/323 (101%)	316 (97%)	10 (3%)	47	46
1	B	332/323 (103%)	321 (97%)	11 (3%)	45	43
All	All	658/646 (102%)	637 (97%)	21 (3%)	47	44

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	LYS
1	A	106	LYS
1	A	182	LEU
1	A	216	LYS
1	A	219	MET
1	A	259	LEU
1	A	305	VAL
1	A	354	GLN
1	A	362	VAL
1	A	374	ASN
1	B	684	ARG
1	B	712	GLN
1	B	713	VAL
1	B	719	MET
1	B	720[A]	LYS
1	B	720[B]	LYS
1	B	727	TRP
1	B	759	LEU
1	B	841	HIS
1	B	843	GLU
1	B	874	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	HIS
1	A	329	ASN
1	A	364	ASN
1	A	374	ASN
1	A	398	GLN
1	A	421	GLN
1	B	653	ASN
1	B	763	HIS
1	B	802	GLN
1	B	814	GLN
1	B	829	ASN
1	B	841	HIS
1	B	854	GLN
1	B	864	ASN
1	B	874	ASN
1	B	898	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	921	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	1	-	5,5,5	0.49	0	5,5,5	1.51	2 (40%)
3	CHD	A	2	-	29,32,32	0.54	0	48,51,51	1.32	6 (12%)
3	CHD	A	3	-	29,32,32	0.69	0	48,51,51	1.28	6 (12%)
5	IMD	A	424	6	3,5,5	0.62	0	4,5,5	0.98	0
6	FDE	A	425	5	20,46,46	1.86	7 (35%)	20,76,76	2.64	6 (30%)
2	FES	A	501	1	0,4,4	0.00	-	0,4,4	0.00	-
3	CHD	B	1	-	29,32,32	0.60	0	48,51,51	1.41	10 (20%)
3	CHD	B	4	-	29,32,32	0.75	0	48,51,51	1.41	10 (20%)
2	FES	B	501	1	0,4,4	0.00	-	0,4,4	0.00	-
5	IMD	B	924	6	3,5,5	0.48	0	4,5,5	0.47	0
6	FDE	B	925	5	20,46,46	2.08	8 (40%)	20,76,76	2.92	8 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1	-	-	0/4/4/4	0/0/0/0
3	CHD	A	2	-	-	0/7/74/74	0/4/4/4
3	CHD	A	3	-	-	0/7/74/74	0/4/4/4
5	IMD	A	424	6	-	0/0/0/0	0/1/1/1
6	FDE	A	425	5	-	0/6/50/50	0/0/8/8
2	FES	A	501	1	-	0/0/4/4	0/1/1/1
3	CHD	B	1	-	-	0/7/74/74	0/4/4/4
3	CHD	B	4	-	-	0/7/74/74	0/4/4/4
2	FES	B	501	1	-	0/0/4/4	0/1/1/1
5	IMD	B	924	6	-	0/0/0/0	0/1/1/1
6	FDE	B	925	5	-	0/6/50/50	0/0/8/8

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	925	FDE	C1D-CHD	-2.73	1.32	1.39
6	B	925	FDE	C4B-NB	-2.67	1.33	1.36
6	A	425	FDE	C1B-NB	-2.55	1.33	1.36
6	A	425	FDE	C1D-CHD	-2.44	1.33	1.39
6	B	925	FDE	C1B-NB	-2.28	1.33	1.36
6	A	425	FDE	C4B-NB	-2.16	1.33	1.36
6	B	925	FDE	C4C-NC	-2.02	1.34	1.36
6	A	425	FDE	C1C-CHC	2.35	1.48	1.43
6	A	425	FDE	C4D-CHA	3.02	1.48	1.39
6	B	925	FDE	C1C-CHC	3.20	1.50	1.43
6	B	925	FDE	C4D-CHA	3.30	1.48	1.39
6	A	425	FDE	C3D-C2D	3.82	1.49	1.37
6	B	925	FDE	C3D-C2D	3.85	1.49	1.37
6	A	425	FDE	C2A-C3A	3.98	1.49	1.37
6	B	925	FDE	C2A-C3A	4.27	1.50	1.37

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	925	FDE	CBD-CAD-C3D	-6.51	100.87	112.53
6	A	425	FDE	CBD-CAD-C3D	-5.70	102.31	112.53
6	A	425	FDE	C1C-C2C-C3C	-4.56	102.68	106.42
6	B	925	FDE	C1C-C2C-C3C	-4.38	102.83	106.42

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1	CHD	C23-C22-C20	-4.27	109.71	114.75
6	B	925	FDE	CMF-C3C-C2C	-3.98	117.32	124.84
6	A	425	FDE	CAA-CBA-CGA	-3.26	106.77	112.75
6	A	425	FDE	CMF-C3C-C2C	-3.01	119.16	124.84
3	B	1	CHD	C6-C5-C4	-2.99	107.71	111.05
6	B	925	FDE	CAA-CBA-CGA	-2.93	107.38	112.75
3	A	3	CHD	C22-C20-C17	-2.80	104.35	110.24
3	B	4	CHD	O12-C12-C13	-2.77	106.63	111.11
3	B	1	CHD	C21-C20-C22	-2.39	106.36	110.35
3	A	3	CHD	C19-C10-C5	-2.23	106.32	110.25
6	B	925	FDE	CME-C3B-C2B	-2.14	120.79	124.84
3	B	4	CHD	C19-C10-C1	-2.08	104.71	108.20
3	B	4	CHD	C17-C13-C12	-2.04	115.87	117.68
3	A	2	CHD	C11-C9-C10	-2.04	111.67	113.79
3	B	1	CHD	C1-C10-C5	2.02	111.12	107.81
3	B	1	CHD	C11-C9-C8	2.03	113.61	110.73
4	A	1	GOL	O2-C2-C3	2.07	118.13	108.65
3	B	1	CHD	C14-C8-C7	2.07	114.61	111.74
3	B	1	CHD	C6-C7-C8	2.12	113.72	111.47
3	B	4	CHD	C6-C7-C8	2.17	113.77	111.47
3	B	1	CHD	C4-C5-C10	2.19	115.07	112.66
3	B	4	CHD	C13-C17-C20	2.25	122.24	119.50
3	B	4	CHD	C23-C22-C20	2.26	117.42	114.75
3	A	3	CHD	C21-C20-C17	2.30	116.79	112.96
3	B	4	CHD	C11-C12-C13	2.30	113.54	111.20
3	A	3	CHD	C11-C9-C8	2.33	114.05	110.73
3	A	2	CHD	C6-C7-C8	2.35	113.96	111.47
3	A	2	CHD	C1-C10-C5	2.36	111.69	107.81
4	A	1	GOL	C3-C2-C1	2.42	120.59	111.12
6	B	925	FDE	CMD-C2D-C3D	2.46	130.38	125.24
3	B	1	CHD	C15-C14-C13	2.47	106.05	103.60
3	A	2	CHD	C11-C9-C8	2.58	114.40	110.73
3	B	4	CHD	C1-C10-C5	2.67	112.19	107.81
3	B	1	CHD	C9-C10-C5	2.67	112.63	108.67
3	A	2	CHD	C15-C14-C13	2.74	106.32	103.60
3	A	2	CHD	C14-C8-C7	2.81	115.64	111.74
3	B	4	CHD	C11-C9-C8	2.94	114.91	110.73
3	A	3	CHD	C15-C14-C13	3.05	106.64	103.60
3	B	4	CHD	C15-C14-C13	3.07	106.65	103.60
3	A	3	CHD	C6-C7-C8	3.27	114.94	111.47
6	B	925	FDE	C2C-C1C-NC	3.34	114.45	110.36
6	A	425	FDE	C2C-C1C-NC	3.70	114.88	110.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	425	FDE	C2B-C1B-NB	5.94	116.52	108.64
6	B	925	FDE	C2B-C1B-NB	7.06	118.00	108.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2	CHD	2	0
6	A	425	FDE	2	0
3	B	1	CHD	3	0
6	B	925	FDE	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	359/359 (100%)	-0.29	4 (1%) 82 83	4, 14, 29, 43	0
1	B	359/359 (100%)	-0.22	8 (2%) 65 66	3, 15, 34, 46	0
All	All	718/718 (100%)	-0.25	12 (1%) 73 73	3, 15, 32, 46	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	600	THR	3.5
1	B	607	LEU	3.1
1	B	923	LEU	3.0
1	A	423	LEU	2.9
1	B	603	ILE	2.9
1	B	565	ARG	2.6
1	A	66	LYS	2.5
1	A	114	ARG	2.2
1	B	614	ARG	2.2
1	B	780	VAL	2.1
1	A	358	LYS	2.1
1	B	712	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CHD	B	4	29/29	0.77	0.23	4.97	27,30,47,50	0
4	GOL	A	1	6/6	0.89	0.27	3.28	21,25,26,27	0
3	CHD	A	3	29/29	0.88	0.15	2.64	17,21,35,38	0
3	CHD	A	2	29/29	0.82	0.23	1.78	37,40,60,61	0
5	IMD	B	924	5/5	0.89	0.15	1.33	28,28,29,29	0
3	CHD	B	1	29/29	0.82	0.23	1.17	37,40,57,59	0
6	FDE	B	925	39/39	0.94	0.15	0.97	18,23,28,30	0
5	IMD	A	424	5/5	0.93	0.11	0.09	15,15,17,17	0
6	FDE	A	425	39/39	0.98	0.09	-0.46	10,14,21,26	0
2	FES	B	501	4/4	0.99	0.06	-1.09	10,11,14,15	0
2	FES	A	501	4/4	0.99	0.07	-1.35	8,10,12,13	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.