



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:12 AM GMT

PDB ID : 3HHZ  
Title : Complex of the vesicular stomatitis virus nucleocapsid and the nucleocapsid-binding domain of the phosphoprotein  
Authors : Green, T.J.; Luo, M.  
Deposited on : 2009-05-18  
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1051 (3.60-3.40)
Clashscore	102246	1157 (3.60-3.40)
Ramachandran outliers	100387	1120 (3.60-3.40)
Sidechain outliers	100360	1121 (3.60-3.40)
RSRZ outliers	91569	1058 (3.60-3.40)
RNA backbone	2183	1050 (4.20-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	87	<div> <div>10%</div> <div>47%</div> <div>31%</div> <div>6%</div> <div>16%</div> </div>
1	B	87	<div> <div>31%</div> <div>62%</div> <div>20%</div> <div>•</div> <div>16%</div> </div>
1	C	87	<div> <div>7%</div> <div>57%</div> <div>23%</div> <div>••</div> <div>16%</div> </div>
1	D	87	<div> <div>34%</div> <div>71%</div> <div>9%</div> <div>•</div> <div>16%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	87	<div><div></div><div>16%</div><div>60%</div><div>21%</div><div>16%</div></div>
2	K	421	<div><div></div><div>3%</div><div>62%</div><div>30%</div><div>7%</div></div>
2	L	421	<div><div></div><div>2%</div><div>61%</div><div>33%</div><div>5%</div></div>
2	M	421	<div><div></div><div>3%</div><div>61%</div><div>32%</div><div>7%</div></div>
2	N	421	<div><div></div><div>5%</div><div>68%</div><div>28%</div><div></div></div>
2	O	421	<div><div></div><div>3%</div><div>62%</div><div>32%</div><div>6%</div></div>
3	R	45	<div><div></div><div>7%</div><div>31%</div><div>80%</div><div>49%</div><div>13%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 20415 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	73	Total	C	N	O	S	0	0	0
			576	368	100	106	2			
1	B	73	Total	C	N	O	S	0	0	0
			576	368	100	106	2			
1	C	73	Total	C	N	O	S	0	0	0
			576	368	100	106	2			
1	D	73	Total	C	N	O	S	0	0	0
			576	368	100	106	2			
1	E	73	Total	C	N	O	S	0	0	0
			576	368	100	106	2			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	179	GLY	-	EXPRESSION TAG	UNP P04880
A	180	SER	-	EXPRESSION TAG	UNP P04880
A	181	HIS	-	EXPRESSION TAG	UNP P04880
A	182	MET	-	EXPRESSION TAG	UNP P04880
B	179	GLY	-	EXPRESSION TAG	UNP P04880
B	180	SER	-	EXPRESSION TAG	UNP P04880
B	181	HIS	-	EXPRESSION TAG	UNP P04880
B	182	MET	-	EXPRESSION TAG	UNP P04880
C	179	GLY	-	EXPRESSION TAG	UNP P04880
C	180	SER	-	EXPRESSION TAG	UNP P04880
C	181	HIS	-	EXPRESSION TAG	UNP P04880
C	182	MET	-	EXPRESSION TAG	UNP P04880
D	179	GLY	-	EXPRESSION TAG	UNP P04880
D	180	SER	-	EXPRESSION TAG	UNP P04880
D	181	HIS	-	EXPRESSION TAG	UNP P04880
D	182	MET	-	EXPRESSION TAG	UNP P04880
E	179	GLY	-	EXPRESSION TAG	UNP P04880
E	180	SER	-	EXPRESSION TAG	UNP P04880
E	181	HIS	-	EXPRESSION TAG	UNP P04880

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	182	MET	-	EXPRESSION TAG	UNP P04880

- Molecule 2 is a protein called Nucleoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	K	421	Total	C	N	O	S	0	0	0
			3327	2118	558	633	18			
2	L	421	Total	C	N	O	S	0	0	0
			3327	2118	558	633	18			
2	M	421	Total	C	N	O	S	0	0	0
			3327	2118	558	633	18			
2	N	421	Total	C	N	O	S	0	0	0
			3327	2118	558	633	18			
2	O	421	Total	C	N	O	S	0	0	0
			3327	2118	558	633	18			


- Molecule 3 is a RNA chain called RNA (45-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	R	45	Total	C	N	O	P	0	0	0
			900	405	90	360	45			


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Phosphoprotein




#### • Molecule 1: Phosphoprotein




#### • Molecule 1: Phosphoprotein




#### • Molecule 1: Phosphoprotein




- Molecule 1: Phosphoprotein



- Molecule 2: Nucleoprotein




- Molecule 2: Nucleoprotein




- Molecule 2: Nucleoprotein







### • Molecule 2: Nucleoprotein




### • Molecule 2: Nucleoprotein







● Molecule 3: RNA (45-MER)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	166.57Å 235.25Å 96.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.50 29.91 – 3.50	Depositor EDS
% Data completeness (in resolution range)	98.5 (30.00-3.50) 98.5 (29.91-3.50)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.83 (at 3.47Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.254 , 0.323 0.248 , 0.302	Depositor DCC
$R_{free}$ test set	2443 reflections (5.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	93.0	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 110.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	1 of 47672 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	20415	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/586	0.57	0/784
1	B	0.39	0/586	0.52	0/784
1	C	0.38	0/586	0.56	0/784
1	D	0.37	0/586	0.55	0/784
1	E	0.39	0/586	0.55	0/784
2	K	0.44	0/3403	0.59	0/4607
2	L	0.43	0/3403	0.64	4/4607 (0.1%)
2	M	0.42	0/3403	0.58	0/4607
2	N	0.43	0/3403	0.65	4/4607 (0.1%)
2	O	0.42	0/3403	0.59	0/4607
3	R	1.01	0/989	1.63	12/1526 (0.8%)
All	All	0.47	0/20934	0.70	20/28481 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	M	0	2

There are no bond length outliers.

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	N	130	ASP	N-CA-C	13.65	147.86	111.00
2	N	131	ASP	N-CA-CB	-8.63	95.06	110.60
2	L	367	ASN	CB-CA-C	-7.90	94.61	110.40
3	R	36	U	O4'-C1'-N1	7.88	114.51	108.20
3	R	9	U	O4'-C1'-N1	7.56	114.25	108.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	M	112	ALA	Peptide
2	M	113	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	576	0	597	15	0
1	B	576	0	597	8	0
1	C	576	0	597	9	0
1	D	576	0	597	5	0
1	E	576	0	597	14	0
2	K	3327	0	3287	104	0
2	L	3327	0	3287	124	0
2	M	3327	0	3287	107	0
2	N	3327	0	3287	74	0
2	O	3327	0	3287	97	0
3	R	900	0	451	48	0
All	All	20415	0	19871	541	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 541 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:113:LEU:HD22	2:M:114:ASP:N	1.62	1.14
2:M:214:ARG:HA	2:M:217:THR:HG22	1.29	1.13
2:L:317:ARG:HD2	3:R:13:U:H2'	1.21	1.11
2:O:317:ARG:HD2	2:O:317:ARG:H	1.16	1.05
2:O:214:ARG:HA	2:O:217:THR:HG22	1.40	1.03

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	71/87 (82%)	51 (72%)	15 (21%)	5 (7%)	1	18
1	B	71/87 (82%)	55 (78%)	14 (20%)	2 (3%)	6	43
1	C	71/87 (82%)	60 (84%)	9 (13%)	2 (3%)	6	43
1	D	71/87 (82%)	52 (73%)	17 (24%)	2 (3%)	6	43
1	E	71/87 (82%)	57 (80%)	13 (18%)	1 (1%)	14	58
2	K	419/421 (100%)	338 (81%)	54 (13%)	27 (6%)	2	20
2	L	419/421 (100%)	326 (78%)	72 (17%)	21 (5%)	3	27
2	M	419/421 (100%)	330 (79%)	65 (16%)	24 (6%)	2	23
2	N	419/421 (100%)	333 (80%)	68 (16%)	18 (4%)	3	31
2	O	419/421 (100%)	315 (75%)	82 (20%)	22 (5%)	2	25
All	All	2450/2540 (96%)	1917 (78%)	409 (17%)	124 (5%)	2	26

5 of 124 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	K	118	PRO
2	K	206	LYS
2	K	215	TYR
2	K	357	PRO
2	L	43	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	64/77 (83%)	57 (89%)	7 (11%)	8	37
1	B	64/77 (83%)	59 (92%)	5 (8%)	16	53
1	C	64/77 (83%)	55 (86%)	9 (14%)	4	24
1	D	64/77 (83%)	60 (94%)	4 (6%)	22	63
1	E	64/77 (83%)	60 (94%)	4 (6%)	22	63
2	K	362/362 (100%)	313 (86%)	49 (14%)	5	26
2	L	362/362 (100%)	325 (90%)	37 (10%)	9	40
2	M	362/362 (100%)	315 (87%)	47 (13%)	5	27
2	N	362/362 (100%)	329 (91%)	33 (9%)	12	46
2	O	362/362 (100%)	320 (88%)	42 (12%)	7	33
All	All	2130/2195 (97%)	1893 (89%)	237 (11%)	8	35

5 of 237 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	321	ASP
2	M	158	ASP
2	O	292	VAL
2	L	327	LEU
2	M	66	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
2	M	266	GLN
2	N	63	ASN
2	O	318	GLN
2	M	347	GLN
2	M	367	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	R	44/45 (97%)	36 (81%)	8 (18%)

5 of 36 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	R	2	U
3	R	4	U
3	R	5	U
3	R	6	U
3	R	7	U

5 of 8 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	R	15	U
3	R	41	U
3	R	32	U
3	R	6	U
3	R	25	U

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	73/87 (83%)	0.65	9 (12%) 5 6	58, 60, 61, 62	0
1	B	73/87 (83%)	1.94	27 (36%) 0 0	57, 62, 64, 65	0
1	C	73/87 (83%)	0.68	6 (8%) 14 12	58, 60, 63, 65	0
1	D	73/87 (83%)	1.89	30 (41%) 0 0	59, 62, 64, 64	0
1	E	73/87 (83%)	1.08	14 (19%) 2 2	55, 61, 64, 65	0
2	K	421/421 (100%)	-0.18	11 (2%) 59 49	53, 63, 69, 73	0
2	L	421/421 (100%)	-0.13	7 (1%) 73 64	57, 65, 69, 72	0
2	M	421/421 (100%)	-0.17	12 (2%) 55 45	55, 64, 67, 69	0
2	N	421/421 (100%)	0.02	19 (4%) 37 29	57, 63, 67, 71	0
2	O	421/421 (100%)	-0.14	13 (3%) 52 43	57, 64, 67, 72	0
3	R	45/45 (100%)	2.75	36 (80%) 0 0	99, 108, 127, 139	0
All	All	2515/2585 (97%)	0.13	184 (7%) 18 14	53, 63, 69, 139	0

The worst 5 of 184 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	193	ALA	10.4
1	B	252	TYR	9.8
1	D	215	GLN	8.6
1	E	193	ALA	7.6
1	D	216	PRO	7.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.