



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:11 PM GMT

PDB ID : 4HIJ  
Title : Anti-Streptococcus pneumoniae 23F Fab 023.102 with bound L-rhamnose-(1-2)-alpha-D-galactose-(3-O)-phosphate-2-glycerol  
Authors : Bryson, S.; Risnes, L.; Damgupta, S.; Thomson, C.A.; Schrader, J.W.; Pai, E.F.  
Deposited on : 2012-10-11  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865



ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	B	304	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 023.102 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	209	Total 1614	C 1008	N 279	O 323	S 4	0	0	0
1	C	212	Total 1635	C 1021	N 283	O 327	S 4	0	0	0


- Molecule 2 is a protein called Fab 023.102 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	203	Total 1534	C 969	N 261	O 298	S 6	0	0	0
2	D	202	Total 1528	C 966	N 260	O 296	S 6	0	0	0

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	B	2	Total 23	C 13	O 10	0	0
3	D	2	Total 23	C 13	O 10	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total O P 4 3 1	0	0
4	D	1	Total O P 4 3 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 5 3 2	0	0
5	D	1	Total C O 5 3 2	0	0


- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	82	Total 82	O 82	0	0
6	B	64	Total 64	O 64	0	0
6	C	91	Total 91	O 91	0	0
6	D	82	Total 82	O 82	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab 023.102 light chain




- Molecule 1: Fab 023.102 light chain




- Molecule 2: Fab 023.102 heavy chain





● Molecule 2: Fab 023.102 heavy chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	143.87Å 66.49Å 118.21Å 90.00° 111.58° 90.00°	Depositor
Resolution (Å)	17.00 – 2.10 16.94 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (17.00-2.10) 80.7 (16.94-2.00)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.45 (at 2.00Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.249 , 0.262 0.251 , 0.263	Depositor DCC
$R_{free}$ test set	2671 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtrriage
Anisotropy	0.180	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 41.1	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	0 of 108059 reflections	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6694	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.05 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.5010e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, MBG, RAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.44	1/1651 (0.1%)	0.82	1/2246 (0.0%)
1	C	0.41	0/1672	0.78	0/2274
2	B	0.41	0/1569	0.80	2/2133 (0.1%)
2	D	0.41	0/1563	0.79	0/2125
All	All	0.42	1/6455 (0.0%)	0.80	3/8778 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	67	SER	CA-CB	-5.40	1.44	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	27	PHE	N-CA-C	-6.23	94.17	111.00
1	A	194	CYS	CA-CB-SG	5.95	124.70	114.00
2	B	99	ASN	N-CA-C	-5.55	96.02	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1614	0	1562	65	0
1	C	1635	0	1584	74	0
2	B	1534	0	1490	67	0
2	D	1528	0	1485	69	0
3	B	23	0	22	2	0
3	D	23	0	22	0	0
4	B	4	0	0	1	0
4	D	4	0	0	0	0
5	B	5	0	5	1	0
5	D	5	0	5	0	0
6	A	82	0	0	3	0
6	B	64	0	0	1	0
6	C	91	0	0	4	0
6	D	82	0	0	7	0
All	All	6694	0	6175	262	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 262 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:28:THR:HB	2:B:31:ARG:NE	1.64	1.12
1:A:24:ARG:HG3	1:A:24:ARG:HH11	0.96	1.08
2:B:28:THR:HB	2:B:31:ARG:HE	0.94	1.06
2:B:64:LYS:HD2	2:B:65:GLY:N	1.74	1.01
1:A:24:ARG:HG3	1:A:24:ARG:NH1	1.76	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/219 (94%)	198 (96%)	8 (4%)	1 (0%)	34	30
1	C	210/219 (96%)	203 (97%)	6 (3%)	1 (0%)	34	30
2	B	197/231 (85%)	188 (95%)	9 (5%)	0	100	100
2	D	196/231 (85%)	188 (96%)	7 (4%)	1 (0%)	34	30
All	All	810/900 (90%)	777 (96%)	30 (4%)	3 (0%)	39	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	138	ASN
1	A	29	VAL
2	D	149	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/184 (98%)	154 (86%)	26 (14%)	4	2
1	C	182/184 (99%)	165 (91%)	17 (9%)	11	7
2	B	170/196 (87%)	152 (89%)	18 (11%)	8	5
2	D	169/196 (86%)	149 (88%)	20 (12%)	6	3
All	All	701/760 (92%)	620 (88%)	81 (12%)	7	3

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	100	LYS
1	C	21	LEU
2	D	178	LEU
2	B	108	LEU
2	B	178	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	93	ASN
1	C	100	GLN
1	C	199	GLN
2	B	76	ASN
2	B	164	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	RAM	B	301	3	10,10,11	0.43	0	14,14,16	0.85	0
3	MBG	B	302	3,4	13,13,13	0.71	1 (7%)	18,18,18	1.27	4 (22%)
3	RAM	D	301	3	10,10,11	0.43	0	14,14,16	0.81	0
3	MBG	D	302	3,4	13,13,13	0.71	1 (7%)	18,18,18	0.91	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RAM	B	301	3	-	0/0/17/20	0/1/1/1
3	MBG	B	302	3,4	-	0/4/24/24	0/1/1/1
3	RAM	D	301	3	-	0/0/17/20	0/1/1/1

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MBG	D	302	3,4	-	0/4/24/24	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	MBG	O1-C1	2.17	1.43	1.40
3	D	302	MBG	O1-C1	2.21	1.44	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	MBG	C7-O1-C1	-2.97	108.44	113.29
3	B	302	MBG	C1-O5-C5	-2.42	109.05	113.75
3	D	302	MBG	C7-O1-C1	-2.34	109.46	113.29
3	B	302	MBG	O5-C5-C6	2.05	111.54	106.36
3	B	302	MBG	O1-C1-C2	2.28	110.90	108.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	RAM	1	0
3	B	302	MBG	1	0

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	PO4	B	303	3,5	0,3,4	0.00	-	0,3,6	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	B	304	4	4,4,5	1.76	1 (25%)	3,3,5	0.98	0
4	PO4	D	303	3,5	0,3,4	0.00	-	0,3,6	0.00	-
5	GOL	D	304	4	4,4,5	1.76	1 (25%)	3,3,5	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PO4	B	303	3,5	-	0/0/0/0	0/0/0/0
5	GOL	B	304	4	-	0/2/2/4	0/0/0/0
4	PO4	D	303	3,5	-	0/0/0/0	0/0/0/0
5	GOL	D	304	4	-	0/2/2/4	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	304	GOL	O1-C1	-3.44	1.23	1.42
5	B	304	GOL	O1-C1	-3.44	1.23	1.42

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	303	PO4	1	0
5	B	304	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	209/219 (95%)	0.31	13 (6%) 24 32	25, 40, 61, 66	0
1	C	212/219 (96%)	0.26	12 (5%) 27 35	25, 38, 58, 67	0
2	B	203/231 (87%)	0.21	10 (4%) 33 42	24, 40, 60, 66	0
2	D	202/231 (87%)	0.12	8 (3%) 42 51	26, 38, 51, 60	0
All	All	826/900 (91%)	0.23	43 (5%) 31 39	24, 39, 58, 67	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	135	THR	5.4
2	B	134	SER	4.3
1	C	2	GLY	4.3
1	A	56	THR	4.0
1	C	67	SER	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MBG	D	302	13/13	0.61	0.28	1.21	61,69,71,72	0
3	RAM	B	301	10/11	0.90	0.15	-0.21	59,61,62,64	0
3	RAM	D	301	10/11	0.92	0.14	-0.25	44,48,51,54	0
3	MBG	B	302	13/13	0.84	0.23	-	68,73,75,76	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	B	304	5/6	0.59	0.52	4.90	83,83,84,85	0
5	GOL	D	304	5/6	0.78	0.66	-	77,77,78,80	0
4	PO4	B	303	4/5	0.96	0.26	-	78,78,79,81	0
4	PO4	D	303	4/5	0.87	0.17	-	74,74,74,75	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.